

2003-04 Crop Year [August 1, 2003 to July 31, 2004]

NI Protein Content of 2003-04 Canada Western Amber Durum Wheat

Statistical Summary by Grade by Province

Final as at December 05, 2003

Canadian Grain Commission

Grain Research Laboratory (GRL)

NI Protein Content, %

Province	Mean	Std. Dev.
1 CWAD		
Manitoba	14.4	1.5
Saskatchewan	13.6	1.5
Alberta & Fort St. John area	13.7	1.4
Western Canada	13.6	1.5
2 CWAD		
Manitoba	13.9	0.9
Saskatchewan	13.3	1.9
Alberta & Fort St. John area	14.1	2.0
Western Canada	13.5	1.9
3 CWAD		
Manitoba	14.7	1.6
Saskatchewan	13.5	1.9
Alberta & Fort St. John area	14.4	1.6
Western Canada	13.8	1.9
All Grades		
Manitoba	14.5	1.4
Saskatchewan	13.6	1.6
Alberta & Fort St. John area	13.8	1.5
Western Canada	13.6	1.6

Notes:

1. NI Protein content is determined by near infrared measurements calibrated against the Combustion Nitrogen Analysis reference method and is expressed on an N x 5.7, 13.5% moisture basis.
2. Results are unlikely to represent samples harvested within two weeks of the date of this report.
3. Samples were supplied by producers and grain companies.
4. A total of 2006 samples were tested.