

2004-05 Crop Year [August 1, 2004 to July 31, 2005]

NI Protein Content of 2004-05 Canada Western Amber Durum Wheat

Statistical Summary by Grade by Province

Final as at November 25, 2004

Canadian Grain Commission

Grain Research Laboratory (GRL)

NI Protein Content, %

Province	Mean	Std. Dev.
1 CWAD		
Saskatchewan	12.8	1.6
Alberta & Fort St. John area	13.6	1.5
Western Canada	13.1	1.6
2 CWAD		
Manitoba	14.4	0.7
Saskatchewan	12.1	1.5
Alberta & Fort St. John area	13.0	1.4
Western Canada	12.3	1.5
3 CWAD		
Manitoba	12.1	0.0
Saskatchewan	12.1	1.6
Alberta & Fort St. John area	12.4	1.3
Western Canada	12.2	1.5
4 CWAD		
Manitoba	12.9	0.5
Saskatchewan	12.3	1.5
Alberta & Fort St. John area	12.5	1.5
Western Canada	12.3	1.5
All Grades		
Manitoba	13.2	0.9
Saskatchewan	12.2	1.5
Alberta & Fort St. John area	12.8	1.5
Western Canada	12.4	1.5

Notes:

1. NI Protein content is determined by near infrared measurements calibrated against the Combustion Nitrogen Analysis reference method and is expressed on an N x 5.7, 13.5% moisture basis.
2. Results are unlikely to represent samples harvested within two weeks of the date of this report.
3. Samples were supplied by producers and grain companies.
4. A total of 1035 samples were tested.