



Regulatory Note

REG2003-03

Fenbuconazole

The fungicide active ingredient fenbuconazole and associated end-use product Indar 75WSP Fungicide for the control of blossom blight and fruit brown rot on Crop Group 12 Stone Fruit—apricot, sweet cherry, tart cherry, nectarine, peach, plum, chickasaw plum, damson plum, Japanese plum, and fresh prune; and black knot on tart cherry, chickasaw plum, damson plum, Japanese plum and fresh prune, have been granted temporary registration under Section 17 of the Pest Control Products Regulations (PCPR).

This Regulatory Note provides a summary of material reviewed and the rationale for the regulatory decision for these products.

(publié aussi en français)

April 28, 2003

This document is published by the Alternative Strategies and Regulatory Affairs Division, Pest Management Regulatory Agency. For further information, please contact:

Publications Coordinator
Pest Management Regulatory Agency
Health Canada
2720 Riverside Drive
A.L. 6605C
Ottawa, Ontario
K1A 0K9

Internet: pmra_publications@hc-sc.gc.ca
www.hc-sc.gc.ca/pmra-arla/

Information Service:
1-800-267-6315 or (613) 736-3799
Facsimile: (613) 736-3798



ISBN: 0-662-34032-9

Catalogue number: H113-7/2003-3E-IN

**© Her Majesty the Queen in Right of Canada, represented by the Minister of Public Works and Government Services
Canada 2003**

All rights reserved. No part of this information (publication or product) may be reproduced or transmitted in any form or by any means, electronic, mechanical photocopying, recording or otherwise, or stored in a retrieval system, without prior written permission of the Minister of Public Works and Government Services Canada, Ottawa, Ontario K1A 0S5.

Foreword

Health Canada's Pest Management Regulatory Agency (PMRA) has issued temporary registration for Indar Technical and the associated end-use product, Indar 75WSP, manufactured by Dow AgroSciences Canada Inc., for the control of blossom blight and fruit brown rot on Crop Group 12 Stone Fruit—apricot, sweet cherry, tart cherry, nectarine, peach, plum, chickasaw plum, damson plum, Japanese plum and fresh prune; and black knot on tart cherry, nectarine, peach, plum, chickasaw plum, damson plum, Japanese plum and fresh prune.

The following maximum residue levels (MRLs) will be recommended for promulgation in Table II, Division 15, of the *Food and Drugs Act* and Regulations:

apricot (0.3 ppm); sweet and tart cherry (0.8 ppm); peach (0.5 ppm); nectarine (0.5 ppm); plum, chickasaw plum, damson plum, Japanese plum, plumcot (0.1 ppm); fresh prune (0.1 ppm); dry prune (0.5 ppm).

Methods for analysing fenbuconazole in environmental media are available to research and monitoring agencies upon request to the PMRA.

Dow AgroSciences Canada Inc. will be carrying out additional environmental studies as a condition of this temporary registration. Following the review of this information, the PMRA will publish a proposed registration decision document (PRDD) and request comments from interested parties before proceeding with a final regulatory decision.

Table of Contents

1.0	The active substance, its properties and uses	1
1.1	Identity of the active substance and impurities	1
1.2	Physical and chemical properties of active substances and end-use product(s)	2
1.3	Details of uses	4
2.0	Methods of analysis	4
2.1	Method for analysis of the active substance as manufactured	4
2.2	Method for formulation analysis	5
2.3	Methods for residue analysis	5
2.3.1	Multi-residue methods for residue analysis	5
2.3.2	Methods for residue analysis of plants and plant products	5
2.3.3	Methods for residue analysis of food of animal origin	6
3.0	Impact on human and animal health	6
3.1	Integrated toxicological summary	6
3.2	Determination of acceptable daily intake (ADI)	8
3.3	Acute reference dose (ARfD)	9
3.4	Toxicology endpoint selection for occupational and bystander risk assessment	9
3.5	Impact on human and animal health arising from exposure to the active substance or to its impurities	10
3.5.1	Operator exposure assessment	10
3.5.2	Bystanders	11
3.5.3	Workers	12
3.5.4	Consumers	13
4.0	Residues	13
4.1	Food residues exposure assessment	13
5.0	Fate and behaviour in the environment	16
5.1	Physical and chemical properties relevant to the environment	16
5.2	Abiotic transformation	17
5.3	Biotransformation	17
5.4	Mobility	19
5.5	Dissipation and accumulation under field conditions	20
5.6	Bioaccumulation	21
5.7	Summary of fate and behaviour in the terrestrial environment	21
5.8	Summary of fate and behaviour in the aquatic environment	23
5.9	Expected environmental concentrations	23
5.9.1	Soil	24
5.9.2	Aquatic systems	24
5.9.3	Vegetation and other food sources	25


6.0	Effects on non-target species	28
6.1	Effects on terrestrial organisms	28
6.2	Effects on aquatic organisms	29
6.3	Effects on biological methods of sewage treatment	30
6.4	Risk characterization	30
6.4.1	Environmental behaviour	30
6.4.2	Terrestrial organisms	31
6.4.3	Aquatic organisms	36
6.5	Risk mitigation	40
7.0	Efficacy	41
7.1	Effectiveness	41
7.1.1	Intended use	41
7.1.2	Mode of action	42
7.1.3	Nature of pest problem	42
7.1.4	Effectiveness against pest	43
7.2	Phytotoxicity to target plants (including different cultivars), or to target plant products (OECD 7.4)	44
7.3	Observations on undesirable or unintended side effects, e.g., on beneficial and other non-target organisms, on succeeding crops, other plants, or parts of treated plants used for propagating purposes (e.g., seeds, cuttings, runners) (OECD 7.5)	44
7.3.1	Impact on succeeding crops (OECD 7.5.1)	44
7.3.2	Impact on adjacent crops (OECD 7.5.2)	45
7.4	Economics	45
7.5	Sustainability	45
7.5.1	Survey of alternatives	45
7.5.2	Compatibility with current management practices including IPM	47
7.5.3	Contribution to risk reduction	48
7.5.4	Information on the occurrence or possible occurrence of the development of resistance	48
7.6	Conclusions	49
7.6.1	Summary	49
8.0	Toxic Substances Management Policy considerations	51
9.0	Regulatory decision	51
	List of abbreviations	53
	References	55
Appendix I	Summary table of toxicology studies for fenbuconazole	64
Appendix II	Food residue chemistry overview of metabolism studies and risk assessment .	77

Appendix III	Food residue chemistry integrated summary table	82
Appendix IV	Environmental assessment	84

1.0 The active substance, its properties and uses

1.1 Identity of the active substance and impurities

Table 1.1.1 Identity of the active substance and preparation containing it

Active substance	Fenbuconazole
Function	Fungicide
Chemical name <ul style="list-style-type: none">International Union of Pure and Applied Chemistry (IUPAC)Chemical Abstracts Service (CAS)	(RS)-4-(4-chlorophenyl)-2-phenyl-2-(1 <i>H</i> -1,2,4-triazole-1-ylmethyl)butyronitrile The active substance is a racemic mixture of R and S isomers. The IUPAC name is from ISO 1750:1981/DAM 2 document. α -[2-(4-chlorophenyl)ethyl]- α -phenyl-1 <i>H</i> -1,2,4-triazole-1-propanenitrile
CAS number	119611-00-6, racemate. This number supersedes 114369-43-6, but both are correct.
Molecular formula	C ₁₉ H ₁₇ ClN ₄
Molecular weight	336.83
Structural formula	 <p>*denotes chiral carbon</p>
Nominal purity of active	98.3%, nominal (label and SPSF) (limits 94.0%, 99.5%)
Identity of relevant impurities of toxicological, environmental, or other significance	The technical grade [fenbuconazole] does not contain any impurities or microcontaminants known to be Toxic Substances Management Policy (TSMP) Track-1 substances.

1.2 Physical and chemical properties of active substances and end-use product(s)

Table 1.2.1 Technical product: Fenbuconazole

Property	Result	Comment																
Colour and physical state	Off-white to white powder																	
Odour	Faintly sulphur-like																	
Melting point or range	126.5–127.0°C																	
Boiling point or range	Not required																	
Density	0.50 g/mL, bulk density																	
Vapour pressure at 25°C (PAI)	0.37×10^{-7} mm Hg (4.9×10^{-6} Pa)	The active ingredient will be non-volatile under field conditions.																
Henry's Law constant at 20°C	4.3×10^{-9} atm·m ³ /mol or 5.57×10^6 (1/H)	The active ingredient will be non-volatile from moist soils or water.																
Ultraviolet (UV)—visible spectrum	<table border="0"> <tr> <td><u>λ max (nm)</u></td> <td><u>ϵ (L·mol⁻¹·cm⁻¹)</u></td> </tr> <tr> <td>196</td> <td>53,000</td> </tr> <tr> <td>262</td> <td>750</td> </tr> <tr> <td>268</td> <td>740</td> </tr> <tr> <td>275</td> <td>480</td> </tr> </table>	<u>λ max (nm)</u>	<u>ϵ (L·mol⁻¹·cm⁻¹)</u>	196	53,000	262	750	268	740	275	480	The active ingredient has a low potential for ultraviolet light-induced phototransformation under normal environmental conditions.						
<u>λ max (nm)</u>	<u>ϵ (L·mol⁻¹·cm⁻¹)</u>																	
196	53,000																	
262	750																	
268	740																	
275	480																	
Solubility in water at 22°C	<table border="0"> <tr> <td><u>pH</u></td> <td><u>Solubility (mg/L)</u></td> </tr> <tr> <td>not provided</td> <td>3.8</td> </tr> </table>	<u>pH</u>	<u>Solubility (mg/L)</u>	not provided	3.8	The solubility of the active ingredient in water is classified as low.												
<u>pH</u>	<u>Solubility (mg/L)</u>																	
not provided	3.8																	
Solubility (g/L) in organic solvents at 25°C	<table border="0"> <tr> <td><u>Solvent</u></td> <td><u>g/L</u></td> </tr> <tr> <td>acetonitrile</td> <td>231</td> </tr> <tr> <td>aromatic 200</td> <td>77</td> </tr> <tr> <td>cyclohexanone</td> <td>445</td> </tr> <tr> <td>ethyl acetate</td> <td>159</td> </tr> <tr> <td>ethyl alcohol</td> <td>39</td> </tr> <tr> <td>heptane</td> <td>01</td> </tr> <tr> <td>1-octanol</td> <td>13</td> </tr> </table>	<u>Solvent</u>	<u>g/L</u>	acetonitrile	231	aromatic 200	77	cyclohexanone	445	ethyl acetate	159	ethyl alcohol	39	heptane	01	1-octanol	13	
<u>Solvent</u>	<u>g/L</u>																	
acetonitrile	231																	
aromatic 200	77																	
cyclohexanone	445																	
ethyl acetate	159																	
ethyl alcohol	39																	
heptane	01																	
1-octanol	13																	

Property	Result	Comment
<i>n</i> -Octanol–water partition coefficient (K_{ow}) (PAI, 99.5%)	1700 ± 300 log K_{ow} = 3.22 ± 0.08	There is potential for the active ingredient to bioaccumulate.
Dissociation constant (pK_a)	Not expected to dissociate in water.	
Stability (temperature, metal)	Stable to temperatures of 220°C or higher Stable when exposed to stainless steel and aluminum metal and to potassium ions and iron (II) and iron (III) oxide	

Table 1.2.2 End-use product: Indar 75WSP Agricultural Fungicide

Property	Result
Colour	Off-white
Odour	Musty, pungent
Physical state	Powder
Formulation type	Wettable powder
Guarantee	Fenbuconazole: 75.0%, nominal (label and SPSF) (limits 72.8%, 77.3%)
Formulants	The product does not contain any EPA List 1 formulants or formulants known to be TSMP Track-1 substances.
Container material and description	Water soluble pouch with an outer waterproof bag
Bulk density	Loose 0.16 g/mL Packed 0.20 g/mL
pH of 1% dispersion in water	7.6
Oxidizing or reducing action	None of the components is an oxidizing or reducing agent.
Storage stability	No significant change in level of active ingredient or in properties after 24 months' storage in commercial packaging (water soluble PVA bag), at 25°C

Property	Result
Explosibility	RH-7592 technical may undergo a dust explosion. K_{st} for technical RH-7592 is 265 bar-m/sec and minimum O_2 required for combustion is 12%. The RH-7592 75WP explosion hazard is expected to be no greater than the technical.

1.3 Details of uses

Indar 75WSP is proposed for application to stone fruits (Use Site Category (USC) 14) to control blossom blight and fruit brown rot of apricot, sweet cherry and tart cherry, nectarines, peaches and plums, as well as leaf spot of sweet cherry and tart cherry, scab of peach, and black knot of sour cherry and plum. The proposed use rate is 140 g product/500 L water/ha applied by ground (usually airblast) sprayer, with up to two sprays at early red bud stage through bloom stage, and a third cover spray pre-pick if needed. In addition it can be applied to fruit up to the time of harvest but not after harvest, and may be applied to foliage post-harvest for leaf spot only. Spray water volume varies considerably with size of trees and is calculated case-by-case to give optimal coverage while maintaining product rate per hectare.

Indar 75WSP should be first applied prior to infection. It is proposed at spray intervals of 7–14 days, depending on disease pressure, and can be applied up to the day of harvest. The applicant has indicated that there can be up to eight spray dates per season; however, the typical number of applications is two per disease.

2.0 Methods of analysis

2.1 Method for analysis of the active substance as manufactured

Product	Analyte	Method ID	Method Type	Linear range (%)	RSD (%)	Method acceptability
Technical	Fenbuconazole	91-100-01	Cap GC	55–105	0.4	Y
Technical	Major impurities	91-100-01	Cap GC	0.1–1.0	2.5–10	Y

2.2 Method for formulation analysis

Product	Analyte	Method ID	Method	Mean recovery (%)	RSD (%)	Method acceptability
Indar 75 WSP	Fenbuconazole	92-119-01	GC	100.1% (n = 2)	0.48% (n = 6)	Acceptable

2.3 Methods for residue analysis

2.3.1 Multi-residue methods for residue analysis

Existing multi-residue methods of analysis that are currently in common usage were not found to be suitable for the determination of fenbuconazole in stone fruit (apricot, cherry, nectarine, peach, plum, and prune).

2.3.2 Methods for residue analysis of plants and plant products

The residue of concern (ROC) was defined as fenbuconazole (RH-7592) and its lactone metabolites (RH-9129 and RH-9130). The petitioner is proposing a single, specific, analytical method (TR 34-90-47R) for supervised residue trials and enforcement of maximum residue limit (MRL). This method can individually determine the residue of RH-7592, RH-9129, and RH-9130 in stone fruit matrices.

The residues of fenbuconazole and its lactone metabolites are extracted in stone fruit using methanol. The extract is separated by filtration and then partitioned with 9.1% sodium chloride and methylene chloride. The eluate is collected, evaporated to dryness, and residues are reconstituted in toluene:acetone (100:10, v:v). Further purification is achieved using silica gel and florisil column chromatography with toluene:acetone (100:30, v:v) as the eluent. The residue is collected, evaporated to dryness, redissolved in toluene:methanol (100:3, v:v) and then analysed by gas liquid chromatography using capillary column and thermionic specific detector optimized for nitrogen selectivity. The limit of detection (LOD) for fenbuconazole (RH-7592) and the metabolites (RH-9129 and RH-9130) was established at 0.01 ppm in stone fruit. The limit of quantification (LOQ) for fenbuconazole (RH-7592) and the lactone metabolites (RH-9129 and RH-9130) was established at 0.05 ppm in stone fruit.

The method validation recoveries were adequate in stone fruit matrices when samples were spiked with a mixture of analytes at levels of 0.01 to 4.0 ppm. The recoveries were $89 \pm 6\%$ for RH-7592, $90 \pm 6\%$ for RH-9130 and $82 \pm 10\%$ for RH-9129. The method/detector response for all analytes was linear over the range of 0.05–1.0 ppm with a correlation coefficient of >0.999 . The method applied external standards as markers for retention time, response, and calibration. The chromatographic peaks were well defined

and symmetrical with no apparent carryover to the subsequent chromatograms in the area of analytical interest for both control and spiked samples.

The inter-laboratory validation (ILV) did support the reliability and reproducibility of the analytical method for the determination of the residues of the RH-7592, RH-9129, and RH-9130 in stone fruit matrices. When samples were spiked with a mixture of analyte at 1.0 ppm and 2.0 ppm, recoveries were $97 \pm 5\%$ for RH-7592, $94 \pm 5\%$ for RH-9130 and $90 \pm 6\%$ for RH-9129 in stone fruit matrices.

2.3.3 Methods for residue analysis of food of animal origin

Neither poultry nor livestock feed items are associated with the use of fenbuconazole on stone fruit. Therefore, no analytical method is needed for animal commodities.

3.0 Impact on human and animal health

3.1 Integrated toxicological summary

A detailed review of the toxicology database available for the new fungicide, fenbuconazole, has been completed. Data submitted were complete and well presented, and included the full battery of studies currently required for registration purposes. The submitted studies were conducted in conformance with currently acceptable international testing protocols.

Following oral dosing, ^{14}C -RH-7592 was rapidly absorbed, distributed and excreted. The feces was the major route of excretion accounting for ~76% to 94% of the administered dose. Recovery from urine ranged from ~5% to 14%. The majority of the radioactivity was excreted within 24 to 48 hours post-dosing. Biliary excretion data indicated that systemic absorption of RH-7592 was high for all dosing groups. Tissue distribution and bioaccumulation were minimal, with <1% of the administered dose recovered in tissues at 96 hours. There was no sex- or dose-related differences in absorption, distribution, or elimination.

There were many metabolites in excreta indicating extensive metabolic breakdown. All major metabolites were derived from enzymatic oxidations on either the benzylic carbon alpha to the chlorophenyl ring or the 3- or 4- position of the phenyl ring. Subsequent non-enzymatic cyclization of the newly formed benzylic alcohol with the adjacent nitrile group, followed by hydrolysis led to the iminolactone/lactone family of metabolites. Conjugation of the OH groups of the alcohol and phenols led to more metabolites, as did combinations of the above-mentioned reactions. A minor metabolic pathway was the cleavage of RH-7592 to yield triazole and RS-5922. There was no significant difference in the total metabolite profile for male and female rats, although some metabolites were present in greater quantities in males, and vice versa. There was a dose-related difference in metabolism, with a higher amount of unmetabolized parent compound in the feces of

the high-dose group, compared to the low-dose and repeated-dose groups, indicating that saturation of the metabolic pathway may be occurring at the high dose.

Acute dosing revealed that technical fenbuconazole was of low toxicity by the oral, dermal, and inhalation routes. It was non-irritating to the skin, minimally irritating when instilled into the eyes, and was not a skin sensitizer (Buehler method). The Indar 75 WP formulation, containing 77.5% fenbuconazole, was of low toxicity by the oral, dermal, and inhalation routes. It was minimally irritating to the skin and eyes and was not a dermal sensitizer (Buehler method).

Short-term (28 days) repeated dermal dosing in rats with technical fenbuconazole did not result in any treatment-related systemic or dermal effects up to and including the highest dose level tested (limit dose) of 1000 mg/kg bw/day.

The target organ for all species tested, after both short-term and long-term exposure to technical fenbuconazole, was the liver. Liver weights were increased at dose levels of ~13 mg/kg bw/day and higher for dogs, and ~25 mg/kg bw/day and higher for mice and rats, with corresponding histopathological changes (hepatocyte hypertrophy and vacuolization in all 3 species; hepatocyte hyperplasia and necrosis in mice only). In addition, male rats exhibited hepatocyte vacuolation at ~5 mg/kg bw/day. Increased liver enzyme activity was observed for mice (SGOT, SGPT) and dogs (alkaline phosphatase, SGPT, SGOT). The thyroid was also a target organ for rats, with increased thyroid weight, follicular cell hypertrophy and focal cystic hyperplasia observed at doses of ~30 mg/kg bw/day and higher. In addition, plasma thyroxine was decreased and thyroid stimulating hormone (TSH) was increased at a dose of 62.07 mg/kg bw/day. Thyroid weight and adrenal weight were increased for dogs at ~45 mg/kg bw/day, but without corresponding histopathological findings.

There was evidence of oncogenic/carcinogenic potential of fenbuconazole in rodents. For rats, there was an increased occurrence of thyroid follicular cell benign tumours and combined thyroid follicular cell benign and malignant tumours in males (28.87 mg/kg bw/day). For mice, this was based on the occurrence of an increased trend for malignant liver tumours in males (85.26 mg/kg bw/day) and an increase in benign liver tumours and combined benign and malignant liver tumours in females (208.84 mg/kg bw/day). The proposed mechanism of action for the thyroid tumours in rats was scientifically supported by sound mechanistic data, i.e., prolonged stimulation of the thyroid by TSH leads to chronic follicular hypertrophy/hyperplasia, progressing to follicular neoplasia of the thyroid. However, data submitted for the proposed mechanistic pathway for liver tumours in mice did not provide a convincing hypothesis. The in vitro and in vivo mutagenicity assays conducted yielded negative results for genotoxic potential. In addition, 2 metabolites of fenbuconazole, i.e., RH-99129 and RH-99130, were negative for genotoxicity when tested according to the *Salmonella*/Ames Assay. It is therefore recommended that for the purpose of risk characterization a low-dose extrapolation model be applied for human risk (Q_1^*). This decision is based on the induction of liver

carcinomas in male mice. The Q_1^* for fenbuconazole is 1.54×10^{-2} (mg/kg bw/day)⁻¹ in human equivalents.

Reproductive performance, maternal toxicity, and offspring toxicity in rats were seen only at the high dose of 66.4 mg/kg bw/day. Reproductive findings were decreased number of dams delivering; decreased litter size; decreased number and percentage of dams with liveborn pups; decreased gestation index; increased number and percentage of dams with stillborn pups; and increased number of litters with no viable offspring (P₁ only). Offspring effects were decreased pup viability, and lower pup body weight on days 0, 4, 7, 14, and 21 postpartum for the F₁ pups. Parental findings were treatment-related mortality in females only; decreased mean body weight, body weight gain, and food consumption; increased liver weights; hepatocyte vacuolation and hypertrophy; increased thyroid weight in males only; follicular cell hypertrophy in both sexes; increased adrenal weight in females only; and hypertrophy of the zona glomerulosa in both sexes. Increased susceptibility of pups was not demonstrated in this study.

Fenbuconazole was not teratogenic to rat or rabbit fetuses at dose levels up to and including 150 mg/kg bw/day (rats) and 30 mg/kg bw/day (rabbits). A meaningful evaluation of soft tissue, visceral or skeletal effects could not be conducted for rabbits in the high dose (60 mg/kg bw/day) group (maternally toxic dose level) since only one litter was produced. Developmental toxicity was noted for rats at 75 and 150 mg/kg bw/day (maternally toxic doses), manifest as a decrease in the number of live fetuses/litter and an increase in post-implantation loss. In addition, there was an increase in early and late resorptions in the 150 mg/kg bw/day group. Fetotoxicity was evident as an increased incidence of incompletely ossified or unossified sternebrae noted for rat fetuses in the 75 and 150 mg/kg bw/day groups, and an increased incidence of rudimentary 14th ribs and partially or unossified pubes in the 150 mg/kg bw/day group. These minor variations were not considered to be adverse, toxicologically significant findings. For rabbits, embryo-fetal mortality (abortions, total resorptions of litters) was observed at 60 mg/kg bw/day (maternally toxic dose). Maternal findings were observed in rabbits at 30 and 60 mg/kg bw/day (decreased food intake, clinical findings at 30 and 60 mg/kg bw/day; loss of body weight, increased mortality at 60 mg/kg bw/day). Maternal toxicity was seen in rats at 75 and 150 mg/kg bw/day (clinical signs, lower body weight gain and lower gravid uterus weight). There was no evidence for increased susceptibility of rat or rabbit fetuses following in utero exposure to fenbuconazole.

3.2 Determination of acceptable daily intake (ADI)

The lowest NOEL was 10 ppm, equal to 1.28/1.59 mg/kg bw/day, established in the 78-week oncogenicity feeding study in mice, based on increased liver weight and liver histopathology at higher dose levels. For the calculation of the ADI, an uncertainty factor (UF) of 100 is proposed. This provides a margin of safety (MOS) of 500× for reproductive toxicity.

The ADI proposed is calculated according to the following formula:

$$\text{ADI} = \frac{\text{NOAEL}}{\text{SF}} = \frac{1.28 \text{ mg/kg bw/day}}{100}$$

$$= 0.0128 \text{ mg/kg bw/day of fenbuconazole.}$$

3.3 Acute reference dose (ARfD)

For an acute reference dose for females 13+ years of age, the studies considered most appropriate in the submitted toxicological data base are the rat and rabbit teratology studies. The dose and endpoint selected for risk assessment is 30 mg/kg bw/day (decrease in live fetuses per litter, and an increase in post-implantation loss at 75 and 150 mg/kg bw/day for rats; increase in abortions and post-implantation loss at 60 mg/kg bw/day for rabbits). For the calculation of the ARfD, an uncertainty factor (UF) of 300 is proposed. This is based on the standard uncertainty factor of 100 with an additional 3-fold uncertainty factor due to the severity of the toxicological endpoint, i.e., increased post-implantation loss and a decrease in the number of live fetuses per litter.

The ARfD proposed is calculated according to the following formula:

$$\text{ARfD} = \frac{\text{NOAEL}}{\text{SF}} = \frac{30 \text{ mg/kg bw/day}}{300}$$

$$= 0.10 \text{ mg/kg bw/day of fenbuconazole.}$$

3.4 Toxicology endpoint selection for occupational and bystander risk assessment

For mixers, loaders and applicators treating stone fruits, the expected duration of exposure, which is predominantly via the dermal route, is intermittent over a 4-month period. For re-entry workers, the expected duration of exposure is intermittent to continuous, intermediate-term (i.e., <6 months). Re-entry exposure would be dermal.

Adequate dermal toxicity data of sufficient duration for the requested orchard use were not available for INDAR 75WSP. Oral studies of sufficient duration have been used for the assessment of risk. After 3 months of dietary exposure to fenbuconazole, a NOAEL of 1.3 mg/kg bw/day (rat) was determined, based on hepatocyte vacuolation at the next dose of 5.1 mg/kg bw/day. The NOAEL established in this study was considered the most appropriate toxicological endpoint for use in the occupational risk assessment.

Maternal toxicity (NOAEL: 10 mg/kg bw/day; rabbit), reproductive toxicity (NOAEL: 6.4 mg/kg bw/day; rat) and developmental toxicity (NOAEL: 30 mg/kg bw/day; rabbit, rat) concerns have been adequately addressed with the NOAEL of 1.3 mg/kg bw/day from the 3-month dietary study (rat).

As fenbuconazole did not demonstrate toxicity concerns on genotoxicity, teratogenicity, or neurotoxicity, the target margin of exposure is 100 for inter- and intra-species variation. There was no evidence of increased sensitivity of the young.

There is evidence of carcinogenicity in animals following long-term exposure to fenbuconazole. It is recommended that, for the purpose of risk characterization, a low-dose extrapolation model be applied for human risk (Q_1^*). This decision is based on the induction of liver carcinomas in male mice. The Q_1^* for fenbuconazole is 1.54×10^{-2} (mg/kg bw/day)⁻¹ in human equivalents.

3.5 Impact on human and animal health arising from exposure to the active substance or to its impurities

Indar 75 WSP would be applied at a rate of 105 g a.i./ha by airblast equipment, up to eight times per season. A mixer/loader/applicator (farmer) would typically treat 30 ha of stone fruit per day. At each application, applicators would handle 3.15 kg a.i. There is potential for intermittent exposure to farmers over a 4-month period.

Workers re-entering treated areas will have potential for post-application exposure. Application times coincide with pruning, hand thinning, harvesting, and other re-entry activities in orchards. There is potential for intermittent to continuous, intermediate-term dermal exposure for re-entry workers.

Dermal absorption

The dermal absorption of ¹⁴C RH-7592 (fenbuconazole) was determined in male rats at doses of 0.002, 0.027, and 1.98 mg/cm². Exposure durations were 0.5, 1, 2, 4, 10, and 24 hours, four rats per dose duration. In addition, one group of animals in the 0.002 mg/cm² dose groups were exposed for 10 hours and then sacrificed after 168 hours (7 days). Recovery at all doses ranged from 73 to 111% of the administered dose.

Absorption was low and increased with duration of exposure. The maximum dermal absorption occurred at the low dose (0.002 mg/cm²). Due to the limitations of the study, particularly the uncertainties regarding the applicability of the study formulations to the proposed product formulation, the highest dermal absorption value from the 0.002 mg/cm² dose group (i.e., 12% at 24 hours) was considered to be the most appropriate value to use for exposure assessments. As the study design did not permit analysis of the fate of skin-bound residues, this dermal absorption value includes residues retained at the skin site (approximately 4%).

3.5.1 Operator exposure assessment

Mixer/loader/applicator exposure was estimated using the Pesticide Handlers' Exposure Database Version 1.1 (PHED 1.1). PHED is a compilation of generic mixer/loader/applicator passive dosimetry data with associated software which facilitates the generation of scenario-specific exposure estimates. With a few exceptions as noted,

the PHED estimates meet criteria for data quality, specificity, and quantity outlined under the North American Free Trade Agreement Technical Working Group on Pesticides. Exposure via the inhalation route was a minor component of overall exposure and was added to the dermal deposition estimates coupled with a default dermal absorption value of 12%.

To estimate exposure for each use scenario, appropriate subsets of A- and B-grade datasets were created from the mixer/loader/applicator database files of PHED. All data were normalized for kg of active ingredient handled. Exposure estimates are presented on the basis of the best-fit measure of central tendency, i.e., summing the measure of central tendency for each body part which is most appropriate to the distribution of data for that body part. Estimates were derived for individuals wearing one layer of clothing and gloves during mixing, loading, and application.

The following exposure estimates and margins of exposure and cancer risk were derived for mixer/loader/applicators:

Table 3.5.1 Mixer/loader/applicator exposure

Occupational scenario	Exposure ¹ (mg/kg bw/day)	Margin of exposure ²	LADD ³ (mg/kg bw/day)	Cancer risk ⁴
Mixer/loader + applicator exposure ⁵				
Stone Fruit—airblast	0.00342	380	4.00 e-05	6.16 e-07

¹ Based on a 70 kg operator, typical North American use patterns of 30 ha/day for stone fruit, and 12% default dermal absorption

² Based on NOAEL 1.3 mg/kg bw/day from a 3-month oral rat study

³ Lifetime Average Daily Dose, calculated assuming 8 days of exposure per year, a life expectancy of 75 years, and a working lifetime of 40 years.

⁴ Based on Q₁* of 1.54 × 10⁻² (mg/kg bw/day)⁻¹

⁵ Individuals wearing one layer of clothing and gloves (exception: groundboom applicator not wearing gloves)

These margins of exposure and cancer risk are acceptable.

3.5.2 Bystanders

Given that the application is restricted to agricultural areas, and that the product would be applied using ground equipment only, exposure and risk to bystanders is expected to be negligible. In addition, a statement will be added on the label that the product is not for use on residential properties.

3.5.3 Workers

Re-entry activities include pruning, thinning, hand-harvesting of fruit, and other pesticide applications. Some of these activities may involve considerable contact with the treated foliage and may coincide with the timing of application of Indar 75 WSP.

In the absence of data, a Tier 1 assessment of re-entry exposure was conducted assuming that 20% of the application rate is dislodgeable and that dislodgeable foliar residues (DFRs) would dissipate at a rate of 10% per day. In addition, for multiple applications, residues were assumed to be cumulative (i.e., any residue remaining before the next application was added and each application was assumed to add the same DFR as the first application). In order to derive an estimate of dermal exposure, the estimated DFR data are coupled with generic transfer coefficients appropriate for re-entry activities conducted in stone fruit orchards. The applicant, Dow AgroSciences, is a member of the Agricultural Re-entry Task Force (ARTF), and as such, the transfer coefficient of 3000 cm²/hr for thinning and harvesting, generated from a study sponsored by ARTF, was used. Thinning and harvesting is considered to be the activity with the highest potential for dermal exposure. Assuming an 8-hour workday and a body weight of 70 kg, dermal deposition of fenbuconazole each day after application was calculated using the following equation:

$$\text{Dermal Deposition (mg/kg bw/day)} = \frac{\text{TC (cm}^2\text{/hr)} \times \text{DFR (}\mu\text{g/cm}^2\text{)} \times 8 \text{ hr} \times 1 \text{ mg}}{70 \text{ kg bw} \times 1000 \mu\text{g}}$$

To estimate the dermally absorbed exposure, a dermal absorption factor of 12% was applied.

Re-entry exposure estimates for the various stone fruit crops were derived based on the proposed spray schedule for use of Indar 75 WSP on stone fruit in Canada (see Intended use section). This table indicates the growth stages at which Indar 75 WSP potentially could be applied (up to 7 applications/season); however, it is unlikely that the product would be applied at every one of those stages, since the typical number of applications to each crop would be approximately 3 or 4. It is assumed that the growing season of peaches, apricots, and nectarines would be 120 days (4 months) and of cherries, 100 days. It is also assumed that re-entry exposure would occur every day during the growing season. The daily time-weighted average exposure, margins of exposure, lifetime average daily dose (LADD) and cancer risk for re-entry workers are presented in Table 3.5.3.

For re-entry workers, MOEs ranged from 200 to 300. These are considered to be acceptable.

Cancer risks ranged from 8×10^{-6} to 1×10^{-5} . These cancer risks are considered acceptable in light of the conservatism in the exposure assessment. These conservatisms include:

- The assumption that the maximum number of applications per year would be applied, when according to typical practices, applications would be 3 to 4 per year;
- The assumption that exposure would occur every day during the growing season (up to 4 months) and that workers would work 8 hours per day for that duration of time;
- Use of the highest transfer coefficient for thinning and harvesting;
- Based on acute toxicity data of fenbuconazole (Toxicity Category IV), a restricted entry interval of 12 hours must be specified on the label.

Table 3.5.3 Re-entry exposure from use of Indar 75 WSP on stone fruits

Crop	Daily time-weighted average exposure ² ($\mu\text{g}/\text{kg bw}/\text{year}$)	Margin of exposure ³	LADD ⁴ ($\text{mg}/\text{kg bw}/\text{day}$)	Cancer Risk ⁵
peaches	4.23	300	7.43e-04	1e-05
apricots nectarines peaches	2.8	500	4.90e-04	8e-06
cherries	6.04	200	8.83e-04	1e-05

¹ Assuming the growing season is days 1 to 120 (peaches, apricots, nectarines), or days 1 to 100 (cherries).

² Including 12% dermal absorption.

³ Based on NOAEL 1.3 mg/kg bw/day from a 3-month oral rat study

⁴ Lifetime Average Daily Dose, calculated assuming 100 or 120 days of exposure per year, a life expectancy of 75 years, and a working lifetime of 40 years.

⁵ Based on Q_1^* of $1.54 \times 10^{-2} (\text{mg}/\text{kg bw}/\text{day})^{-1}$

Note: Post-application estimates for plums were not determined since it was assumed to be covered off by other crops.

3.5.4 Consumers

Not applicable.

4.0 Residues

4.1 Food residues exposure assessment

Nature of the residue in plants

Fenbuconazole (RH-7592, 6.8% a.i. emulsifiable concentrate) radiolabelled as phenyl- ^{14}C]RH-7592 or triazole- ^{14}C]RH-7592 was applied five times at a rate of 200 g a.i./ha (1 kg a.i./ha/season) on peach trees. In the experiments conducted with phenyl-labelled RH-7592, the parent compound (RH-7592) and the lactone metabolite (RH-9129) were the only two compounds identified. These compounds were present at residue levels of

0.036 ppm and 0.011 ppm, respectively. Since the lactone metabolite contains two asymmetric carbons, it can exist in the form of two stereo-isomers. The other stereoisomer of lactone metabolite, RH-9130, was not identified in any peach metabolism sample. The glucose conjugates of RH-4911 were also found at levels of 0.006 ppm. In the experiments conducted with triazole-labelled RH-7592, triazole alanine (RH-3968) and parent compound (RH-7592) were predominant residues found at levels of 0.062 ppm and 0.020 ppm, respectively. The lactone metabolite (RH-9129) and triazole acetic acid (RH-4098) at levels <0.01 ppm were also identified. Based on the peach metabolism study, the residue of concern is defined as the parent RH-7592 and its lactone metabolites, RH-9129 and RH-9130.

Confined accumulation in rotational crops

Stone fruit are perennial crops. No other food or feed crop is grown in a stone fruit orchard. Therefore, there is no concern for secondary residues in any other food or feed.

Field accumulation in rotational crops

Stone fruit are perennial crops. No other food or feed crop is grown in a stone fruit orchard. Therefore, there is no concern for secondary residues in any other food or feed.

Nature of the residue in animals

No poultry or livestock feed item is associated with the use of fenbuconazole on stone fruit. Therefore, no animal metabolism study is needed.

Methods for residue analysis of food of animal origin

No poultry or livestock feed item is associated with the use of fenbuconazole on stone fruit. Therefore, no analytical method for animal commodities is needed.

Methods for residue analysis of plants and plant products

The analytical method (TR 34-90-47R) was proposed for supervised residue trials and tolerance enforcement. This method determines fenbuconazole (RH-7592) and its metabolites (RH-9129 and RH-9130) by gas liquid chromatography using capillary column and thermionic specific detector. The limit of detection (LOD) for RH-7592, RH-9129, and RH-9130 in stone fruit was established at 0.01 ppm. The limit of quantitation (LOQ) for the RH-7592, RH-9129, and RH-9130 in stone fruit was established at 0.05 ppm. This method was found to give adequate recoveries in stone fruit matrices. The recoveries were $89 \pm 6\%$ for RH-7592, $90 \pm 6\%$ for RH-9130 and $82 \pm 10\%$ for RH-9129. The detector response for all analytes was linear over the range of 0.05–1.0 ppm. The interlaboratory validation of method supported the reliability and reproducibility of the analytical method to determine the residue of fenbuconazole and its metabolites in stone fruit matrices.

Storage stability data

Samples of stone fruit spiked with fenbuconazole (RH-7592) and its metabolites (RH-2930 and RH- 2929) at a level of 0.5 ppm were stored at approximately -10°C for a duration of up to 54.5 months. All samples were analysed at 0, 92, 182, 214, 365, 555,

723, 909, 1088, 1263, 1471, and 1634 days of freezer storage. Residues of fenbuconazole (RH-7592) and metabolites (RH-9129 and RH-9130) were stable in stone fruit at approximately -10°C up to 54.5 months.

Crop field trials

Supervised crop field trials were conducted on apricot, cherry, peach, plum, and fresh prune in the United States (U.S.). Results indicated that maximum combined residues of fenbuconazole and the metabolites were in apricot 0.27 ppm, cherries 0.749 ppm, peaches 0.5 ppm, plum and prune 0.08 ppm, when plants were treated six to twelve times with fenbuconazole formulated as 2F or Indar 75 WP at a rate of 672–1344 g a.i./ha/season (0.9–1.82 × proposed GAP). The residue trials submitted for peaches are accepted as bridging data to support the use of Indar 75 WSP on nectarines.

Consequently, MRLs in/on apricot (0.3 ppm), cherry (0.8 ppm), nectarine (0.5 ppm), peach (0.5 ppm), plum (0.1 ppm), fresh prune (0.1 ppm), and dry prune (0.5 ppm) are recommended to cover residues of RH-7592, RH-9129, and RH-9130.

Processed food/feed

Fenbuconazole (2F formulation; 23.5% a.i.) was applied to plum at a rate of 672 or 694 g a.i./ha (approximately 1× the proposed Canadian season rate). The fresh plums were processed into dry prunes. A comparison of the residues in the raw agricultural commodity (RAC) with those in processed fruit demonstrated concentration in dry prune. When treated according to the proposed Canadian use pattern, the concentration factor for dry prune is not expected to exceed 5. The residues of fenbuconazole and its lactone metabolites in dry prune will be covered under the recommended MRL of 0.5 ppm.

Meat/milk/poultry/eggs

No poultry or livestock feed item is associated with the use of fenbuconazole on stone fruit. It is expected that no residue of fenbuconazole will be present in poultry and livestock food commodities as a result of this use.

Dietary risk assessment

The proposed domestic use of fenbuconazole on apricot, cherry, nectarine, peach, plum, and prune does not pose an unacceptable chronic, acute, or lifetime dietary cancer risk from food and water to any segment of the population, including infants, children, adults, and seniors. The available refinements (Canadian Supervised Trial Median Residues (STMRs), U.S. mean residue, processing study data, and estimated percent crop treated information) were used. For chronic risk from dietary exposure to the residues of fenbuconazole and its metabolites from food and water, the potential daily intake (PDI) was less than 3% of the acceptable daily intake (ADI) for all population subgroups including infants, children, adults, and seniors. For acute dietary intake at the 95th percentile, exposure to the residues of fenbuconazole and its metabolites represented less than 2% of the ARfD for the women 13+ years of age. The life time cancer risk from dietary exposure to the residues of fenbuconazole and its metabolites from food and water was estimated to be 1.72e-06 for all infants (<1 year) and children 1–6 years, and < 8e-07

for the rest of the subgroups. It is expected that further refinement would result in a lifetime risk less than the level of concern of 1.00e-06.

5.0 Fate and behaviour in the environment

5.1 Physical and chemical properties relevant to the environment

Fenbuconazole has low solubility in water and is not expected to dissociate in water. The vapour pressure and Henry's Law constant indicate that fenbuconazole is non-volatile. Furthermore, fenbuconazole has a low potential for ultraviolet light-induced phototransformation under normal environmental conditions. The octanol-water partitioning coefficient is relatively high indicating that there is potential for the active ingredient to bioaccumulate in organisms (Table 5.1.1). Data were not available on the physicochemical properties of environmental transformation products.

Table 5.1.1 Physical and chemical properties of the active ingredient relevant to the environment

Property	Value	Comments
Solubility in water	3.8 mg/L	The solubility of the active ingredient in water is classified as low.
Vapour pressure	0.37×10^{-7} mm Hg (0.005 mPa)	According to Kennedy and Talbert (1977), the active ingredient is non-volatile under field conditions.
Henry's Law constant	4.3×10^{-9} atm·m ³ /mol or 5.57×10^6 (1/H)	The active ingredient is non-volatile from water or moist soil.
log K _{ow}	K _{ow} = 1700 ± 300 log K _{ow} = 3.22 ± 0.08	There is potential for the active ingredient to bioaccumulate.
pK _a	No value provided	The active ingredient is not expected to dissociate in water.

Property	Value		Comments
UV-visible absorption	<u>λ max (nm)</u>	<u>ϵ (L·mol⁻¹·cm⁻¹)</u>	The active ingredient has a low potential for ultraviolet light-induced phototransformation under normal environmental conditions.
	196	53 000	
	262	750	
	268	740	
	275	480	

5.2 Abiotic transformation

Laboratory studies on the hydrolysis, phototransformation on soil, and phototransformation in water were submitted to determine the effect of abiotic processes on fenbuconazole.

In an hydrolysis study, the half-lives of fenbuconazole were extrapolated to 2210, 3740, and 1340 days (equivalent to 6.1, 10.2, and 3.7 years) at pHs 5, 7, and 9, respectively. This indicates that fenbuconazole is stable to hydrolysis at environmentally-relevant pHs.

The phototransformation on soil was slow, with a half-life of 79 d under conditions of 12 h light:12 h dark. Only two unidentified minor transformation products were measured on soil reaching maxima of 3.03% and 2.75% of the applied radioactivity. Fenbuconazole did not phototransform in water. The half-life for phototransformation in water was estimated at 1280 d (~3.5 years) under conditions of 12 h light to 12 h dark. Transformation products were not detected in water. Phototransformation is not an important route of transformation of fenbuconazole on soil and is not a route of transformation in water. Data on the phototransformation in air were not required as volatilization is not expected.

Fenbuconazole does not undergo hydrolysis or photolysis in water; therefore, it can be considered to be stable to abiotic processes in water. Fenbuconazole undergoes very limited phototransformation on soil with two minor, unknown transformation products produced from this process; therefore, fenbuconazole is not subject to abiotic transformation mechanisms.

5.3 Biotransformation

Laboratory studies on the biotransformation of fenbuconazole in aerobic soil, anaerobic soil, and aerobic water/sediment systems were reviewed to determine the effect of biotic (microbial) processes on fenbuconazole.

Biotransformation processes were examined in two aerobic soils. An average of 45.7% of the parent compound transformed under aerobic conditions between day 7 and day 363 of the study (silty clay loam: 53.6%; sandy loam: 37.6%). The half-lives were 258 d and 367 d in silty clay loam and sandy loam soils, respectively. The major transformation

products detected were RH-9129 and free triazole in the silty clay loam, with major transformation products not detected in the sandy loam samples. Minor transformation products RH-9130 and RH-6467 were formed in silty clay loam, while in sandy loam, the minor transformation products were free triazole, RH-9129, RH-9130, and RH-6467, with high quantities (approximately 20%) of unidentified radioactivity (products). Biotransformation is a route of transformation of RH-7592 in aerobic soils, but this is a slow process. According to the classification scheme of Goring et al. (1975), fenbuconazole is persistent in soil under aerobic conditions.

The transformation of fenbuconazole was examined in two anaerobic soils. Following 30 days of aerobic aging, less than 3.2% of fenbuconazole had mineralized to CO₂. By the end of the study, 7.8% of the parent compound had transformed. The half-lives of fenbuconazole in anaerobic silty clay loam and sandy loam were 451 d and 655 d, respectively, exceeding the half-lives in aerobic soils. Major transformation products were not detected. The minor transformation products RH-9129, RH-9130, and RH-6467 were detected with high quantities of unidentified radioactivity. As only 7.8% of the parent compound transformed under anaerobic conditions after 60 days, biotransformation is a route of transformation in anaerobic soil; however, this is a very slow process. According to the classification scheme of Goring et al. (1975), fenbuconazole is persistent in soil under anaerobic conditions.

In a study on aerobic aquatic biotransformation, fenbuconazole strongly partitioned to sediment. For example, the concentration of fenbuconazole in river water decreased from 96% of the applied radioactivity (AR) at initiation to 3.6% AR after 105 days. As concentrations in water decreased, the concentration in sediments increased from 11% AR to 79% AR. A similar effect was observed in the pond system. Major transformation products were detected in neither water nor sediment. Naming/identification conventions were inconsistent and only certain transformation products were specifically identified as RH-6467 or RH-129/130 (also called RH-29/30 or RH-99129/RH-99130). The maximum dissipation times in water were 4.3 and 1.2 days in river and pond water, respectively. The study did not present dissipation times for sediment; however, for the entire system, the maximum DT_{50s} exceeded 1000 days. The results indicate faster dissipation of fenbuconazole in the pond system. This may be a direct consequence of the amount of total organic carbon (TOC) in the water as the pond water contained approximately twice the level of TOC as the river water. Thus, the study's author speculated that adsorption to suspended colloidal organic particles may be a process by which scavenging of fenbuconazole takes place. This process may augment partitioning to the sediment layer. Another explanation for the more rapid dissipation in the pond system may be that there is more microbial activity in pond water than in river water. Although transformation of fenbuconazole appeared to occur in water, aerobic biotransformation is not an important route of transformation in water/sediment systems. The primary route of dissipation of fenbuconazole in aerobic water/sediment systems is partitioning to sediment. According to the classification scheme of McEwen and Stephenson (1979), fenbuconazole is non-persistent in water and persistent in sediment.

A request for a waiver was granted in place of an anaerobic sediment/water study as the aquatic biotransformation behaviour of RH-7592 under anaerobic conditions is described by other submitted studies.

Biotransformation is a route of transformation of fenbuconazole under aerobic and anaerobic conditions in soil, but this is a slow process. In aquatic environments, fenbuconazole appears to be rapidly removed from water; however, it does not undergo significant biotransformation in water/sediment systems. Rather, the compound is partitioned to sediments, acting as a sink for this compound. Based on studies of biotransformation, according to the classification schemes of Goring et al. (1975) and McEwen and Stephenson (1979), fenbuconazole is persistent in soil and water/sediment systems (i.e., in the sediment portion). Major transformation products (RH-9129 and free triazole) were only detected in aerobic soils.

5.4 Mobility

The adsorption/desorption characteristics of fenbuconazole were studied in five types of soil from the United States by a batch equilibrium experiment. After 84 h of equilibration, 86%, 80%, 74%, 62%, and 48% of the applied RH-7592 was adsorbed in sandy loam, loam, silty clay loam, clay, and sand, respectively. The adsorption K_d values ranged from 5.1 (clay) to 115 mL/g (sandy loam) with corresponding adsorption K_{oc} values ranging from 2185 (clay) to 9042 mL/g (sandy loam). Based on these laboratory results, fenbuconazole is immobile in loam and sandy loam and has slight to low potential for mobility in clay, sand, and silty clay loam. Adsorption appears to be associated with the percentage of organic matter present in soil. According to the classification scheme of McCall et al. (1981), fenbuconazole will be slightly mobile in soils containing a low percentage of organic carbon (generally $\leq 1\%$) and relatively immobile in soils with higher levels of organic carbon. The results from the desorption phase were not accepted for Canadian regulatory purposes.

An aged-soil leaching study was conducted on the sandy loam soil used in the adsorption/desorption study. Less than one percent of the recovered activity was detected beyond the 0–6 cm segment, 0.2% of the activity was found in the leachate, and $<1.0\%$ $^{14}\text{CO}_2$ evolved during the aging process. Three minor transformation products (RH-99129, RH-99130, and RH-96467) were detected in the aged soil residues, accounting for less than 10% of the total activity. RH-99129 and RH-99130 are diastereomers. The K_{oc} value was calculated to be >3445 . Thus, according to the classification scheme of McCall et al. (1981), fenbuconazole has a slight potential for mobility in sandy loam soil. In the adsorption/desorption study, the adsorption K_{oc} value for sandy loam was 9000, for which RH-7592 is classified as immobile according to McCall et al. (1981).

Thus, from the laboratory studies of mobility, fenbuconazole is relatively immobile in soils and would not be expected to leach. Based on the Henry's Law constant and vapour pressure (Table 1.2.1), fenbuconazole is not expected to volatilize from soils or moist

surfaces, including water. The high K_{oc} and K_{ow} indicate that fenbuconazole is expected to partition to sediments. Partitioning to sediments was confirmed in the study on biotransformation in aerobic water/sediment.

5.5 Dissipation and accumulation under field conditions

Under terrestrial field conditions, the DT_{50} s of fenbuconazole were calculated by the registrant to be 161 days and 314 days at the Midwest and northern California field sites, respectively. These DT_{50} s are in general agreement with those reported in the aerobic soil biotransformation study (285 and 367 d in silty clay loam and sand loam, respectively). Examination of the data by the PMRA indicates that the DT_{50} at the Midwest site was >364 d at one of three replicate plots and could not be determined at the two other plots. Similarly, the PMRA determined that the DT_{50} s at the northern California plots were 198 d at one of the three replicate plots and >364 d at two of the replicate plots. The data indicate that fenbuconazole is persistent under field conditions according to the classification scheme of Goring et al. (1975). Major transformation products were not detected in the field studies, although four minor transformation products (RH-9130, RH-9129, RH-6467, and 1,2,4-triazole) were measured.

Based on the long DT_{50} s, carryover of residues of fenbuconazole and its transformation products will occur if repeated applications are made from season to season. If the compound's use is extended to field crops, these residues may be available for uptake by rotated crops. Residues of some transformation products were infrequently detected below 30 cm; however, there was no general trend for leaching over the 12- to 18-month duration of the field trials. Despite the lack of leaching, the pattern of dissipation of fenbuconazole was not clearly identified in the two sites applicable to Canadian conditions; therefore, the guideline requirements for a terrestrial field dissipation study have not been fulfilled.

There is good agreement between laboratory and field data in the classification of the persistence, lack of leaching, and immobility of fenbuconazole in soil. The exact dissipation route of fenbuconazole in soil has not been confirmed between laboratory and field studies.

In a separate field study, the percent interception of fenbuconazole by orchard trees was calculated to be 56% and the DT_{50} on turf was determined to be 6.7 days. Note that this study was not conducted according to any international protocol and good laboratory practices or quality assurance were not provided. Therefore, the results should be interpreted with caution.

Data are not available on the dissipation of fenbuconazole under aquatic field conditions.

5.6 Bioaccumulation

The octanol–water partition coefficient for fenbuconazole is 1700 ± 300 . Given that the $\log K_{ow}$ exceeds three ($\log K_{ow} = 3.22 \pm 0.08$), there is potential for the compound to bioaccumulate in biological organisms. The PMRA identified the parent compound (fenbuconazole) and RH-9129/RH-9130 (the lactone stereoisomers) as the residues of concern in plants. No residue of concern was identified for animals because stone fruit are not fed to animals.

A laboratory study with rats indicated that fenbuconazole was rapidly absorbed, distributed, and excreted primarily via the feces within 24 to 48 hours post-dosing. Tissue distribution and bioaccumulation were minimal. Elimination was biphasic with an initial rapid phase (24–48 hours post-dosing) followed by a slower decline (48–96 hours post-dosing).

Although three studies on the bioaccumulation in fish were submitted, only two studies were reviewed (see Appendix I for rationale for not reviewing the additional study). Similar results were found for fish, although the elimination period was longer than in rats. Bioconcentration factors for bluegill sunfish were 170 \times , 50 \times , and 330 \times in whole fish, fillets, and visceral tissues, respectively, with 95–98% of concentrated residues depurated over a 14-day period. Five transformation products were identified: lactone A (RH-9129), a ketone (RH-6467), two polar stereoisomers, and the sulfate conjugate of a benzyl alcohol intermediate in the pathway leading to formation of the lactone and ketone. An unknown transformation product was tentatively identified as the glucuronide conjugate of the proposed benzylic alcohol intermediate.

5.7 Summary of fate and behaviour in the terrestrial environment

Fenbuconazole undergoes very limited phototransformation on soil with a DT_{50} of 79 d. Therefore, fenbuconazole is not subject to abiotic transformation mechanisms in the terrestrial environment.

Biotransformation is a route of transformation of fenbuconazole in aerobic and anaerobic soils, although this is a slow process in aerobic soils (half-lives of 258 and 367 d for aerobic silty clay loam and sandy loam soils, respectively) and a very slow process in anaerobic soils (half-lives of 451 and 655 d in anaerobic silty clay loam and sandy loam, respectively). Thus, fenbuconazole is classified as persistent in soil under both aerobic and anaerobic conditions. Two major transformation products were detected in aerobic soils (RH-9129 and free triazole). As fenbuconazole is non-volatile, studies on the biotransformation in air are not required.

The mobility of fenbuconazole was reviewed in laboratory studies of adsorption/desorption and aged-soil leaching. The adsorption K_d values for fenbuconazole range from 5.1 to 115 mL/g while corresponding adsorption K_{oc} values range from 2185 to 9042 mL/g. Fenbuconazole is classified as immobile in loam and sandy loam and has

slight to low potential for mobility in clay, sand, and silty clay loam. Adsorption appears to be associated with the percentage of organic matter present in soil. Fenbuconazole will be slightly mobile in soils containing a low percentage of organic carbon (generally $\leq 1\%$) and relatively immobile in soils with higher levels of organic carbon. The results from the desorption phase were not accepted for Canadian regulatory purposes. In an aged-soil leaching study, less than one percent of the recovered activity was detected beyond the 0–6 cm segment, in the leachate, or as $^{14}\text{CO}_2$. A K_{oc} value was calculated to be greater than 3445, indicating that fenbuconazole has slight potential for mobility in sandy loam soil. In the adsorption/desorption study, the adsorption K_{oc} value for sandy loam was 9000, for which RH-7592 was classified as immobile.

Thus, from the laboratory studies of mobility, fenbuconazole is relatively immobile in soils and would not be expected to leach. Based on the Henry's Law constant and vapour pressure (Table 1.2.1), fenbuconazole is not expected to volatilize from soils or moist surfaces, including water. The high K_{oc} and K_{ow} indicate that fenbuconazole may be expected to partition to sediments. Partitioning to sediments was confirmed in the study on biotransformation in aerobic water/sediment.

Under terrestrial field conditions, the registrant reported DT_{50} s of 161 and 314 d for fenbuconazole at two field sites. These half-lives are in general agreement with those reported in the aerobic soil biotransformation study. Examination of the data by the PMRA indicates that the DT_{50} at the Midwest site was >364 days at one of three replicate plots and could not be determined at the two other plots. Similarly, the PMRA determined that the DT_{50} s at the northern California plots were 198 d at one of the three replicate plots and >364 d at two of the replicate plots. The data indicate that fenbuconazole is persistent under field conditions. Major transformation products were not detected in the field studies, although four minor transformation products (RH-9130, RH-9129, RH-6467, and 1,2,4-triazole) were measured. No leaching was observed; however, based on the reportedly long half-lives, carryover of residues of fenbuconazole and its transformation products will occur if repeated applications are made from season to season. If the compound's use is extended to field crops, these residues may be available for uptake by rotated crops. Although there is good agreement between laboratory and field data in the classification of the persistence, lack of leaching, and immobility of fenbuconazole in soil, the exact dissipation route of fenbuconazole in soil has not been confirmed between laboratory and field studies. As the pattern of dissipation of fenbuconazole was not clearly identified, the guideline requirements for a terrestrial field dissipation study have not been fulfilled. However, as the laboratory data support the persistence of fenbuconazole, no terrestrial field study is required at this time. In a separate field study, the percent interception of fenbuconazole by orchard trees was calculated to be 56% and a half-life on turf was determined to be 6.7 days.

Given the octanol–water partition coefficient for fenbuconazole of 1700 ± 300 ($\log K_{ow} = 3.22 \pm 0.08$), there is potential for the compound to bioaccumulate in biological organisms. Although fenbuconazole accumulated in both rats and fish, the compound was depurated from both organisms with a more rapid elimination in rats than in fish. The

transformation products in fish included RH-9129 (lactone), RH-6467 (ketone), two polar stereoisomers, and the sulfate conjugate of a benzyl alcohol intermediate in the pathway leading to formation of the lactone and ketone. A third transformation product was tentatively identified as the glucuronide conjugate of the benzylic alcohol intermediate.

5.8 Summary of fate and behaviour in the aquatic environment

Fenbuconazole may be expected to enter the aquatic environment through direct overspray, spray drift from orchard blast application, and/or runoff via sorption to soil particles.

Fenbuconazole does not undergo hydrolysis or phototransformation in water at environmentally relevant pHs. No major or minor transformation products were formed in either of the studies on the hydrolysis or phototransformation of fenbuconazole in water.

Studies on the biotransformation of fenbuconazole in aerobic water and in anaerobic sediment/water were not submitted because fenbuconazole partitions to sediment and the behaviour in sediment was described in the anaerobic segment of the aerobic water/sediment study. In a study on the biotransformation of fenbuconazole in aerobic water/sediment, the longest dissipation times for water were 4.3 days and 1.2 days in river and pond water, respectively, indicating that fenbuconazole is rapidly removed from water (non-persistent). Although dissipation times for sediment were not presented, the DT_{50s} for the river and pond water/sediment systems exceeded 1000 days. Major transformation products were not detected in the aerobic water/sediment study. Although minor transformation products were formed, aerobic biotransformation in water/sediment is not an important route of transformation of fenbuconazole. The fate of fenbuconazole in aerobic water/sediment systems is partitioning to sediment where it is classified as persistent.

Data are not available on the dissipation of fenbuconazole under aquatic field conditions.

5.9 Expected environmental concentrations

The expected environmental concentrations (EEC) of fenbuconazole in environmental compartments of concern were estimated based on calculations made using simple scenarios. These concentrations were used as initial approximations for estimating the potential exposure to wildlife. It was assumed that fenbuconazole was applied at the maximum proposed Canadian label rate of 0.105 kg a.i./ha. The application pattern was based on the tentative application uses supported by the PMRA: 7 applications per season with applications 7 days apart with the exception of a 60-day interval between Application Numbers 5 and 6. The scenario assumes that the concentrations in the various environmental compartments were obtained immediately following the last of the applications.

5.9.1 Soil

The EEC of fenbuconazole in soil was calculated assuming application to bare soil with a soil bulk density of 1.5 g/cm³ and a soil depth of 15 cm. Seven applications at the maximum proposed Canadian label rate of 0.105 kg a.i./ha were used in the pattern outlined in Section 5.9. Using the most conservative registrant-reported DT₅₀ of 314 d in soil (field dissipation study), the concentration of fenbuconazole in soil immediately following the seventh application is equivalent to a cumulative application of 0.650 kg a.i./ha. Based on the maximum cumulative application rate, the EEC in soil was estimated to be 0.29 mg a.i./kg soil dry weight (dw).

5.9.2 Aquatic systems

The DT₅₀ of 4.3 d from the study on the biotransformation of fenbuconazole in aerobic water/sediment was used to calculate the EEC resulting from direct overspray of fenbuconazole to aquatic systems. Using the application pattern outlined in Section 5.9 (7 applications at the maximum proposed Canadian label rate of 0.105 kg a.i./ha at 7-d and 60-d intervals), the EEC of fenbuconazole in water immediately following the seventh application is the equivalent of a cumulative application of 0.150 kg a.i./ha. Assuming a scenario in which a body of water 30 cm deep is oversprayed with the equivalent of the cumulative application rate, the EEC in water is 0.05 mg a.i./L water. Although this scenario may be unrealistic for ground application, it is useful as a first approximation and is used to compare the EECs in aquatic systems and the no observed effect concentrations (NOECs) from environmental toxicology studies.

Based on the potential use pattern of fenbuconazole in areas where stone fruit are grown, residues of fenbuconazole in potential drinking water sources in these areas (i.e., groundwater and reservoirs) were modelled using the models LEACHM for groundwater and PRZM/EXAMS for surface water. As the proposed use pattern does not include use in the prairie region, concentrations of fenbuconazole in dugouts were not determined.

Because the modelling results from the Level I screening model failed the human health assessment conducted by the PMRA, a refined analysis was conducted (Level II). The Level II analysis represents a less conservative approach to predicting drinking water concentrations of the active ingredient as it more accurately reflects the use pattern of the chemical. Three scenarios typical of stone fruit-growing regions were used in the water modelling: an apple-growing region in British Columbia, a grape-growing region in Niagara, Ontario, and an apple-growing region in Nova Scotia. Because, in general, the most suitable soils for stone fruit production in Canada are sandy loam soils, the Level II assessment used input parameters specific to sandy loam soil, where applicable.

The most conservative estimates of the EECs in drinking water sources were 2.2 µg a.i./L and 0.25 µg a.i./L for acute and chronic exposures, respectively. These values were provided for the human health assessment. As a Level II drinking water assessment was

conducted, any further use expansion beyond stone fruit-growing regions (e.g., Okanagan Valley, British Columbia, or the Niagara Region, Ontario) will require a re-evaluation of drinking water concentrations for two reasons: concentrations in dugouts were not examined and the use of fenbuconazole may be expanded to new crops for which drinking water scenarios were not modelled at Level II.

5.9.3 Vegetation and other food sources

Data were not provided on concentrations of fenbuconazole on foliar crops immediately after application. In the absence of these data, concentrations of fenbuconazole on vegetation and insects resulting from direct over-spray were estimated using a nomogram developed by the U.S. EPA from the data of Hoerger and Kenaga (1972) and Kenaga (1973), for use in ecological risk assessment (Urban and Cook 1986). EECs were determined for two scenarios:

- (1) no transformation between first and last (seventh) application and no interception of airblast spray by orchard trees (Table 5.9.3a)
- (2) dissipation from turf using a DT_{50} of 6.7 d and 56% interception of airblast spray (from Batra 1995, DACO 8.2.3.6). This refined scenario was applied to all vegetative matter in the diets of wild birds and mammals (Table 5.9.3b).

A fresh weight to dry weight conversion was calculated for both scenarios.

Table 5.9.3a Maximum EEC in vegetation and insects after a direct over-spray assuming no transformation and no interception of airblast spray by orchard trees

Matrix	EEC (mg a.i./kg fw) ^a	Fresh/dry weight ratios	EEC (mg a.i./kg dw)
Short range grass	157	3.3 ^b	519
Leaves and leafy crops	82.3	11 ^b	906
Long grass	72.0	4.4 ^b	317
Forage crops	88.2	5.4 ^b	476
Small insects	38.2	3.8 ^c	145
Pods with seeds	7.86	3.9 ^c	30.7
Large insects	6.54	3.8 ^c	24.9
Grain and seeds	6.54	3.8 ^c	24.9
Fruit	9.85	7.6 ^c	74.9

^aBased on correlations reported in Hoerger and Kenaga (1972) and Kenaga (1973)

^bFresh/dry weight ratios from Harris (1975)

^cFresh/dry weight ratios from Spector (1956)

Table 5.9.3b Maximum EEC in vegetation after a direct over-spray assuming dissipation on vegetation and interception of airblast spray by orchard trees

Matrix	EEC (mg a.i./kg fw) ^a	Fresh/dry weight ratios	EEC (mg a.i./kg dw)
Short range grass	18.6	3.3 ^b	61
Leaves and leafy crops	9.74	11 ^b	107
Long grass	8.5	4.4 ^b	38
Forage crops	10.4	5.4 ^b	56
Pods with seeds	0.93	3.9 ^c	3.6
Grain and seeds	0.77	3.8 ^c	2.9
Fruit	1.17	7.6 ^c	8.9

^aBased on correlations reported in Hoerger and Kenaga (1972) and Kenaga (1973)

^bFresh/dry weight ratios from Harris (1975)

^cFresh/dry weight ratios from Spector (1956)

Wild birds and mammals could be exposed to residues of fenbuconazole as a result of the consumption of sprayed vegetation and/or contaminated prey. The EECs of fenbuconazole in the diets of bobwhite quail (*Colinus virginianus*), mallard ducks (*Anas platyrhynchos*), rats, mice, and rabbits were calculated for the two scenarios presented in Tables 5.9.3a and 5.9.3b. The calculations of the EECs used in the risk assessment for wild birds and mammals are outlined below and presented in Table 5.9.4.

Wild Bird Scenario 1 (no transformation and no interception assumed):

The EECs of fenbuconazole in the diets of bobwhite quail (*Colinus virginianus*) and mallard duck (*Anas platyrhynchos*) were calculated based on the maximum proposed application rate of 0.735 kg a.i./ha (Table 5.9.4). The diet of bobwhite quail consists of approximately 55% grain, 30% small insects, and 15% forage crops. The EECs of fenbuconazole on these items were calculated to be 13.7, 43.6, and 71.4 mg a.i./kg dry weight of diet, respectively, based on the maximum application rate of 0.735 kg a.i./ha. Therefore, the EEC of fenbuconazole in the diet of the bobwhite quail is 129 mg a.i./kg dw. For mallard ducks, the diet consists of approximately 70% grain and 30% arthropods. The EECs of fenbuconazole on these items were calculated to be 17.4 and 7.5 mg a.i./kg dw of diet, respectively, based on the maximum application rate of 0.735 kg a.i./ha. Therefore, the EEC of fenbuconazole in the diet of the mallard duck is 25 mg a.i./kg dw.

Wild Bird Scenario 2 (assumes transformation using DT₅₀ for turf and 56% interception):

The EECs in diets of bobwhite quail and mallard duck decrease significantly when transformation on vegetation and interception of the airblast spray by orchard trees are

taken into account. For bobwhite quail, the EEC decreases from 129 mg a.i./kg dw to 53.5 mg a.i./kg dw. For mallard duck, the EEC in the diet is reduced from 25 mg a.i./kg dw to 9.5 mg a.i./kg dw. The refined EECs from Wild Bird Scenario 2 are approximately 59% and 62% less than those from the unrefined Wild Bird Scenario 1 for the bobwhite quail and mallard duck, respectively.

Wild Mammal Scenario 1 (no transformation and no interception assumed):

The rat diet consists of approximately 70% short grass, 20% grain/seeds, and 10% large insects. The EECs of fenbuconazole on these items were calculated to be 363, 5.0, and 2.5 mg a.i./kg dw, respectively. Therefore, the EEC in the diet of the rat is approximately 371 mg a.i./kg dw. For mice, the diet consists of approximately 50% grain/seeds, 25% short grass, and 25% leaves/leafy crops for which the EECs were calculated as 12.4, 130, and 226 mg a.i./kg dw, respectively. Therefore, the EEC in the diet of the mouse is approximately 369 mg a.i./kg dw. For rabbits, the diet consists of approximately 25% of each of short grass, leaves/leafy crops, long grass, and forage crops. The EECs on these items were calculated to be 130, 226, 79.2, and 119 mg a.i./kg dw, respectively. Therefore, the EEC in the diet of the rabbit is approximately 554 mg a.i./kg dw.

Wild Mammal Scenario 2 (assumes transformation using DT₅₀ for turf and 56% interception):

For the three mammalian species, the EECs in diets decrease significantly when transformation on vegetation and interception of the airblast spray by orchard trees are taken into account. For rats, the EEC decreases from 371 mg a.i./kg dw to 50 mg a.i./kg dw. For mice, the EEC in the diet is reduced from 369 mg a.i./kg dw to 54 mg a.i./kg dw. For rabbits, the EEC is reduced from 554 mg a.i./kg dw to 170 mg a.i./kg dw. The refined EECs from Wild Mammal Scenario 2 are approximately 87%, 85%, and 69% less than those from the unrefined Wild Mammal Scenario 1 for the rat, mouse, and rabbit, respectively.

Table 5.9.4 Maximum EEC in diets of birds and mammals with and without the assumption of transformation on vegetation and interception of airblast spray by orchard trees

Organism	Matrix	Maximum EEC (mg a.i./kg dw diet)	
		assuming no transformation and no interception	assuming transformation and interception
Bobwhite quail	30% small insects 15% forage crops 55% grain	129	53.5

Organism	Matrix	Maximum EEC (mg a.i./kg dw diet)	
		assuming no transformation and no interception	assuming transformation and interception
Mallard duck	30% large insects 70% grain	25	9.5
Rat	70% short grass 20% grain/seeds 10% large insects	371	50
Mouse	25% short grass 50% grain/seeds 25% leaves and leafy crops	369	54
Rabbit	25% short grass 25% leaves and leafy crops 25% long grass 25% forage crops	554	170

6.0 Effects on non-target species

6.1 Effects on terrestrial organisms

The toxicity of fenbuconazole was studied with two types of invertebrates: earthworms and honey bees. As fenbuconazole had no significant effect on earthworm survival at any concentration tested, fenbuconazole is considered to be non-toxic to earthworms up to a concentration of 98 mg a.i./kg dw. There were no compound-related toxicity effects in an acute contact study with honey bees and fenbuconazole is classified as relatively non-toxic to honey bees. The LD₅₀ of >292 µg a.i./bee is equivalent to an application rate of >327 kg a.i./ha, which exceeds the proposed single application rate of 0.105 kg a.i./ha by more than 3000 times. Data on the toxicity to beneficial predators and parasites were not submitted. As beneficial invertebrates may be expected to be present during the time(s) of application of the end-use product in orchards, studies on the toxicity of fenbuconazole to beneficial predators and parasites are required to assess potential toxic effects.

Studies on the acute oral toxicity, acute dietary toxicity, and reproductive toxicity to birds were reviewed. Although compound-related effects were observed in all studies, fenbuconazole is classified as practically non-toxic to bobwhite quail on an acute oral basis, based on the LD₅₀ of >2150 mg a.i./kg bw. The single-dose acute oral NOEL was 1470 mg a.i./kg bw. In addition to four mortalities observed in the highest dose group

(out of 10 birds), a variety of sublethal effects were observed. In the acute dietary studies, fenbuconazole would be classified as slightly toxic to bobwhite quail and mallard duck on an acute dietary basis based on the LC₅₀s of 4954 and 2013 mg a.i./kg dw of diet, respectively. The NOEC was 312 mg a.i./kg diet for both species, although various sublethal effects were noted in the three highest test concentrations and physiological changes were observed on gross necropsy in some birds. In the reproduction toxicity studies, the NOEC of fenbuconazole to both bobwhite quail and mallard duck was 150 mg a.i./kg dw. No overt sign of toxicity was noted; however, significant reductions in growth (body weight gain) and biologically significant decreases in egg production and reduced hatchability were found at 600 mg a.i./kg dw. A toxicant-mediated effect on egg shell thickness was identified for bobwhite quail; therefore, there are toxicant-mediated effects on reproduction in avian species.

Supplemental herbicide screening reports indicate that fenbuconazole does not cause symptoms of injury (phytotoxic effects) to the 11 crops tested at application rates of 75, 150, and 750 g a.i./ha, however, insufficient information was provided for a full scientific review. The most sensitive monocot and dicot species could not be determined from the submitted information. The EC₂₅ and EC₅₀ for both monocots and dicots was >750 g a.i./ha. The NOEC was 750 g a.i./ha, which exceeds the maximum proposed seasonal application rate of 735 g a.i./ha. Additional studies on the toxicity of fenbuconazole to terrestrial vascular plants were not submitted.

6.2 Effects on aquatic organisms

Based on an acute toxicity with *Daphnia magna*, fenbuconazole is classified as moderately toxic to *Daphnia magna*. In a chronic study with *Daphnia magna*, fenbuconazole had an effect on the reproduction at concentrations greater than 0.078 mg a.i./L. In a subchronic study with midge larvae (*Chironomus riparius*), effects on survival and emergence were observed. Given the high partitioning of this product to sediments, and its relatively high toxicity, additional data on toxicity to benthic organisms will be required if the use pattern is expanded beyond what is currently proposed for stone fruit.

Fenbuconazole was found to be acutely moderately toxic to rainbow trout and highly toxic to bluegill sunfish. Sublethal effects were also observed in both species. In an early-life stage test, fenbuconazole did not significantly affect hatching success or larval survival of early life stages of fathead minnows; however, standard length was significantly reduced at the highest test concentration and wet weight was significantly reduced at the two highest test concentrations (0.16 and 0.33 mg a.i./L). Other sublethal effects were observed. In a full life-cycle toxicity study with fathead minnow, the most sensitive endpoint identified was the time to first spawn, with an NOEC of 0.027 mg a.i./L. Other measurement endpoints were also affected at higher concentrations, including number of eggs produced, survival of parents and offspring, and numbers of eggs per spawn.

In fish, results on the bioaccumulation of fenbuconazole were found to be similar to those in rats, although the elimination period was longer in fish. Bioconcentration factors for bluegill sunfish were 170×, 50×, and 330× in whole fish, fillets, and visceral tissues, respectively, with 95–98% of concentrated residues were depurated over a 14-day period. Five transformation products were identified: lactone A (RH-9129), a ketone (RH-6467), two polar stereoisomers, and the sulfate conjugate of a benzyl alcohol intermediate in the pathway leading to formation of the lactone and ketone. A third transformation product was tentatively identified as the glucuronide conjugate of the benzylic alcohol intermediate.

Studies on the acute toxicity to two species of freshwater green algae, one diatom, and one blue-green algae were reviewed. The most sensitive algal species was *Selenastrum capricornutum* Printz, with a NOEC of 0.270 mg a.i./L and the percent growth inhibition in the treated algal culture as compared to the control ranged from 0 to 100%. Abnormalities were noted in the higher concentrations tested including bloating and irregularly shaped algal cells.

Studies on the acute toxicity to a marine crustacean and toxicity to mollusk embryo larvae were reviewed. A study of the acute toxicity of fenbuconazole to *Mysidopsis bahia* determined that fenbuconazole is highly toxic to mysid shrimp. Significant differences in shell deposition were observed for the Eastern Oyster exposed to fenbuconazole. The NOEC and LOEC were 0.53 and 0.69 mg a.i./L, respectively, with a 32% decrease in shell deposition at the LOEC level. As an EC₅₀ was not determined, fenbuconazole is classified as highly toxic to the Eastern Oyster based on the LOEC. In a 96-h acute toxicity study on sheepshead minnow, *Cyprinodon variegatus*, the LC₅₀ and NOEC were 1.8 and 0.89 mg a.i./L, respectively; thus, fenbuconazole is classified as moderately toxic to sheepshead minnow. Sublethal effects were also observed. Data were not submitted on the toxicity to marine algae.

6.3 Effects on biological methods of sewage treatment

As data are not required, data were not submitted.

6.4 Risk characterization

6.4.1 Environmental behaviour

As summarized in Sections 5.7 and 5.8, fenbuconazole is persistent in soil and sediment. In addition to acute exposure, there is, therefore, potential for prolonged exposure of soil- and sediment-dwelling organisms to residues of fenbuconazole. Further exposure of terrestrial organisms can be expected from the consumption of contaminated vegetation. Results from laboratory and field studies, as well as groundwater modelling (Section 5.9.2) indicate that fenbuconazole is not likely to leach into groundwater; however, fenbuconazole may be expected to enter the aquatic environment through direct overspray, spray drift from orchard blast application, and/or runoff via sorption to soil

particles. Therefore, there is potential for exposure of fenbuconazole to non-target terrestrial and aquatic organisms. Based on the physicochemical properties of fenbuconazole, volatilization is not an expected route of exposure of non-target organisms.

In order to minimize entry into the environment, a summary of storage, disposal, and decontamination procedures for the active ingredient is required. The instructions for storage, disposal, and decontamination are consistent with avoiding entry of fenbuconazole into the environment.

6.4.2 Terrestrial organisms

The submitted toxicity studies were conducted using the parent compound. Data were not submitted on the toxicity of transformation products of fenbuconazole to terrestrial organisms. Therefore, the terrestrial risk assessment is for the parent compound, fenbuconazole.

The degree of risk to terrestrial (and aquatic) organisms was classified according to the following index:

Margin of safety (MOS)	Risk qualifier
≥ 10	Negligible risk
1 to <10	Low risk
0.1 to <1	Moderate risk
0.01 to <0.1	High risk
0.001 to <0.01	Very high risk
<0.001	Extremely high risk

6.4.2.1 Non-target terrestrial invertebrates

6.4.2.1.1 Earthworms

One scientifically valid and acceptable toxicity study with earthworms was submitted. The NOEC was 98 mg a.i./kg soil. The EEC of fenbuconazole in soil (0.29 mg a.i./kg soil) is below the NOEC. The margin of safety is 340; therefore, fenbuconazole poses a negligible risk to earthworms at the proposed application rate (Appendix IV, Table 5).

6.4.2.1.2 Bees

One scientifically valid and acceptable acute contact toxicity study with honey bees was submitted. According to the classification scheme of Atkins et al. (1981), fenbuconazole is relatively non-toxic to bees.

The LD₅₀ for honey bees was >292 µg a.i./bee, which is equivalent to >327 kg a.i./ha. The maximum seasonal application rate of 0.735 kg a.i./ha is lower than the LD₅₀. As the margin of safety is 445, fenbuconazole poses a negligible risk to honey bees at the proposed application rate (Appendix IV, Table 5).

6.4.2.1.3 Predators and parasites

As data were not submitted on the toxicity of fenbuconazole to beneficial predators and parasites, the risk to these organisms cannot be assessed at this time.

6.4.2.2 Birds

Wild birds, such as bobwhite quail (*Colinus virginianus*) and mallard duck (*Anas platyrhynchos*), could be exposed to residues of fenbuconazole as a result of the consumption of sprayed vegetation and/or contaminated prey. From Table 5.9.4, assuming no transformation and no interception of orchard blast spray (Wild Bird Scenario 1, see Section 5.9.3), the EECs of fenbuconazole in the diets of the bobwhite quail and mallard duck are 129 and 25 mg a.i./kg dw, respectively. When transformation on vegetation and interception by trees are taken into account (Wild Bird Scenario 2), these EECs decrease to 53.5 and 9.5 mg a.i./kg dw, respectively. Individual risk assessments were carried out for acute oral exposure to bobwhite quail, acute dietary exposure to bobwhite quail and mallard duck, and chronic exposure for reproductive effects with both avian species.

One acceptable toxicity study was submitted on the acute oral exposure of wild birds to fenbuconazole. In an acute oral study with bobwhite quail, the LD₅₀ was >2150 mg a.i./kg bw while the NOEL was 1470 mg a.i./kg bw. The average body weight per individual (BWI) of the control group in the study was 204 g and the daily individual (ind) food consumption (FC) was 0.015 kg dw. Therefore, the daily intakes of fenbuconazole (DI = FC × EEC) are 1.94 and 0.80 mg a.i./ind/d under Wild Bird Scenarios 1 and 2, respectively. Expressed on a per individual basis, the LD_{50 (ind)} and NOEL_(ind) were 439 and 300 mg a.i./ind, respectively. Based on the predicted daily intake of the active ingredient and the LD_{50 (ind)}, the number of days of intake of fenbuconazole by a bobwhite quail in the wild to attain a dose equivalent to that administered by gavage that killed 50% of the individuals in the laboratory population is 227 and 547 days under Wild Bird Scenarios 1 and 2, respectively. Similarly, based on the predicted daily intake and the NOEL_(ind), the maximum number of days of intake of fenbuconazole by a wild bobwhite quail, to attain a dose equivalent to that administered by gavage that had no observable effect on the laboratory population, is 155 and 374 days, under Wild Bird Scenarios 1 and 2,

respectively. These values indicate that the application of fenbuconazole at the maximum proposed label rate poses a negligible risk to wild bird populations, such as the bobwhite quail, that are acutely exposed to fenbuconazole (Appendix IV, Table 5).

The LD₅₀s from separate acute dietary studies with bobwhite quail and mallard duck were 4954 mg a.i./kg diet and 2013 mg a.i./kg diet, respectively. According to the U.S. EPA classification scheme, fenbuconazole is considered slightly toxic when birds are acutely exposed. The NOECs were 625 and 312 mg a.i./kg dw diet for bobwhite quail and mallard duck, respectively. Under Wild Bird Scenario 1, the EECs in the diet of the bobwhite quail and the mallard duck are expected to be 129 and 25 mg a.i./kg dw, respectively. Thus, the margins of safety for bobwhite quail and mallard duck are 4.8 and 13, respectively. Therefore, under Wild Bird Scenario 1, fenbuconazole is considered to pose a low dietary risk to bobwhite quail and a negligible dietary risk to mallard ducks at the proposed maximum application rate (Appendix IV, Table 5). Under the refined Wild Bird Scenario 2, which takes into account transformation on vegetation and interception by orchard trees, the EECs in the diet of the bobwhite quail and the mallard duck are expected to be 53.5 and 9.5 mg a.i./kg dw, respectively. Thus, the margins of safety for bobwhite quail and mallard duck are 12 and 33, respectively. Therefore, according to Wild Bird Scenario 2, fenbuconazole poses a negligible dietary risk to wild birds at the proposed maximum application rate (Appendix IV, Table 5).

Two chronic studies were submitted that examined reproductive effects in bobwhite quail and mallard duck. For both species, the NOEL and LOAEL were 150 and 600 mg a.i./kg diet, respectively. For bobwhite quail, under Wild Bird Scenario 1, the NOEL slightly exceeds the EEC in the diet of 129 mg a.i./kg diet. This results in a margin of safety of 1.2, indicating a low risk of reproductive effects in bobwhite quail. Consequently, a refined dietary risk assessment was conducted for bobwhite quail using the EEC of 53.5 mg a.i./kg diet (Wild Bird Scenario 2). The refined scenario for bobwhite quail resulted in a margin of safety of 2.8, which is interpreted as a low risk of reproductive effects. For mallard ducks, the NOEL also exceeds the EECs of 25 and 9.5 mg a.i./kg diet from Wild Bird Scenarios 1 and 2, respectively. The resultant margins of safety are 6.0 and 16 for Wild Bird Scenarios 1 and 2, respectively (Appendix IV, Table 5). The margin of safety for Wild Bird Scenario 2 indicates a negligible risk of reproductive effects occurring in mallard duck following long-term dietary exposure to fenbuconazole.

6.4.2.3 Wild mammals

Wild mammals such as rats and mice could be exposed to residues of fenbuconazole as a result of the consumption of sprayed vegetation and/or contaminated prey. From Table 5.9.4, assuming no transformation and no interception of orchard blast spray (Wild Mammal Scenario 1, see Section 5.9.3), the EECs of fenbuconazole in the diets of rats, mice, and rabbits are 371, 369, and 554 mg a.i./kg dw, respectively. When transformation on vegetation and interception by trees are taken into account (Wild Mammal Scenario 2, see Section 5.9.3), these EECs decrease to 50, 54, and 170 mg a.i./kg dw, respectively.

For rats, body weights per individual (BWI) of 0.192 and 0.157 kg and food consumptions (FC) of 0.028 and 0.020 kg dw per individual rat were used for males and females, respectively. Therefore, the daily intakes (DI = FC × EEC) of fenbuconazole are 10 and 7.4 mg a.i./ind/d for males and females, respectively, under Wild Mammal Scenario 1, and 1.4 and 1.0 mg a.i./ind/d for males and females, respectively, under Wild Mammal Scenario 2. Two acute oral toxicity studies were reviewed by the Health Evaluation Division (HED): one for the active ingredient and one for the formulated end-use product. The LD₅₀s in these studies were 5000 and 4000 mg a.i./kg bw for the active and end-use product, respectively. Expressed on a per individual basis, the LD_{50(ind)}s (LD₅₀ × BWI) are 960 and 785 mg a.i./ind for males and females, respectively, in the active ingredient study and 768 and 628 mg a.i./ind for males and females, respectively, in the end-use product study. As NOELs were not available for either study, one-tenth of the LD₅₀ was used as the NOEL. The calculated NOELs are, therefore, 500 and 400 mg a.i./kg bw and the NOEL_(ind)s (NOEL × BWI) are 96.0 and 78.5 mg a.i./ind for males and females, respectively, in the active ingredient study, and 76.8 and 62.8 mg a.i./ind for males and females, respectively, in the end-use product study.

Using the data from the oral toxicity study with the active ingredient, the daily intakes from Wild Mammal Scenarios 1 and 2, and the LD₅₀ of individual rats, it would take more than 92 and 106 days of continuous feeding ($LD_{50(ind)} \div DI$) under Wild Mammal Scenario 1, and more than 686 and 785 days of continuous feeding under Wild Mammal Scenario 2, for wild male and female rats, respectively, to attain a dose equivalent to that administered by gavage in the laboratory that killed 50% of the laboratory population. From study with the end-use product, it would take more than 74 and 85 days of continuous feeding under Wild Mammal Scenario 1, and more than 549 and 628 days of continuous feeding under Wild Mammal Scenario 2, for wild male and female rats, respectively, to attain a dose equivalent to that administered by gavage in the laboratory that killed 50% of the laboratory population.

As the NOELs used in the risk assessment are one-tenth of the LD₅₀s, the maximum number of days of intake of fenbuconazole by a wild rat to attain a dose equivalent to that administered by gavage in the laboratory that had no observable effect on the laboratory population is also one-tenth of the number of days of intake to accumulate a dose equivalent to that administered by gavage that killed 50% of the laboratory population. Thus, from the active ingredient study, the maximum number of days of intake to reach the laboratory dosage that had no observable effect is 9.2 and 11 days for male and female rats, respectively, under Wild Mammal Scenario 1, and 69 and 79 days for male and female rats, respectively, under Wild Mammal Scenario 2. Similarly, from the end-use product study, the maximum number of days of intake to reach the laboratory dosage that had no observable effect is 7.4 and 8.5 days for male and female rats, respectively, under Wild Mammal Scenario 1, and 55 and 63 days for male and female rats, respectively, under Wild Mammal Scenario 2. The results from the most sensitive rat (males in end-use product study) are presented in Appendix IV, Table 5.

For mice, one acute oral toxicity study was reviewed. Body weights per individual (BWI) of 0.031 and 0.023 kg were used for males and females, respectively. Using a mean FC rate of 0.006 kg dw per individual mouse per day, the daily intake ($DI = FC \times EEC$) of fenbuconazole is 2.2 or 0.3 mg a.i./ind/d for Wild Mammal Scenario 1 and the refined Wild Mammal Scenario 2, respectively. The LD_{50} was 5000 mg a.i./kg bw; when expressed on a per individual basis, the $LD_{50(ind)}$ ($LD_{50} \times BWI$) is 155 and 115 mg a.i./ind for males and females, respectively. Based on the daily intakes and the LD_{50} of individual mice, it would take more than 70 and 52 days of continuous feeding ($LD_{50(ind)} \div DI$) under Wild Mammal Scenario 1, or 478 and 355 days under Wild Mammal Scenario 2 for wild male and female rats, respectively, to attain a dose equivalent to that administered by gavage in the laboratory that killed 50% of the laboratory population. As for the rat studies, since the NOEL was not available, one-tenth of the LD_{50} was used in the risk assessment of the acute toxicity to mice. The calculated NOEL is, therefore, 500 mg a.i./kg bw and the $NOEL_{S(ind)}$ ($NOEL \times BWI$) are 15.5 and 11.5 mg a.i./ind for males and females, respectively. Therefore, the maximum numbers of days of intake of fenbuconazole by a wild rat to attain a dose equivalent to that administered by gavage in the laboratory with no observable effect on the laboratory population are 7.0 and 5.2 days under Wild Mammal Scenario 1, and 48 and 36 days under Wild Mammal Scenario 2, for males and females, respectively. The results from the most sensitive mouse (females) are presented in Appendix IV, Table 5.

Based on the above assessments, application of fenbuconazole at the maximum proposed label rate poses a negligible acute risk to populations of wild mammals that are exposed to fenbuconazole on vegetation in their diet.

In the dietary studies with male and female rats, the most sensitive NOEC was 20 mg a.i./kg dw (3-month study, male rats). Using an EEC of 371 mg a.i./kg dw (Wild Mammal Scenario 1), the margin of safety is 0.05, which indicates a high risk to rats. A refined assessment was performed using the EEC of 50 mg a.i./kg dw from Wild Mammal Scenario 2. In the refined scenario, which takes into account transformation and interception of the air blast spray by orchard trees, the margin of safety is 0.40 indicating moderate risk to rats (Appendix IV, Table 5).

A similar assessment was performed for the dietary studies with male and female mice. The most sensitive NOEC was 10 mg a.i./kg dw for both male and female mice (78-week study). Using an EEC of 369 mg a.i./kg dw (Wild Mammal Scenario 1), the margin of safety is 0.027, which indicates a high risk to mice. A refined assessment was performed using the EEC of 54 mg a.i./kg dw from Wild Mammal Scenario 2. In the refined scenario, the margin of safety is 0.19 indicating moderate risk to mice (Appendix IV, Table 5).

From reproductive studies with rats, the most sensitive NOEC was 80 mg a.i./kg dw (for both male and female systemic toxicity and female reproductive toxicity). Using an EEC of 371 mg a.i./kg dw (Wild Mammal Scenario 1), the margin of safety is 0.22, which indicates a moderate reproductive risk to rats. A refined assessment using the EEC of

50 mg a.i./kg dw from Wild Mammal Scenario 2 was performed. From the refined scenario, the margin of safety is 1.6, which indicates a low reproductive risk to rats (Appendix IV, Table 5).

Based on the studies with rats and mice, fenbuconazole may pose dietary and reproductive risks to mammals in the wild.

6.4.2.4 Non-target terrestrial plants

Three herbicidal screening studies were submitted on the phytotoxicity of fenbuconazole to several species of terrestrial plants. No effect was observed in the Tier I testing at a rate of 750 g a.i./ha. As this exceeds the maximum cumulative application rate of 735 g a.i./ha, the risks to terrestrial vascular plants are expected to be negligible.

6.4.2.5 Summary of risk to terrestrial organisms

An assessment of the environmental safety associated with the use of fenbuconazole has identified risks to avian and mammalian species. The most sensitive endpoints are reported in Appendix IV, Table 5. Using the proposed pattern of seven applications per year at a maximum rate of 0.105 kg a.i./ha and a refined risk assessment, fenbuconazole poses a low reproductive risk to upland game birds and wild mammals and a moderate dietary risk to wild mammals. Note that the risk assessments for wild birds and mammals took into account both a reduced amount of fenbuconazole reaching the ground due to interception by orchard trees and transformation on the vegetation consumed by the birds and mammals (e.g., grasses, leaves, leafy crops, forage crops, grains, seeds, and fruit). Thus, the dietary and reproductive risks may be greater if these two factors are not taken into account. These dietary and reproductive risks are attributed to the systemic and reproductive toxicity of the parent compound to birds and mammals.

Negligible acute risk is posed to earthworms, bees, wild birds, and wild mammals. The absence of data means the risks to beneficial predators and parasites cannot be determined at this time. The risks to terrestrial vascular plants are expected to be negligible. As fenbuconazole is persistent in soil, natural soil fungi and their processes on decomposition may be affected; however, the absence of data means that the degree to which fungal processes may be affected cannot be assessed. Furthermore, the risks to terrestrial organisms resulting from exposure to major transformation products of fenbuconazole are unknown.

6.4.3 Aquatic organisms

As for terrestrial organisms, data were not submitted on the toxicity of transformation products of fenbuconazole to aquatic organisms. Therefore, the aquatic risk assessment is for the parent compound, fenbuconazole. The degree of risk to aquatic organisms was classified according to the classification scheme identified in Section 6.4.2.

6.4.3.1 Non-target freshwater invertebrates

One accepted study was submitted on the acute toxicity of fenbuconazole to *Daphnia magna*. The 48-h EC₅₀ was 2.3 mg a.i./L. Therefore, according to the U.S. EPA classification scheme, fenbuconazole is classified as moderately toxic to daphids. The EEC of fenbuconazole in water (0.05 mg a.i./L) is below the 48-h NOEC of 0.78 mg a.i./L from the acute study. The margin of safety is 16; therefore, fenbuconazole poses a negligible acute risk to pelagic freshwater invertebrates at the proposed application rate (Appendix IV, Table 6). Note, however, that this result cannot be generalized to all freshwater invertebrates as fenbuconazole partitions to sediment where it may accumulate.

Although a study of the subchronic toxicity of fenbuconazole to *Chironomus riparius* was reviewed, at the present time, the toxicity to benthic species cannot be accurately assessed as current risk assessment methods do not allow a determination of the EECs in sediment and pore water.

One valid study was submitted to illustrate the chronic toxicity of fenbuconazole to *Daphnia magna*. In a chronic life cycle toxicity test, 21-d NOEC and LOEC for number of young per adult and adult growth was 0.078 mg a.i./L.

6.4.3.2 Non-target marine invertebrates

One acceptable study was submitted on the acute toxicity of fenbuconazole to *Mysidopsis bahia*, a pelagic marine crustacean. The 96-h LC₅₀ was 0.63 mg a.i./L. Therefore, according to the U.S. EPA classification scheme, fenbuconazole is classified as highly toxic to mysids. The EEC of fenbuconazole in water (0.05 mg a.i./L) is below the 96-h NOEC of 0.16 mg a.i./L. The margin of safety is 3.2; therefore, fenbuconazole poses a low risk to mysids at the proposed application rate (Appendix IV, Table 6).

A study on the acute toxicity of fenbuconazole to the Eastern Oyster, *Crassostrea virginica*, was accepted. An EC₅₀ was not determined. Based on the 96-h LOEC of 0.69 mg a.i./L, fenbuconazole is classified as highly toxic to this marine mollusk according to the U.S. EPA classification scheme. The EEC of fenbuconazole in water (0.05 mg a.i./L) is below the 96-h NOEC of 0.53 mg a.i./L. The margin of safety is 11; therefore, fenbuconazole poses a negligible acute risk to *C. virginica* at the proposed application rate (Appendix IV, Table 6).

As for freshwater organisms, the results for mysids and mollusks cannot be generalized to all marine invertebrates as fenbuconazole partitions to sediment where it may accumulate and benthic species may be exposed.

6.4.3.3 Fish

Freshwater

Two acceptable studies were submitted on the acute toxicity of fenbuconazole to freshwater fish. For coldwater fish (*Onchorynchus mykiss*, rainbow trout), the 96-h LC₅₀ was 1.4 mg a.i./L. Therefore, according to the U.S. EPA classification scheme, fenbuconazole is classified as moderately toxic to coldwater fish. The EEC of fenbuconazole in water (0.05 mg a.i./L) is below the 96-h NOEC of 0.7 mg a.i./L for rainbow trout. The margin of safety is 14; therefore, fenbuconazole poses a negligible risk to coldwater fish at the proposed application rate (Appendix IV, Table 6).

For warmwater fish (*Lepomis macrochirus*, bluegill sunfish), the 96-h LC₅₀ was 0.68 mg a.i./L. Therefore, according to the U.S. EPA classification scheme, fenbuconazole is classified as highly toxic to warmwater fish. The EEC of fenbuconazole in water (0.05 mg a.i./L) is below the 96-h NOEC of 0.42 mg a.i./L for bluegill sunfish resulting in a margin of safety of 8.4. Therefore, fenbuconazole poses a low risk to warmwater fish at the proposed application rate (Appendix IV, Table 6).

Two early-life stage toxicity tests with fathead minnow (*Pimephles promelas*) were submitted but only one study was found to be acceptable. In the accepted study, various endpoints were affected including wet weight, standard length, spinal curvature, and erratic swimming behaviour. The NOEC for the most sensitive endpoint (wet weight at 30 days post-hatch) was 0.082 mg a.i./L.

One whole life-cycle toxicity study with fathead minnow (*Pimephles promelas*) was submitted and accepted. In this study, various endpoints were affected including time to first spawn, number of eggs produced, survival of parents and offspring, and number of eggs per spawn. The NOEC for the most sensitive endpoint (time to first spawn) was 0.027 mg a.i./L.

The EEC of fenbuconazole in water (0.05 mg a.i./L) is less than the early-life-stage NOEC of 0.082 mg a.i./L resulting in a margin of safety of 1.6. Therefore, fenbuconazole may pose a low risk to early-life stages of freshwater fish at the proposed application rate (Appendix IV, Table 6). A study of the bioconcentration of fenbuconazole in fish indicates fenbuconazole may accumulate in tissues (BCFs of 330, 170, and 50 in viscera, whole fish, and fillets, respectively).

Marine/estuarine

One acceptable study on the toxicity to marine/estuarine fish (*Cyprinodon variegatus*, sheepshead minnow) was submitted. The 96-h LC₅₀ was 1.8 mg a.i./L. Therefore, according to the U.S. EPA classification scheme, fenbuconazole is classified as moderately toxic to marine/estuarine fish. The EEC of fenbuconazole in water (0.05 mg a.i./L) is below the 96-h NOEC of 0.89 mg a.i./L for sheepshead minnow, resulting in a margin of safety of 18. Therefore, fenbuconazole poses a negligible risk to marine/estuarine fish at the proposed application rate (Appendix IV, Table 6).

6.4.3.4 Algae

Freshwater

Three studies on the phytotoxicity of fenbuconazole to freshwater green alga (*Scenedesmus subspicatus* and *Selenastrum capricornutum*) were submitted, of which two were acceptable for review—one study for each species. One study on the toxicity to a diatom (*Navicula pelliculosa*) and one study on the toxicity to blue-green algae (cyanophyta, *Anabaena flos-aquae*) were also reviewed.

The most sensitive species was *S. capricornutum*. The EEC of fenbuconazole in water (0.05 mg a.i./L) is below the 120-h NOEC of 0.27 mg a.i./L. This results in a margin of safety of 5.4. Therefore, fenbuconazole poses a low risk to freshwater algae at the proposed application rate (Appendix IV, Table 6).

Marine

As data were not submitted on the toxicity to marine species of algae, the risk to marine algae cannot be assessed at this time.

6.4.3.5 Aquatic vascular plants

As the toxicity study with the aquatic vascular plant *Lemna gibba* was found to be unacceptable, the risk of fenbuconazole to aquatic vascular plants has not been determined at this time.

6.4.3.6 Summary of risk to aquatic organisms

Several areas of concern have been identified for aquatic species following an assessment of the environmental safety associated with the use of fenbuconazole (Appendix IV, Table 6). Using the proposed pattern of seven applications per year at a maximum rate of 0.105 kg a.i./ha, fenbuconazole poses a low acute risk to warmwater fish, freshwater algae, and marine invertebrates. There is a low risk of effects in early-life stages of freshwater fish following exposure to fenbuconazole. Furthermore, the risk posed to freshwater benthic invertebrates (e.g., chironomids) based on exposure to fenbuconazole in water (i.e., pore water) or sediment has not been determined. Risks to benthic species may be present as a result of the partitioning of fenbuconazole to sediments. Although data were not submitted on the risk to marine benthic species, this risk may be expected to exceed that of the pelagic species of marine organism that was reviewed (mysid) because of the persistence of fenbuconazole in sediments. Note, however, that exposure to marine organisms is expected to be limited in Canada as the areas in which stone fruit are grown are distant from marine waters. The potential risk to aquatic vascular plants has not been determined at this time. As for terrestrial organisms, the risks to aquatic organisms resulting from exposure to major transformation products of fenbuconazole are unknown.

6.5 Risk mitigation

Environmental concerns

Based on the data submitted and on the existing data requirements for Use Site Category 14 (Terrestrial Food Crops), an assessment of the environmental safety associated with the use of fenbuconazole has identified several areas of concern. Application of fenbuconazole using the proposed pattern of 7 applications per year at a maximum rate of 0.105 kg a.i./ha will pose a potential risk to the following organisms:

Low Risk

- Upland game bird (Bobwhite quail)—Reproduction
- Wild mammals (rats)—Reproduction
- Bluegill sunfish—Acute
- Fathead minnow—Early-life stage
- Freshwater algae—Acute
- Marine invertebrates (Mysid)—Note that, since the risk from exposure to fenbuconazole in marine sediment has not been assessed, the risk to benthic species may exceed that for the pelagic species of marine invertebrate that was reviewed.

Moderate Risk

- Wild mammals (rats, mice)—Dietary

Unknown Risk

- Predatory arthropod (beneficial predator)
- Parasitic arthropod (beneficial parasite)
- Freshwater benthic invertebrates (chironomids)
- Aquatic vascular plants

Fenbuconazole is a persistent compound and limited environmental toxicity data have been reviewed. Very high amounts of carryover are expected. The PMRA recommends that this product not be applied in consecutive years to reduce the environmental loading of fenbuconazole. Toxicological concerns with the major transformation products are unknown.

Any further use expansion beyond stone fruit-growing regions (e.g., Okanagan Valley, British Columbia, or the Niagara Region, Ontario) will require a re-evaluation of drinking water concentrations.

Buffer zones

Based on the proposed application rates, buffer zones to protect sensitive terrestrial and aquatic habitats are not required.

Additional required label statements

Under the ENVIRONMENTAL HAZARDS section, the statement “This product is toxic to fish.” (“*highly toxic*” on the label for the TGAI) on the labels for the technical active ingredient and end-use product must be replaced with the following:

This product is toxic to fish, aquatic invertebrates, and algae.

On the label for the technical active ingredient, under the ENVIRONMENTAL HAZARDS section, delete the phrase “*unless in accordance with the permitting authority*” at the end of the second sentence so that the modified sentence reads as follows:

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters.

On the label for the technical grade active ingredient, under the ENVIRONMENTAL HAZARDS section, delete following statement:

Do not discharge effluent containing this product into sewer systems without previously notifying the sewage treatment plant authority.

Based on an orchard use pattern and absence of data on the risks to beneficial predators and parasites, the following statement must be added to the ENVIRONMENTAL HAZARDS section of the label for the end-use product:

The toxicity to beneficial predators and parasites has not been assessed.

Based on the persistence of fenbuconazole and high potential for carryover, which may detrimentally affect organisms, the following statement must be added to the ENVIRONMENTAL HAZARDS section of the label for the end-use product:

Fenbuconazole is persistent and will carry over; it is recommended that this product not be used in areas treated with INDAR 75WSP FUNGICIDE during the previous season.

7.0 Efficacy

7.1 Effectiveness

7.1.1 Intended use

Indar 75WSP is proposed for application to stone fruits (USC 14) to control blossom blight and fruit brown rot of apricot, tart cherry and sweet cherry, nectarines, peaches and plums, as well as leaf spot of tart cherry and sweet cherry, scab of peach, and black knot of tart cherry and plum. The proposed use rate is 140 g product/500 L water/ha applied by

ground (usually airblast) sprayer, with up to two sprays at early red bud through bloom stage, and a third cover spray pre-pick if needed. In addition it can be applied to fruit up to time of harvest but not post-harvest, and may be applied to foliage after harvest for leaf spot only. Spray water volume varies considerably with size of trees and is calculated case-by-case to give optimum coverage while maintaining product rate per hectare.

Indar 75WSP should be first applied prior to infection. It is proposed at spray intervals of 7 to 14 days depending on disease pressure, and can be applied up to day of harvest. The applicant has indicated that there can be up to eight spray dates per season but the typical number of applications is two per disease (see Table 7.1).

Table 7.1. Proposed spray schedule for use of Indar on stone fruit in Canada

Crop/ Disease	Bloom	Shuck	Cover I	Cover II	Cover III	Pre-harvest I	Pre-harvest II	Post-harvest (leaves)	Typical no. of sprays
Apricot/ nectarine/peach Blossom blight Fruit brown rot	X	X				X	X		12
Cherries, sweet and tart Blossom blight Fruit brown rot Leaf spot	X	X	X X	X X	X X	X	X	X	120
Cherries, tart Black knot		X	X	X	X				2
Peach Scab		X	X	X					0
Plums Fruit brown rot Black knot	X	X	X	X	X	X	X		22

7.1.2 Mode of action

Indar 75WSP contains 75% fenbuconazole, a sterol inhibitor (Group 3) fungicide. Active ingredients in this group interfere with production of ergosterol for fungal membranes. Treated fungal spores will germinate but vegetative growth does not occur. Fenbuconazole is rapidly absorbed by green tissues and translocated acropetally.

7.1.3 Nature of pest problem

Blossom blight and fruit rot are a widespread and significant problem on stone fruit crops. In Canada, they are really two phases of a disease caused by the same or similar

pathogens (*Monilinia* spp.), first causing infection of blossoms and twigs, and later decay of ripening fruit or harvested fruit, with up to 100% loss. Blossom blight symptoms are prevalent on peach, nectarine and apricot but less common on cherry or plum. Infected plant parts, both on and off the tree, become a source of further inoculum which spreads during warm, wet weather. Fungicides are needed to reduce initial inoculum on blossoms, to protect fruit from later infection and to prevent rot in storage. The different crops each have their own critical period when coverage is most important, and only sweet cherries may require weekly sprays for the whole season. Fungicides are used primarily to control this disease, with additional sprays to control other diseases only if these are likely to be severe.

Black knot of sour cherry and plum is caused by *Dibotryon morbosum* and affects American, Japanese and European plum varieties as well as wild plums and cherries. Infection of shoots, twigs and branches occurs when spores are released in wet weather between early April and June. Lesions develop into cankers which may take two years to produce more spores. Cankers appear as black outgrowths resulting in progressive girdling of branches, death of parts of the tree and yield loss. Black knot is prevalent in eastern Canada.

Leaf spot of cherry can cause defoliation before or after fruit, reduced flower and fruit production, and cause poor fruit quality. Subsequent yield losses may be 80–100%. Scab of peach causes superficial spots on the fruit, reducing marketability. These are less critical diseases in Canadian production than blossom blight, fruit brown rot or black knot.

7.1.4 Effectiveness against pest

Five reports of Canadian brown rot trials with Indar products (75WSP or 240 g/L flowable) on peach, nectarine, or sour cherry were provided. Blossom blight was not assessed directly but is reflected by the level of brown rot. In the untreated check, up to 17% of fruit on the tree showed symptoms, while after storage for 5–11 days 41–96% of fruit was affected. Indar provided 93–98% control of disease in stored fruit when applied alone (4–7 applications), and up to 83% when applied in a mixed program. Efficacy of Indar was consistently as good as or better than commercial standards including myclobutanil and is expected to have similar activity against brown rot on all stone fruit. These data support the application of Indar 75WSP at a rate of 105 g a.i./ha to stone fruit beginning at early bloom in a program of 4 to 7 sprays, alone or alternated with other protectant fungicides.

One black knot trial on European plum and one on sour cherry were provided. In these trials, Indar was applied to inoculated trees at 100 or 135 g a.i./ha in May and June (4–5 sprays) before symptoms developed. The percentage of shoots infected with black knot was assessed the following spring. Indar provided 82–98% control, similar to Bravo and Captan, compared with 20–41% infected shoots in the untreated check.

Claims for cherry leaf spot and peach leaf scab were withdrawn by the applicant. As cherry leaf spot was the only disease requiring post-harvest sprays to trees, there is no need for this final (8th) application. The spray interval of up to 14 days is also not applicable. Claims of control of blossom blight, brown rot, and black knot are accepted as proposed at 140 g a.i./ha.

In efficacy trials, initial sprays were described as percent bloom; however, more specific growth stages are used on the proposed label, consistent with the U.S. label and presumably familiar to Canadian growers. There were insufficient comparative trials to determine a lowest effective rate; however, the proposed rate was shown to be effective for the 75WSP formulation and is consistent with the rate used commercially in the U.S.

From an efficacy standpoint, there is no restriction on the number of applications per season; however, Indar 75WSP would typically be alternated with other products, resulting in a use pattern of up to 4 sprays per season, and would provide benefit wherever it is included in the spray program. Since data were not submitted to confirm that a final spray on harvest day is necessary for control, a longer pre-harvest interval would also be acceptable.

7.2 Phytotoxicity to target plants (including different cultivars), or to target plant products (OECD 7.4)

In field trials, slight phytotoxicity to peach blossoms was noted at one site where Indar 75WSP plus surfactant were applied around a time of near-frost conditions in April. This did not affect fruit set or yield. The product has apparently been used in Europe and the U.S. without notable crop damage.

7.3 Observations on undesirable or unintended side effects, e.g., on beneficial and other non-target organisms, on succeeding crops, other plants, or parts of treated plants used for propagating purposes (e.g., seeds, cuttings, runners) (OECD 7.5)

Indar may be compatible with the biocontrol insects and arthropods used in orchards. It has been shown to have low toxicity to honeybees and was screened against five insect species (from Lepidoptera, Coleoptera, Homoptera), as well as the two-spotted spider mite (*Tetranychus urticae*) and green peach aphid (*Myzus persicae*) with no activity as a spray or soil application. However, since it was not tested specifically against beneficial species, the impact of fenbuconazole on beneficial species in IPM programs cannot be determined at this time.

7.3.1 Impact on succeeding crops (OECD 7.5.1)

Not applicable to orchard use.

7.3.2 Impact on adjacent crops (OECD 7.5.2)

Indar was tested in orchards typically undersown with sod crops such as ryegrass, without any noted effect. Observations of other orchard crops which might be grown near stone fruit were not reported.

7.4 Economics

Approximately 5800 ha of stone fruit are grown in Ontario and British Columbia combined, which comprises 98% of the national crop (2001, Statistics Canada). Farm value in Ontario alone is \$28 million, according to the applicant. Brown rot is the major disease affecting these crops and can cause up to 100% loss. This typically requires 5–10 fungicide applications per season for control, at an estimated cost of \$40–\$142 per hectare per spray, depending on the product used. For comparable cost, Indar 75WSP will contribute to disease control and help maintain marketable quality and yields. Its value to the grower is in providing another effective disease control option.

7.5 Sustainability

7.5.1 Survey of alternatives

The stone fruit diseases are currently managed by fungicides and sanitation. Some crops are less affected and certain varieties may have disease tolerance due to time of ripening or other physiological characteristics; however, resistant varieties are generally not available.

7.5.1.1 Non-chemical control practices

Both brown rot and black knot pathogens are capable of producing large numbers of spores and therefore removing as much infected material as possible is critical to managing disease. For example, any cankers or infected shoots should be pruned out and general pruning should promote an open, ventilated canopy. Unharvested fruit should be removed from trees in autumn, when possible, thinning of fruit should be timed so that it decays quickly on the orchard floor, and cultivation under trees in spring may be used to disrupt fungal fruiting structures that develop. Culled fruit should not be piled in one area near to trees.

7.5.1.2 Chemical control practices

Registered products for these two diseases on stone fruit are: captan, iprodione, chlorothalonil, sulphur, thiophanate-methyl, triforine, myclobutanil, propiconazole, and cyprodinil (Table 10.5.1). Application rates are in the range of 125–4500 g a.i. per application (18 kg for sulphur) and a program can include up to 10 sprays of different products per season. Individual products may have limitations with respect to effectiveness, crop safety, or potential for resistance. The demethylation inhibitor

products and iprodione are favoured for blossom blight control, whereas triforine and sulphur are rarely used.

Table 7.5.1 Alternative disease control products

Active Ingredient	Example end-use products	Fungicide activity site	Application rate (a.i./ha)		Comments
			Brown rot	Black knot	
Captan	Maestro, Captan	multi-site	3–3.5 kg	3–3.5 kg	Poor rainfastness Phytotoxic on certain sweet cherry and plum varieties
Chlorothalonil	Bravo	multi-site	2.5–4.5 kg	2.5–4.5 kg	Phytotoxic with oil
Cyprodinil	Vanguard	amino acids	280–560 g		Prone to resistance Not for cherries
Iprodione	Rovral	cell division, DNA, RNA	750–875 g		Some post-infection activity Resistance found in other regions
Myclobutanil	Nova	sterol demethylation	136 g		Also controls powdery mildew Prone to resistance for other diseases Not used for brown rot
Propiconazole	Topas	sterol demethylation	125 g		Some post-infection activity Prone to resistance for other diseases

Active Ingredient	Example end-use products	Fungicide activity site	Application rate (a.i./ha)		Comments
			Brown rot	Black knot	
Sulphur	Kumulus, MicroNiasul	multisite	18 kg		Harmful to some beneficial mites, can be phytotoxic, skin irritant
Thiophanate-methyl	Senator, Easout	tubulin formation	1.22 kg		Group is prone to resistance for brown rot
Triforine	Funginex	sterol demethylation	488 g		Prone to resistance for other diseases

7.5.2 Compatibility with current management practices including IPM

Integrated pest management (IPM) practices are strongly recommended in Ontario and British Columbia. These include monitoring for inoculum of stone fruit diseases to determine the level of disease pressure. Although thresholds are not established for initial fungicide application, sampling of flowers, twigs, and fruit for brown rot symptoms can provide feedback on effectiveness of spray programs and assist in decision-making regarding further applications during the season. Some products have post-infection activity and may prevent sporulation from recent infections, but it is best not to rely on this feature and to apply all products preventatively. Indar has been assessed in combination with captan and in sequence with other registered fungicides without apparent incompatibility, provided that the water-soluble pouches are dissolved in the spray mixture before other products are added as a tankmix.

Another aspect of IPM is to alternate use of fungicide chemical groups and reduce the number of sprays per season where possible. According to the applicant, Indar 75WSP has a role in replacing one myclobutanil or triforine early spray and one captan or iprodione spray late in the season for brown rot control. By providing improved control, this may delay the need for frequent (3–7 day) applications of less effective protectants. As fenbuconazole is in the same chemical class as myclobutanil and propiconazole (sterol inhibitors), these three would not be used exclusively due to risk of promoting resistance, particularly for brown rot.

Indar has been shown to have low toxicity to honeybees and was screened against five insect species (from Lepidoptera, Coleoptera, Homoptera), as well as the two spotted spider mite (*Tetranychus urticae*) and green peach aphid (*Myzus persicae*) with no

activity as a spray or soil application. It may be compatible with the biocontrol insects and arthropods used in orchards. However, since it was not tested specifically against beneficial species, the impact of fenbuconazole on beneficial species in IPM programs cannot be determined at this time.

7.5.3 Contribution to risk reduction

Indar provides an alternative to current fungicide products, some of which may present greater hazards to human health and the environment.

7.5.4 Information on the occurrence or possible occurrence of the development of resistance

The Fungicide Resistance Action Committee (FRAC) has not reported specific recommendations for stone fruit pathogens; however, general principles of resistance management should be included on the Indar 75WSP label, as per Regulatory Directive DIR99-06. Fenbuconazole is a Group 3 fungicide (demethylation inhibitors) and may be prone to resistance in pathogens that have been exposed to other fungicides in this group, such as myclobutanil and propiconazole. It is recommended that Indar 75WSP be alternated with products from other fungicide groups, such as captan, iprodione, and cyprodinil, rather than with other Group 3 products.

The following statements are required on the Indar 75WSP label:

GROUP	3	FUNGICIDE
-------	---	-----------

For resistance management, please note that Indar 75WSP contains a Group 3 fungicide. Any fungal population may contain individuals naturally resistant to Indar 75WSP and other Group 3 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same orchards. Other resistance mechanisms that are not linked to site of action but are specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance:

Where possible, rotate the use of Indar 75WSP or other Group 3 fungicides with different groups that control the same pathogens.

Avoid application of consecutive sprays of Indar 75WSP or other fungicides in the same group in a season.

Fungicide use should be based on an IPM program that includes scouting, historical information related to pesticide use and cover crop, and considers cultural, biological, and other chemical control practices.

Monitor treated fungal populations for resistance development.

If disease continues to progress after treatment with this product, do not increase the use rate. Discontinue use of this product, and switch to another fungicide with a different site of action, if available.

Contact your local extension specialist or certified crop advisors for any additional pesticide resistance management or IPM recommendations for specific crops and pathogens.

For further information and to report suspected resistance, contact Dow AgroSciences Canada Inc. at 1-800-667-3852.

7.6 Conclusions

Indar 75WSP is effective for control of blossom blight and fruit brown rot on apricot, peach, nectarine, plum and cherry and for control of black knot on plum and cherry. Indar 75WSP may be applied at a rate of 140 g per hectare from early red bud stage up to day of harvest. The maximum number of applications is seven per season; however, for resistance management, Indar 75WSP should be alternated with products from different fungicide activity groups. For comparable cost, Indar 75WSP will contribute to disease control and help maintain marketable quality and yields. Its value to the grower is in providing another effective disease control option.

7.6.1 Summary

Table 7.6.1 Summary of label proposals and recommendations

Proposed		Recommendation (based on value assessment)
Crops/diseases	Crop Group 12 stone fruits: apricot, sweet cherry, tart cherry, nectarine, peach, chickasaw plum, damson plum, Japanese plum and fresh prune—fruit brown rot and blossom blight	As proposed
	tart cherry, Japanese plum, chickasaw plum, damson plum, and fresh prune—black knot	As proposed

Proposed		Recommendation (based on value assessment)
Rate	140 g/ha (1 pouch per 0.4 hectares) in 500 L of water or sufficient water to ensure thorough coverage	As proposed
Application method	Add a wetting agent or non-polymer spray adjuvant. Ground application only	As proposed Add contraindication for aerial
Timing of applications	Blossom blight—begin at early red bud stage before infection occurs Fruit Brown Rot—begin applications 3 weeks before harvest using a 7- to 10-day spray interval. Black knot—begin at petal fall (cherry) or white popcorn stage (plum) Apply every 7 to 10 days. Under high disease pressure or favourable weather conditions, use shortest spray interval.	As proposed
Conditions	Do not use more than 8 applications per season. May be applied up to day of harvest.	Do not use more than 7 applications per season (or as per residue data). PHI of 1 day also acceptable

8.0 Toxic Substances Management Policy considerations

During the review of **Indar 75WSP**, the PMRA has taken into account the federal Toxic Substances Management Policy¹ and has followed its Regulatory Directive DIR99-03². It has been determined that this product does not meet TSMP Track-1 criteria because of the following:

- **Fenbuconazole** is not bioaccumulative. Studies have shown that the bioconcentration factor (BCF) is 330, which is below the TSMP Track-1 cut-off criterion of $BCF \geq 5000$. Furthermore, the octanol–water partition coefficient ($\log K_{ow}$) is 3.22 ± 0.08 , which is below the TSMP Track-1 cut-off criterion of ≥ 5.0 .
- **Fenbuconazole** (technical grade) does not contain any impurity known to be a toxic microcontaminant as identified in Part 2.13.4 of DIR98-04 or any of the TSMP Track-1 substances listed in App. II of DIR99-03. Impurities of toxicological concern are not expected to be present in the raw materials nor are they expected to be generated during the manufacturing process.

The end-use product is not known to contain any U.S. EPA inert List 1 or 2 formulants or any known TSMP Track-1 substances.

9.0 Regulatory decision

The fungicide active ingredient fenbuconazole and associated end-use product Indar 75WSP Fungicide for the control of blossom blight and fruit brown rot on Crop Group 12 Stone Fruit—apricot, sweet cherry, tart cherry, nectarine, peach, plum, chickasaw plum, damson plum, Japanese plum and fresh prune; and black knot on tart cherry, plum, chickasaw plum, damson plum and fresh prune, have been granted temporary registration under Section 17 of the Pest Control Products Regulations, pending the generation of the following studies:

- a validated method for animal biota (preferably for fish) (DACO 8.2.2.4).
- studies on the toxicity to beneficial predators (DACO 9.2.5) and beneficial parasites (DACO 9.2.6)

¹ The federal Toxic Substances Management Policy is available through Environment Canada's web site at www.ec.gc.ca/toxics.

² The PMRA's *Strategy for Implementing the Toxic Substances Management Policy*, DIR99-03, is available through the Pest Management Information Service: phone 1-800-267-6315 within Canada or 1-613-736-3799 outside Canada (long distance charges apply); fax (613) 736-3798; e-mail pminfoserv@hc-sc.gc.ca or through our web site at www.hc-sc.gc.ca/pmra-arla.

The following MRLs will be recommended for promulgation in the Table II, Division 15 of the *Food and Drugs Act* and Regulations:

apricot (0.3 ppm); sweet and tart cherry (0.8 ppm); peach (0.5 ppm);
nectarine (0.5 ppm); plum, chickasaw plum, damson plum, Japanese plum,
plumcot (0.1 ppm); fresh prune (0.1 ppm); dry prune (0.5 ppm).

List of abbreviations

a.i.	active ingredient
ADI	acceptable daily intake
ALAT	alanine aminotransferase
ARfD	acute reference dose
BCF	bioconcentration factor
bw	body weight
CAS	Chemical Abstracts Service
d	day(s)
DACO	Data Code
DFR	dislodgeable foliar residue
DNA	deoxyribonucleic acid
EEC	expected environmental concentration
EPA	United States Environmental Protection Agency (U.S.)
EXAMS	Exposure Analysis Modelling System
F ₀	parental animals
F ₁	1 st generation offspring
F ₂	2 nd generation offspring
GAP	good agricultural practices
HED	Health Evaluation Division
K _{ow}	<i>n</i> -octanol–water partition coefficient
K _d	adsorption quotient
K _{oc}	adsorption quotient normalized to organic carbon
LC ₅₀	lethal concentration 50%
LD ₅₀	lethal dose 50%
LEACHM	Leaching Estimation and Chemistry Model
LOAEL	lowest observed adverse effect level
LOQ	limit of quantification
MAS	maximum average score (at 24, 48, and 72 h)
MMAD	mass median aerodynamic diameter
MOE	margin of exposure
MOS	margin of safety
MRL	maximum residue level
NOAEL	no observed adverse effect level
NOEC	no observed effect concentration
OECD	Organisation for Economic Cooperation and Development
PAI	pure active ingredient
PHED	Pesticide Handlers' Exposure Database
PHI	pre-harvest interval
pK _a	dissociation constant
PMRA	Pest Management Regulatory Agency (Health Canada)
ppm	parts per million
PRZM	Pesticide Root Zone Model
RAC	raw agricultural commodity
RNA	ribonucleic acid

SGOT	serum glutamic-oxaloacetic transaminase
SGPT	serum glutamate pyruvate transaminase
SPSF	Statement of Product Specification Form
STMR	Supervised Trial Median Residue
T3	tri-iodothyronine
T4	thyroxine
TGAI	technical grade active ingredient
TSH	thyroid stimulating hormone
TSMP	Toxic Substances Management Policy
μg	microgram
μL	microlitre
U.S.	United States of America
USC	Use Site Category

References

Studies reviewed in support of food residue chemistry:

Batra, R. 1993. RH-7592 Storage Stability in Soil, Wheat Grain, Wheat Straw, Stone Fruit, Pecans and Apples—Data to 18 months. Rohm and Haas Company. Analytical Report No. AR 34A-93-16. MRID # 42925901. Unpublished.

Batra, R. October 4, 1993. Fenbuconazole (RH-7592) Formulation Bridging Studies on Apricots, Cherries & Peaches: Zero day Treatment to Sampling Interval; RAR 93-0070, 0075, 0084, 0087, 0089. Rohm and Haas Company. Technical Report No. 34A-93-19. Unpublished.

Batra, R. 1994. RH-7592 Storage Stability in Stone Fruit—Data 30 months. Rohm and Haas Company. Analytical Report No. AR 34A-94-14. MRID # 43333501. Unpublished.

Batra, R. December 3, 1996. Fenbuconazole (RH-7592) Residues in Gerber[®] Peach Puree—Analysis of RAC, In-process Samples and Finished Product (Peach Baby Food): 95-0086. Rohm and Haas Company and Centre Analytical Laboratories, Inc. Technical Report No. 34-96-69. Unpublished.

Batra, R. December 17, 1996. Fenbuconazole (RH-7592) Formulation 75W Commercial Packing House Residues in Trials Peaches: RAR 95-0113, 95-0121, 95-0143, 95-0183. Rohm and Haas Company and Centre Analytical Laboratories, Inc. Technical Report No. 34-96-65. Unpublished.

Batra, R. and B. Cressman, October 18, 1996. RH-7592 Storage Stability in Stone Fruit. Rohm and Haas Company. Technical Report No. TR 34-96-145. Unpublished.

Batra, R. February 11, 1997. An Evaluation of Fenbuconazole Prepared for the 1997 FAO Panel of Experts on Pesticide Residues in Food and the Environment, Section 2: Metabolism and Environment Fate. Report No. TR 34-96-143. Unpublished.

Burnett, T.F. October 28, 1988. RH-7592 Parent Residues Data for Plums, RAR 90-0083, 90-0126, 90-0149. Rohm and Haas Company. Technical Report No. 34A-91-11. Unpublished.

Burnett, T.F. March 6, 1991. RH-7592 Total Residues Data for Cherry, RAR 90-0058, 90-0066, 90-0116. Rohm and Haas Company. Technical Report No. 34A-91-02. Unpublished.

Burnett, T.F. March 22, 1991. RH-7592 Total Residues Data for Peach RAR 90-0062, 90-0085, 90-0125, 90-0130, 90-0131. Rohm and Haas Company. Technical Report No. 34A-91-08. Unpublished.

Burnett, T.F. March 28, 1991. RH-7592 Total Residues Data for Fresh and Dried Prune PAR 90-0126, 90-0164, 90-0245. Rohm and Haas Company. Technical Report No. 34A-91-12. Unpublished.

Burnett, T.F. September 27, 1991. RH-7592 Residue Analytical Confirmation Method for Pecan and Stone fruit. Rohm and Haas Company. Technical Report No. 34-91-67. MRID 420535-01. Unpublished.

Burnett, T.F. April 20, 1992. RH-7592: Total Residues Data for Stone fruit (Peach, Cherry, Plum and Prune) at zero (0) Day after Treatment: RAR 87-0110, 87-0245, 87-0258, 87-0263, 87-0269, 87-0333, 87-0367, 88-0015. Rohm and Haas Company. Technical Report No. 34A-92-06. Unpublished.

Burnett, T.F., N. Ding and J. J. Martin, April 04, 1991. RH-7592 Total Residues Data for Cherries, RAR 87-0110, 87-0113, 87-0194, 87-0211, 87-0263, 87-0269, 87-0333. Rohm and Haas Company. Technical Report No. 34A-91-14. Unpublished.

Burnett, T.F., N. Ding and J. J. Martin, April 04, 1991. RH-7592 Total Residues Data for Peaches, RAR 87-0156, 87-0237, 87-0245, 87-0276. Rohm and Haas Company. Technical Report No. 34A-91-16. Unpublished.

Burnett, T.F., Ding, N. and Martin, J. J. April 04, 1991. RH-7592 Total Residues Data for Fresh Plums, RAR 87-0204, 87-0206, 87-0258, 87-0367 and 88-0055. Rohm and Haas Company. Technical Report No. 34A-91-15. Unpublished.

Chamberlin, William. March 27, 1991. FDA MultiScreen Pesticide Protocols for RH-7592, RH-9129, RH-9130 and RH-6467. Rohm and Haas Company. Technical Report No. 34-91-25 MRID # 418750-44. Unpublished.

Hawkins, D.R. September 15, 1988. Metabolism of ¹⁴C-7592 in Peaches. Rohm and Haas Company Technical Report No. 34S-88-24. MRID # 41073508. Unpublished.

Martin, John J. November 29, 1988. A Residue Analytical Method for Parent RH-7592 in Stone fruit. Rohm and Haas Company. Technical Report No. 34S-88-20. MRID # 41031239. Unpublished.

Martin, John J. August 9, 1990. Residue Analytical Method for Parent RH-7592 and its Lactone Metabolites RH-9129 and RH-9130 in Stone fruit. Rohm and Haas Company. Technical Report No. 34-90-47. MRID # 41875038. Unpublished.

Martin, John J. March 15, 1991. Independent Laboratory Validation of the Residue Analytical Method for Parent RH-7592 and its Lactone Metabolites RH-9129 and RH-9130 in Stone fruit. Rohm and Haas Company. Technical Report No. 34-91-08. MRID # 41875032. Unpublished.

Martin, John J. (Revised by Burnett, T.F.) December 8, 1993. Revised Residue Analytical Method for Parent RH-7592 and its Lactone Metabolites RH-9129 and RH-9130 in Stone fruit. Rohm and Haas Co. Technical Report No. 34-90-47R. Unpublished.

Martin, J.J. and T.F. Burnett, October 26, 1988. RH-7592 Parent Residues Data and Half-Life of Decline for Peaches, RAR 87-0237, 87-0276, 87-0156, 87-0245, 87-0170. Rohm and Haas Company. Technical Report No. 34A-88-55. Unpublished.

Martin, J.J. and T.F. Burnett, (1988) RH-7592 Parent Residues Data for Cherry, RAR 87-0194, 87-0263, 87-0269, 87-0333, 87-0110, 87-0113, 87-0211. Rohm and Haas Company. Technical Report No. 34A-88-62 (October 28, 1988). Unpublished.

Martin, J.J. and T. F. Burnett, (1988) RH-7592 Parent Residues Data for Plums, RAR 87-0206, 87-0258, 87-0204, 87-0163. Rohm and Haas Company. Technical Report No. 34A-88-69. (November 7, 1988) Unpublished.

O'Dowd, M.L. May 19, 1988. RH-7592: Residue Decline in Peaches. Rohm and Haas Company. Technical Report No. 34S-88-12. Unpublished.

Sharma, Ashok K. April 23, 1992. RH-7592 Peanut Metabolism. Rohm and Haas Company Technical Report No. 34-92-38. MRID# 42447701. Unpublished.

Sharma, Ashok K. November 18, 1993. RH-7592 Supplementary Crop Metabolism—Supplement to Rohm and Haas Technical Reports 34-92-38 (MRID 42447701), 34-89-48 (MRID 41875034) and 34S-88-24 (MRID 41073508). Technical Report No. 34-93-82. Unpublished.

Stavinski, S.S. April 2, 1991. RH-7592 Storage Stability Summary. Rohm and Haas Company. Technical Report No. TR 34-91-26. MRID # 41875039. Unpublished.

Stavinski, S.S. September 23, 1991. RH-7592 (Indar) Confirmatory Method/Interference Study. Rohm and Haas Company. Technical Report No. 34-91-66. Unpublished.

Stavinski, S. S. August 14, 1992. RH-7592 (fenbuconazole) Fungicide, Summary of Storage Stability Information. Rohm and Haas Company. Technical Report No. TR 34-92-59. Unpublished.

Stavinski, S.S., T.F. Burnett, R.O. Deakyne and J.J. Martin. March 25, 1991. RH-7592 Storage Stability in Stone fruit; Cherry, Plum and Peach. Rohm and Haas Company and Craven Laboratory, Technical Report No. TR 34-91-16. MRID # 4185037. Unpublished.

Stavinski, S.S., T.F. Burnett, R.O. Deakyne and J.J. Martin. March 29, 1991. RH-7592 Storage Stability, Aged Field Residues. Rohm and Haas Company and Centre Analytical Laboratories, Inc. Technical Report No. TR 34-91-22. MRID # 41875042. Unpublished.

Stavinski, S.S. August 30, 1993. Rohm and Haas Company Response to EPA HED and EFGWB Fenbuconazole Reviews of Stone fruit Petition No. 1F 3989 and Pecan Petition No. 1F 3995. Rohm and Haas Company. Technical Report No. TR 34-93-86. Unpublished.

Studies reviewed and references used in support of environmental assessment

Atkins, E.L., E.A. Greywood, and R.L. Macdonald. 1975. Toxicity of Pesticides and Other Agricultural Chemicals to Honey Bees: Laboratory Studies. University of California, Division of Agricultural Sciences.

Atkins, E. L., D. Kellum, and K.W. Atkins. 1981. Reducing pesticide hazards to honey bees: Mortality prediction techniques and integrated management strategies. University of California, Division of Agricultural Sciences, Leaflet 2883. p. 22.

Atkins, E.L. 1988. RH-7592 Technical: Bee Adult Toxicity Dusting Test. Performed by College of Natural and Agricultural Sciences, University of California, Riverside, CA, U.S. for Rohm and Haas Company, Spring House, PA, U.S. Rohm and Haas Report No. 88RC-0066. 13-Dec-88. DACO 9.2.4.1.

Batra, R. 1995. Indar[®] (RH-7592) Residues on Non-target Orchard Turf under Air Blast Application Method. Performed by QC, Inc., Southampton, PA, U.S. and Rohm and Haas Company, Spring House, PA, U.S. Rohm and Haas Co. Lab Memo LM 34-95-128. 11-Sep-95. DACO 8.2.3.6.

Baur, L.O., Jr. 1994. Photolysis of ¹⁴C-RH-7592 in Pond Water. Study performed by XenoBiotic Laboratories, Inc., Plainsboro, NJ, U.S. Sponsor: Rohm and Haas Company, Spring House, PA, U.S. Rohm and Haas Technical Report No. 34-94-135. 30-Sep-94. DACO 8.2.3.3.2-2.

Beavers, J.B., T. Ross, G.J. Smith, and M.J. Jaber. 1991a. RH-7592 Technical: A one-generation reproduction study with the northern bobwhite (*Colinus virginianus*). Prepared by Wildlife International Ltd., Easton, MD, U.S. Project No. 129-142. Sponsored by Rohm and Haas Company, Spring House, PA, U.S. Rohm and Haas Report No. 89RC-0081. DACO 9.6.3.1.

Beavers, J.B., T. Ross, G.J. Smith, and M.J. Jaber. 1991b. RH-7592 Technical: A one-generation reproduction study with the mallard duck (*Anas platyrhynchos*). Prepared by Wildlife International Ltd., Easton, MD, U.S. Project No. 129-143. Sponsored by Rohm and Haas Company, Spring House, PA, U.S. Rohm and Haas Report No. 89RC-0082. DACO 9.6.3.2.

Bieber, W.-D. 1990. Interim Report: Field Dissipation Study on RH-7592. (Draft). Sponsored by Rohm and Haas Deutschland GmbH. Project No. NA 89 9611. Rohm and Haas Technical Report No. NATEC-89-9611. September 1990. DACO 8.6.1.

Blakemore, C.G. and D. Burgess. 1991. Chronic Toxicity of RH 7592 to *Daphnia magna* under Flow-Through Test Conditions. Final Report No. 3814. Rohm and Haas Report No. 89RC-0084. Prepared by Analytical Bio-Chemical Laboratories, Inc. Columbia, MO. DACO 9.3.3.

Burgess, D. 1988. Acute flow-through toxicity of RH7592 Technical to *Daphnia magna*. Prepared by Analytical Bio-Chemistry Laboratories, Inc., Columbia, MO. Report No. 88RC-022. U.S. EPA Accession No. 410735-07. DACO 9.3.2.

- Burgess, D. and J.W. Blasberg. 1991. Acute toxicity of RH-7592 to *Selenastrum capricornutum* Printz. Performed by Analytical Bio-Chemistry Laboratories, Inc., Columbia, MO, U.S. for Rohm and Haas Company, Spring House, PA, U.S. ABC Laboratory Project ID: Final Report #38928. Rohm and Haas Report No. 90-RC-0110. 01-Apr-91. DACO 9.8.2.2.
- Deakyne, R.O. and S.S. Stavinski. 1991. RH-7592 (Indar) Terrestrial Field Dissipation—Final Report. Performed by Pan-Agricultural Laboratories, Madera, CA, U.S., Quality Control Laboratory, Southampton, PA, U.S., Centre Analytical Laboratories, Inc., State College, PA, U.S., and Rohm and Haas Company, Spring House, PA, U.S. Rohm and Haas Technical Report No. 34-91-64. 04-Oct-91. DACO 8.3.2.2.1.
- Deakyne, R.O. and S.S. Stavinski. 1993. RH-7592 Single Application Terrestrial Field Dissipation Study (fenbuconazole). Performed by Pan-Agricultural Laboratories, Madera, CA, U.S. Rohm and Haas Technical Report No. 34-93-35. July 1993. DACO 8.3.2.2.2.
- Dione, E. 1993. RH 57592 Technical—Acute Toxicity to Eastern Oyster (*Crassostrea virginica*) under flow-through conditions. Laboratory Report No. 93-10-4971. Rohm and Haas Report No. 93-RC-0073. Prepared by Springborn Laboratories, Inc. Wareham, MA. DACO 9.4.3.
- Douglas, M.T. and R.W.S. Hall. 1990. The algastatic activity of RH-7592 Technical. Performed by Huntingdon Research Centre Ltd., Huntingdon, Cambridgeshire, England, for Rohm and Haas Company, Spring House, PA, U.S. Rohm and Haas Company Report No. 90-RC-01111. 19-Sep-90. DACO 9.8.2.1.
- England, D.C. 1990. Acute Toxicity of RH-7592 Technical to Earthworms (*Eisenia foetida*). Performed by Analytical Bio-Chemistry Laboratories, Inc., Columbia, MO, U.S. ABC Final Report #38896. Sponsored by Rohm and Haas Company, Spring House, PA, U.S. Rohm and Haas Study No. 90RC-0112. 26-Oct-90. DACO 9.2.3.1.
- Fletcher, D.W. 1988a. 21-day Acute Oral LD₅₀ Study with RH-7592 Technical in Bobwhite Quail. Biolife Associates, Ltd., Neillsville, WI. Report No. 88RC-0021. Submitted by Rohm and Haas Company, Spring House, PA. DACO 9.6.2.1.
- Fletcher, D.W. 1988b. 8-day Acute Dietary Study with RH-7592 Technical in Bobwhite Quail. Prepared by Bio-life Associates, Ltd., Neillsville, WI. Report No. 88RC-0020. Submitted by Rohm and Haas Company, Spring House PA. DACO 9.6.2.4.
- Fletcher, D.W. 1988c. 8-day Acute Dietary Study with RH-7592 Technical in Mallard Ducklings. Prepared by Bio-life Associates, Ltd., Neillsville, WI. Report No. 88RC-0019. Submitted by Rohm and Haas Company, Spring House, PA. DACO 9.6.2.5.
- Forbis, A.D. 1987. Uptake, Depuration and Bioconcentration of 14C-RH-7592 to Bluegill Sunfish (*Lepomis macrochirus*). Analytical Bio-Chemistry Laboratories, Inc., Columbia, MO, U.S. ABC Laboratory Project ID #35932. Sponsored by Rohm and Haas Company, Spring House, PA, U.S. Rohm and Haas Technical Report No. 31S-87-15. 05-Aug-87. DACO 9.5.6.1.

- Goring, C.A.I., D.A. Laskowski, J.W. Hamaker, and R.W. Meikle. 1975. Principles of pesticide degradation in soil. In (Haque, R. and V.H. Freed, eds.) *Environmental dynamics of pesticides*. Plenum Press, New York, pp. 135–172.
- Hanauer, R. 1991. RH-7592 Aqueous Photolysis. Rohm and Haas Technical Report No. 34-90-34. 15-May-90. DACO 8.2.3.3.2.3.
- Hicks, S.L. and M.A. Muckerman. 1994. Acute toxicity of RH-57,592 Technical to *Selenastrum capricornutum* Printz (Supplement to “Acute toxicity of RH-7592 to *Selenastrum capricornutum* Printz,” ABC Laboratories’ Report #38928; Rohm and Haas Report #90-RC-0110; MRID #418750-09.). Performed by ABC Laboratories, Inc., Columbia, MO, U.S. for Rohm and Haas Company, Spring House, PA, U.S. ABC Laboratories’ Final Report #41330. Rohm and Haas Report No. 93RC-0247. 14-Dec-94. DACO 9.8.2.3.
- Hoerger, F. and Kenaga, E.E. 1972. Pesticide residues on plants: correlation of representative data as basis for estimation of their magnitude in the environment. In Coulston F and F. Korte (eds). *Global aspects of chemistry, toxicology and technology as applied to the environment*, Vol. I. Thieme, Stuttgart, and Academic Press, New York. pp. 9–28.
- Kenaga, E.E. 1973. Factors to be considered in the evaluation of the toxicity of pesticides to birds in their environment. In Coulston F. and F. Dote (eds). *Global aspects of chemistry, toxicology and technology as applied to the environment*, Vol. II. Thieme, Stuttgart, and Academic Press, New York. pp. 166–181.
- Kennedy, J.M. and R.E. Talbert. 1977. Comparative persistence of dinitroaniline-type herbicides on the soil surface. *Weed Science* **25**:373–381.
- LSS report. 1999. Part 2: Chemistry requirements for the registration of a technical grade of active ingredient (TGAI) or an integrated system product (ISP). Laboratory Services Subdivision, Pest Management Regulatory Agency, Ottawa, ON. DACO 2.
- Machado, M. 1993a. RH 7592 Technical—Acute Toxicity to mysid shrimp (*Mysidopsis bahia*) under flow-through conditions. Laboratory Report No. 93-11-5012. Rohm and Haas Report No. 93-RC-0072. Prepared by Springborn Laboratories, Inc., Wareham, MA. DACO 9.4.2.
- Machado, M.W. 1993b. RH-57,692 Technical—Acute toxicity to Sheepshead minnow (*Cyprinodon variegatus*) under flow-through conditions. Prepared by Springborn Laboratories, Inc. Wareham, MA. Report No. 93RC-0071. Submitted by Rohm and Haas Company, Spring House, PA. MRID No. 430580-01. DACO 9.5.2.4.
- MacLeod, D.A. 1994. Statistical Analysis Methods for Avian Reproduction Experiments. Technical Report Series No. 211. Canadian Wildlife Services, National Wildlife Research Centre, Environment Canada. Hull, Quebec.

- Mamouni, A. 1992. Degradation of 14C-labelled fenbuconazole in three soils. Performed by RCC Umweltchemie AG, Itengen, Switzerland. Report No. RCC Project 320264. Rohm and Haas Technical Report No. 34-92-91. 15-Oct-92. DACO 8.2.3.4.2.
- McCall, P.J., D.A. Laskowski, R.L. Swann and H.J. Dishburger. 1981. Measurement of sorption coefficients of organic chemicals and their use in environmental fate analysis. *In Test Protocols for Environmental Fate & Movement of Toxicants. Proceedings of a symposium.* Association of Official Analytical Chemists. 94th Annual Meeting, October 21–22, 1980, Washington, DC, pp. 89–109.
- McEwen, F.L. and G.R. Stephenson. 1979. *The Use and Significance of Pesticides in the Environment.* John Wiley and Sons, Inc., Toronto, p. 282.
- Mineau, P., Jobin, B., and Baril, A. 1994. A critique of the avian 5-day dietary toxicity test (LC₅₀) as the basis of the avian risk assessment. Environment Canada, Technical Report Series No. 215. 23 pp.
- Musco, V.A. 1992a. Greenhouse Plantback Test with Indar (RH-7592). Rohm and Haas Company Memo to Dr. W.S. Hurt, Spring House, PA, U.S. Rohm and Haas HER Memo No. 92-166. 26-Oct-92. DACO 9.8.4.
- Musco, V.A. 1992b. Greenhouse Plantback Test with Indar (RH-7592)—Report II. Rohm and Haas Company Memo to Dr. W.S. Hurt, Spring House, PA, U.S. Rohm and Haas HER Memo No. 92-196. 30-Nov-92. DACO 9.8.4.
- Musco, V.A. 1993. Greenhouse Plantback Test with Indar (RH-7592) —Report III. Rohm and Haas Company Memo to Dr. W.S. Hurt, Spring House, PA, U.S. Rohm and Haas HER Memo No. 93-101. 02-Aug-93. DACO 9.8.4.
- O’Dowd, M.L. 1988a. RH-7592: Hydrolysis study. Rohm and Haas Company, Spring House, PA, U.S. Rohm and Haas Technical Report No. 34S-88-05. 26-May-88. DACO 8.2.3.2.1.
- O’Dowd, M.L. 1988b. Laboratory Study of Pesticide Accumulation in Fish: RH-7592 Metabolism in Bluegill Sunfish. Rohm and Haas Company, Spring House, PA, U.S. Rohm and Haas Technical Report No. 34-88-26. 04-Oct-88. DACO 9.5.6.2.
- O’Dowd, M.L. 1990a. Amendment to RH-7592 Hydrolysis Study. Rohm and Haas Company, Spring House, PA, U.S. Supplemental Report to Rohm and Haas Technical Report No. 34S-88-05. Laboratory Project ID: Rohm and Haas Technical Report No. 34-90-55. 27-Jul-90. DACO 8.2.3.2.2.
- O’Dowd, M.L. 1990b. Supplement to TR 34S-88-26: RH-7592 Metabolism in Bluegill Sunfish. Rohm and Haas Company, Spring House, PA, U.S. Rohm and Haas Technical Report No. 34-90-14. 17-Aug-90. DACO 9.5.6.3.

-
- Rhodes, J. 2002a. Toxicity of Indar™ 75 WSP Fungicide to the freshwater diatom, *Navicula pelliculosa*. ABC Laboratories, Inc. ABC Study No. 46761. 24-Apr-02. Rohm and Haas Report No. 01RC-0155. DACO 9.8.4.
- Rhodes, J. 2002b. Toxicity of Indar™ 75 WSP Fungicide to the blue-green alga, *Anabaena flos-aquae*. ABC Laboratories, Inc. Unpublished report. ABC Study No. 46760. 08-Apr-02. Rohm and Haas Report No. 01RC-0154. DACO 9.8.4.
- Rhodes, J. 2002c. Toxicity of Indar™ 75 WSP Fungicide to the blue-green alga, *Anabaena flos-aquae*: Raw Data Package ABC Laboratories, Inc. Unpublished report. ABC Study No. 46760R. 08-Apr-02. Rohm and Haas Report No. 01RC-0154A. DACO 9.8.4.
- Rhodes, J. 2002d. Toxicity of Indar™ 75 WSP Fungicide to duckweed, *Lemna gibba* G3. ABC Laboratories, Inc. Unpublished report, ABC Study No. 46762. 14-May-02. Rohm and Haas Report No. 01RC-0156. DACO 9.8.5.
- Rhodes, J.E., K. Friesen, and W.A. McAllister. 1991. Early life-stage toxicity of RH-7592 Technical to the fathead minnow (*Pimephales promelas*) under flow-through conditions. ABC Report No. 38142. Rohm and Haas Report No. 89RC-0083. Conducted by Analytical Bio-Chemistry Laboratories, Inc., Columbia, MO. DACO 9.5.3.1.
- Rhodes, J.E. and T. Leak. 1996. Full Life-Cycle Toxicity of RH-57,592 to the Fathead minnow (*Pimephales promelas*) under Flow-Through Conditions. Prepared by ABC Laboratories, Inc., Columbia, MO. Report No. 93RC-0226. Submitted by Rohm and Haas Company, Spring House, PA. MRID No. 442158-01. DACO 9.5.3.2.
- Schieber, C. 1988a. Soil Metabolism of RH-7592. Performed by Rohm and Haas Company, Spring House, PA, U.S. Rohm and Haas Technical Report No. 34S-88-13. 26-Sept-88. DACOs 8.2.3.4.2 and 8.2.3.4.4.
- Schieber, C. 1988b. Adsorption and Desorption Study of RH-7592. Performed by Rohm and Haas Company, Spring House, PA, U.S. Rohm and Haas Technical Report No. 34S-88-06. 12-Jul-88. DACO 8.2.4.2.
- Schieber, C. 1988c. Aged Leach Study of RH-7592. Performed by Rohm and Haas Company, Spring House, PA, U.S. Rohm and Haas Technical Report No. 34S-88-09. 12-Jul-88. DACO 8.2.4.3.2.
- Specht, W. 1992. See page behaviour of Indar/Corbel (a.i. fenbuconazole) in three Standard Soils. Performed by Chemische Laboratorien GmbH, St. Anscharplatz 10, Hamburg, Germany. Project Identity: ROH-9110, Az. 88223/91. 11-Mar-92. DACO 8.2.4.3.1.
- Swigert, J.P. 1988a. Acute flow-through toxicity of RH-7592 Technical to Rainbow trout (*Salmo gairdneri*). Prepared by Analytical Bio-Chemistry Laboratories, Inc., Columbia, MO. Report No. 88RC-0025. Submitted by Rohm and Haas Company, Spring House, PA. Accession No. 410312-35. DACO 9.5.2.1.

- Swigert, J.P. 1988b. Acute flow-through toxicity of RH-7592 Technical to Bluegill sunfish (*Lepomis macrochirus*). Prepared by Analytical Bio-Chemistry Laboratories, Inc., Columbia, MO. Report No. 88RC-0024. Submitted by Rohm and Haas Company, Spring House, PA. Accession No. 410735-06. DACO 9.5.2.2.
- Sword, M.C. and J.L. Stratton. 1991. Early life-stage toxicity of RH-7592 technical to fathead minnow (*Pimephales promelas*) under flow-through conditions. ABC Report No. 39266. Rohm and Haas Report No. 91RC-0007. Study conducted by Analytical Bio-Chemistry Laboratories, Inc., Columbia, MO. DACO 9.5.3.1.
- Todt, K. 1991. Final Report: Assessment of RH-7592 Residues in Soil. Sponsored by Rohm and Haas Deutschland GmbH. Project No. NA 90 9210. January 1991. DACO 8.6.2.
- Urban, D.G. and N.J.Cook. 1986. Ecological Risk Assessment. U.S. Environmental Protection Agency, EPA 540.9-85-001. Washington, D.C. 96 pp.
- van der Kolk, J. 1995. Indar®: Chronic effects on midge larvae (*Chironomus riparius*) in a water/sediment system. Springborn Laboratories (Europe) AG. Horn, Switzerland. Rohm and Haas Report No. 94RC-0189. Rohm and Haas Company, Springhouse PA. DACO 9.3.4.
- Völkl, S. 1992. ¹⁴C-Fenbuconazole: Metabolism in aquatic systems. Performed by RCC Umweltchemie Ag, Itingen, Switzerland. Study Project No. RCC Project 313560. Rohm and Haas Technical Report No. 34-92-68. 23-Jul-92. DACO 8.2.3.5.4.
- Wang, W.W. 1991a. Soil photolysis of ¹⁴C-RH-7592. Study performed by XenoBiotic Laboratories, Inc., Princeton, NJ, U.S. Sponsor: Rohm and Haas Company, Spring House, PA, U.S. Rohm and Haas Technical Report No. 34-91-05. 07-Feb-91. DACO 8.2.3.3.1.
- Wang, W.W. 1991b. Aqueous photolysis of ¹⁴C-RH-7592. Study performed by XenoBiotic Laboratories, Inc., Princeton, NJ, U.S. Sponsor: Rohm and Haas Company, Spring House, PA, U.S. Rohm and Haas Technical Report No. 34-91-04. 07-Feb-91. DACO 8.2.3.3.2.1.

Appendix I Summary table of toxicology studies for fenbuconazole

METABOLISM—TECHNICAL			
<p>In a series of rat metabolism studies, ¹⁴C-RH-7592 (≥98% radiochemical purity) uniformly labelled in the unsubstituted phenyl ring, was administered to Sprague Dawley rats (3–5/sex/dose) as a single gavage dose at 1 or 100 mg/kg bw or as a repeated gavage dose of 14 daily doses (10 ppm in the diet) of unlabelled RH-7592 (purity 96.4%) followed by a single oral dose of 1 mg/kg bw of ¹⁴C-RH-7592 on day 15. In addition, another group of bile duct-cannulated rats (5 rats/sex) received a single gavage dose of ¹⁴C-RH-7592 at 1 mg/kg bw.</p> <p>Following oral dosing, ¹⁴C-RH-7592 was rapidly absorbed, distributed and excreted. The feces was the major route of excretion, accounting for ~76% to 94% of the administered dose. Recovery from urine ranged from ~5% to 14%. The majority of the radioactivity was excreted within 24 to 48 hours post-dosing. Biliary excretion data indicated that systemic absorption of RH-7592 was high for all dosing groups. Tissue distribution and bioaccumulation were minimal, with <1% of the administered dose recovered in tissues at 96 hours. Radioactivity in the blood peaked at 3 to 6 hours. Elimination was biphasic with an initial rapid phase (24–48 hours post-dosing) followed by a slower decline (48–96 hours post-dosing). There was no sex- or dose-related difference in absorption, distribution or elimination.</p> <p>There were many metabolites were in excreta, indicating extensive metabolic breakdown. All major metabolites were derived from enzymatic oxidations on either the benzylic carbon alpha to the chlorophenyl ring or the 3- or 4- position of the phenyl ring. Subsequent non-enzymatic cyclization of the newly formed benzylic alcohol with the adjacent nitrile group, followed by hydrolysis led to the iminolactone/lactone family of metabolites. Conjugation of the OH groups of the alcohol and phenols led to more metabolites, as did combinations of the above-mentioned reactions. A minor metabolic pathway was the cleavage of RH-7592 to give triazole and RS-5922. There was no significant difference in the total metabolite profile for male and female rats, although some metabolites were present in greater quantities in males, and vice versa. There was a dose-related difference in metabolism, with a higher amount of unmetabolized parent compound in the feces of the high-dose group, compared to the low-dose and repeated-dose groups, indicating that saturation of the metabolic pathway may be occurring at the high dose.</p>			
STUDY	SPECIES/STRAIN AND DOSES	NOAEL and LOAEL mg/kg bw/day	TARGET ORGAN/SIGNIFICANT EFFECTS/COMMENTS
ACUTE STUDIES—TECHNICAL			
Oral	Mouse—CD-1, 5/sex/group; 0 and 5000 mg/kg bw	LD ₅₀ >5000 mg/kg bw	No treatment-related finding LOW TOXICITY
Oral	Rat—CrI:CD BR, 5/sex/group; 1, 2, 3, 4, and 5 g/kg bw	LD ₅₀ >5 g/kg bw	<p>≥ 1 g/kg bw: Lower body weight gain, both sexes, week 1 only</p> <p>≥ 2 g/kg bw: Ataxia, lacrimation, salivation, passiveness, and arched back. Recovery was complete by day 8.</p> <p>5 g/kg bw: One male and 2 females died; reddened glandular stomach and enlarged adrenal glands at necropsy.</p> <p>LOW TOXICITY</p>
Dermal	Rat—CrI:CD BR, 5/sex/group; 0 and 5000 mg/kg bw	LD ₅₀ >5000 mg/kg bw	No treatment-related finding No skin irritation LOW TOXICITY

STUDY	SPECIES/STRAIN AND DOSES	NOAEL and LOAEL mg/kg bw/day	TARGET ORGAN/ SIGNIFICANT EFFECTS/COMMENTS
Inhalation	Rat—Sprague-Dawley, 5/sex; 2.1 mg/L	LC ₅₀ >2.1 mg/L	MMAD = 9.8 µm, GSD not calculated; however, 18.6% of particles < 3 µm. Clinical observations observed were apathy, hunched posture, laboured respiration, piloerection, and chromodacryorrhea. Complete recovery by study day 3. LOW TOXICITY
Skin Irritation	Rabbit—NZW, 3/sex; 0.5 g dose	MAS = 0.00/8.0	NON-IRRITATING
Eye Irritation	Rabbit—NZW, 9 males; 0.1 g dose	MAS = 0.00/110	MINIMALLY IRRITATING
Skin Sensitization (Buehler Closed Patch Method)	Guinea pig—Hartley; 10/sex in test group, 5/sex in positive and negative control groups. Test material administered 25% for induction and 20% for challenge. Positive control DNCB	Test material did not elicit any dermal reactions. No evidence of sensitization Positive control was sensitizing, demonstrating responsiveness of assay.	NOT A SENSITIZER
ACUTE STUDIES - FORMULATION (INDAR 75WP)			
Oral	Rat—CrI: CD BR, 5/sex/group; 1, 2, 3, 4, and 5 g/kg bw	Combined LD ₅₀ >4 g/kg bw (confidence limits of 3.6 and 4.5 g/kg bw).	1 g/kg bw: Lower body weight gain, study week 1, males only ≥ 2 g/kg bw: Treatment-related mortality; lower body weight gain, both sexes, week 1 only. Ataxia, lacrimation, passiveness, yellow/brown-stained ano-genital area, unthrifty appearance, reddened extremities, and arched back; recovery was complete by day 7. Necropsy findings (decadents only): reddened intestines and stomach, black/red areas attached to the mucosal surface of the stomach. LOW TOXICITY
Dermal	Rat—CrI: CD BR, 6/sex; 2 g/kg bw	LD ₅₀ > 2 g/kg bw	Reddened skin, tan stained skin, desiccation and pinpoint scabs first observed on study day 1 with complete recovery by day 11. LOW TOXICITY

STUDY	SPECIES/STRAIN AND DOSES	NOAEL and LOAEL mg/kg bw/day	TARGET ORGAN/SIGNIFICANT EFFECTS/COMMENTS
Inhalation	Rat—Sprague-Dawley, 5/sex; 4.4 mg/L	LC ₅₀ > 4.4 mg/L	MMAD = 2.62 µm, GSD = 1.76 µm. Ruffled fur, both sexes; red serous secretion from the nose and laboured breathing, females only. Complete recovery by day 3. White gray/black brown/dark red foci on the lungs (all males, 1 female). LOW TOXICITY
Skin Irritation	Rabbit—NZW, 6 males; 0.5 g dose	MAS = 0.17/8.0	Very slight erythema which cleared by 24 hours. Treatment did not result in edema. MINIMALLY IRRITATING
Eye Irritation	Rabbit—NZW, 6 males; 0.1g dose	MAS = 3.3/110	Slight conjunctival irritation, cleared by 48 hours. Treatment did not affect the cornea or iris. MINIMALLY IRRITATING
Skin Sensitization (Buehler Closed Patch Procedure)	Guinea pig—Hartley; 10/sex in test group, 5/sex in positive and negative control groups. Test material administered 20% for induction and for challenge. Positive control DNCB	Test material did not elicit any evidence of sensitization. Positive control was sensitizing, demonstrating responsiveness of assay.	NOT A SENSITIZER
SHORT TERM - TECHNICAL			
28-day dermal	Rats—CrI: CD BR 6/sex/group; No treatment (sham control), RH-57592 formulation blank, RH-57592 technical at 1.0 g a.i./kg bw/day, RH-57592 2F at 0.0625 g a.i./kg bw/day, RH-57592 2F at 0.25 g a.i./kg bw/day or RH-57592 2F at 1.0 g a.i./kg bw/day	Systemic toxicity: LOAEL could not be determined since there were no treatment-related systemic effects. NOAEL = 1.0 g a.i./kg bw/day. Dermal toxicity: LOAEL could not be determined since there were no treatment-related systemic effects. NOAEL = 1.0 g a.i./kg bw/day.	No systemic treatment-related effects at any dose level tested. Dermal findings: Acanthosis, parakeratosis, eschar or superficial exudate, and necrosis of the epidermis. These findings were attributed to the non-active ingredients of the 2F formulation and not to the active ingredient itself.

STUDY	SPECIES/STRAIN AND DOSES	NOAEL and LOAEL mg/kg bw/day	TARGET ORGAN/SIGNIFICANT EFFECTS/COMMENTS
3-month dietary	Mouse—CD-1; 10/sex/group; 0, 540, 1000, 3000 and 10 000 ppm (equal to 0, 85.59, 158.40, 465.37, and 964.01 mg/kg bw/day for males, and 0, 113.46, 201.93, 595.30, and 2014.99 mg/kg bw/day for females) Compound consumption in the 10 000 ppm group is based on data from the first 2 weeks of the study only, because of treatment-related mortality.	LOAEL = 85.59/113.46 mg/kg bw/day NOAEL could not be determined since there were treatment-related effects observed at all dose levels tested.	85.59/113.46, 158.40/201.93 and 465.37/595.30 mg/kg bw/day: Decreased body weight gain (465.37 only; males); increased liver weights; hepatocyte hypertrophy, hepatocyte vacuolation and hepatocyte necrosis; decreased triglycerides; decreased cholesterol; increased ALAT (465.37 only; males) 964.01/2014.99 mg/kg bw/day: Not tolerated. Resulted in 80% and 100% mortality for males and females, respectively, by study week 3.
3-month dietary	Mouse—CD-1; 10/sex/group; 0, 20, 60, 180, and 540 ppm (equal to 0, 3.8, 11.1, 28.6, and 99.1 mg/kg bw/day for males, and 0, 5.7, 17.6, 50.4, and 139.2 mg/kg bw/day for females).	Males: LOAEL = 28.6 mg/kg bw/day NOAEL = 11.1 mg/kg bw/day Females: LOAEL = 139.2 mg/kg bw/day NOAEL = 50.4 mg/kg bw/day	3.8/5.7 and 11.1/17.6 mg/kg bw/day: No treatment-related finding 28.6 mg/kg bw/day: Males only affected—increased liver weight, hepatocyte hypertrophy and single cell necrosis; increased SGPT 99.1/139.2 mg/kg bw/day: Increased liver weight, hepatocyte hypertrophy and vacuolation; focal/single cell necrosis; increased SGPT and SGOT
3-month dietary	Rat—CrI:CD BR;10/sex/group; 0, 20, 80, 400, and 1600 ppm (equal to 0, 1.3, 5.1, 25.3, and 103.0 mg/kg bw/day for males, and 0, 1.5, 6.3, 31.5, and 123.9 mg/kg bw/day for females)	Males: LOAEL = 5.1 mg/kg bw/day NOAEL = 1.3 mg/kg bw/day Females: LOAEL = 31.5 mg/kg bw/day NOAEL = 6.3 mg/kg bw/day	1.3/1.5 mg/kg bw/day: No treatment-related findings 5.1 mg/kg bw/day: Males only affected—hepatocyte vacuolation 25.3/31.5 mg/kg bw/day: Increased liver weight; hepatocyte hypertrophy and vacuolation 103.0/123.9 mg/kg bw/day: Decreased body weight gain (females only); increased liver weight; hepatocyte hypertrophy and vacuolation; decreased triglycerides; increased cholesterol and GGT (females only); follicular cell hypertrophy in the thyroid

STUDY	SPECIES/STRAIN AND DOSES	NOAEL and LOAEL mg/kg bw/day	TARGET ORGAN/SIGNIFICANT EFFECTS/COMMENTS
3-month dietary	Dog—Beagle; 4/sex/group; 0, 30, 100, 400, and 1600 ppm (equal to 0, 0.97, 3.30, 13.27, and 50.40 mg/kg bw/day for males, and 0, 1.05, 3.48, 13.98, and 53.27 mg/kg bw/day for females)	LOAEL = 13.27/13.98 mg/kg bw/day NOAEL = 3.30/3.48 mg/kg bw/day	0.97/1.05 and 3.30/3.48 mg/kg bw/day: No treatment-related findings 13.27/13.98 mg/kg bw/day: Increased liver weight; hepatocyte vacuolation (males only); hepatocyte hypertrophy 50.40/53.27 mg/kg bw/day: Decreased body weight gain, food intake and food efficiency; increased liver weight; hepatocyte hypertrophy; increased alkaline phosphatase; increased SGPT and GGT (females only); decreased total protein, albumin and globulin (females only); increased triglycerides (males only); marginally decreased RBC count, Hgb and HCT (females only)
1-year dietary	Dog—Beagle; 4/sex/group; 0, 15, 150, and 1200 ppm (equal to 0, 0.54, 5.2, and 47.8 mg/kg bw/day for males, and 0, 0.62, 5.2, and 46.4 mg/kg bw/day for females).	LOAEL = 47.8/46.4 mg/kg bw/day NOAEL = 5.2 mg/kg bw/day	0.54/0.62 and 5.2 mg/kg bw/day: No adverse treatment-related effects 47.8/46.4 mg/kg bw/day: Decreased body weight gain; increased liver, thyroid and adrenal weights; hepatocyte hypertrophy; hepatocyte pigment; increased alkaline phosphatase; decreased total protein and albumin

STUDY	SPECIES/STRAIN AND DOSES	NOAEL and LOAEL mg/kg bw/day	TARGET ORGAN/SIGNIFICANT EFFECTS/COMMENTS
CHRONIC TOXICITY/ONCOGENICITY—TECHNICAL			
78-week dietary	Mouse—CD-1 mice, 60/sex/group; 0, 10, 200, and 650 ppm for males (equal to 0, 1.28, 26.28, and 85.26 mg/kg bw/day) and 0, 10, 650, and 1300 ppm for females (equal to 0, 1.59, 104.64, and 208.84 mg/kg bw/day)	<p>Chronic effects Males: LOAEL = 26.28 mg/kg bw/day NOAEL = 1.28 mg/kg bw/day Females: LOAEL = 104.64 mg/kg bw/day NOAEL = 1.59 mg/kg bw/day</p> <p>Oncogenicity Males: An increased incidence of hepatocellular carcinomas was noted at 85.26 mg/kg bw/day. Females: An increased incidence of hepatocellular adenomas and combined adenomas and carcinomas was noted at 208.84 mg/kg bw/day</p>	<p>1.28/1.59 mg/kg bw/day: No adverse treatment-related effects 26.28 mg/kg bw/day (males): Increased liver weight; hepatocyte enlargement and vacuolization 85.26/104.64 mg/kg bw/day: Increased liver weight; enlarged livers (males only); hepatocyte enlargement and vacuolization 208.84 mg/kg bw/day (females): Increased liver weight; enlarged livers; hepatocyte enlargement and vacuolization</p>
2-year dietary	Rat—Sprague-Dawley, 70/sex/group; 0, 8, 80, and 800 ppm (equal to 0, 0.30, 2.91, and 29.46 mg/kg bw/day for males and 0, 0.38, 3.89, and 42.29 mg/kg bw/day for females)	<p>Chronic effects LOAEL = 29.46/42.29 mg/kg bw/day NOAEL = 2.91/3.89 mg/kg bw/day</p> <p>Oncogenicity Males: An increased incidence of thyroid follicular cell adenomas and combined thyroid follicular cell adenomas and carcinomas was noted at 29.46 mg/kg bw/day. Females: No evidence of treatment-related oncogenicity</p>	<p>0.30/0.38 mg/kg bw/day and 2.91/3.89 mg/kg bw/day: No adverse treatment-related effects 29.46/42.29 mg/kg bw/day: Decreased body weight gain (males only); increased liver and thyroid weights; increased cholesterol (females); enlarged thyroids (males); hepatocyte enlargement and vacuolization; thyroid focal cystic hyperplasia (males)</p>

STUDY	SPECIES/STRAIN AND DOSES	NOAEL and LOAEL mg/kg bw/day	TARGET ORGAN/SIGNIFICANT EFFECTS/COMMENTS
2-year dietary; supplementary study	Male rats—Sprague-Dawley, 60/sex/group; 0, 800, and 1600 ppm (equal to 0, 28.87, and 62.07 mg/kg bw/day)	<p>Chronic effects LOAEL = 28.87 mg/kg bw/day NOAEL could not be determined since there were treatment-related findings at both dose levels tested</p> <p>Oncogenicity An increased incidence of thyroid follicular cell adenomas and combined thyroid follicular cell adenomas and carcinomas was noted at 62.07 mg/kg bw/day</p>	<p>28.87 mg/kg bw/day: Decreased food efficiency; increased liver weight; hepatocyte enlargement and vacuolization; increased cholesterol</p> <p>62.07 mg/kg bw/day: Decreased body weight gain; decreased food efficiency; increased liver weight; hepatocyte enlargement and vacuolization; thyroid follicular cell hypertrophy; increased cholesterol. (Decreased plasma thyroxine, week 103; increased TSH, weeks 85 and 103)</p>

STUDY	SPECIES/STRAIN AND DOSES	NOAEL and LOAEL mg/kg bw/day	TARGET ORGAN/SIGNIFICANT EFFECTS/COMMENTS
REPRODUCTION/DEVELOPMENTAL TOXICITY—TECHNICAL			
Two-generation dietary, one litter per generation	Rat—CrI:CD BR, 25/sex/group; 0, 8, 80, and 800 ppm (equal to 0, 0.6, 5.8, and 61.3 mg/kg bw/day for males, and 0, 0.6, 6.4, and 66.4 mg/kg bw/day for females)	<p>Systemic Toxicity LOAEL= 61.3/66.4 mg/kg bw/day NOAEL=5.8/6.4 mg/kg bw/day</p> <p>Reproductive toxicity Males: LOAEL could not be determined since there were no treatment-related effects at any dose level tested NOAEL = 61.3 mg/kg bw/day Females: LOAEL = 66.4 mg/kg bw/day NOAEL = 6.4 mg/kg bw/day</p> <p>Offspring toxicity LOAEL = 61.3/66.4 mg/kg bw/day NOAEL = 5.8/6.4 mg/kg bw/day</p>	<p>0.6 mg/kg bw/day: No treatment-related effects</p> <p>6.4 mg/kg bw/day (females): Slightly increased liver weight (P₂ females only; non-adverse)</p> <p>61.3/66.4 mg/kg bw/day: Mortality (females only); decreased body weight gain and food intake; increased liver weight; increased thyroid weight (males); increased adrenal weight (females); hepatocyte hypertrophy and vacuolation; thyroid follicular cell hypertrophy; hypertrophy of zona glomerulosa of the adrenal (P₁ females; P₂ males and females)</p> <p>Males: No treatment-related reproductive effect at any dose level tested</p> <p>Females: 66.4 mg/kg bw/day: Decreased number delivering; decreased litter size; decrease in the number and percentage of dams with liveborn pups; decreased gestation index; increase in the number and percentage of dams with stillborn pups; increase in number of litters with no viable offspring (P₁ only)</p> <p>61.3/66.4 mg/kg bw/day: Lower pup viability; lower pup body weight</p>

STUDY	SPECIES/STRAIN AND DOSES	NOAEL and LOAEL mg/kg bw/day	TARGET ORGAN/SIGNIFICANT EFFECTS/COMMENTS
Teratogenicity oral gavage	Female rats—CrI: CD BR, 25/group; 0, 30, 75 or 150 mg/kg bw/day.	<p>Maternal toxicity LOAEL = 75 mg/kg bw/day NOAEL = 30 mg/kg bw/day</p> <p>Developmental toxicity LOAEL = 75 mg/kg bw/day NOAEL = 30 mg/kg bw/day</p> <p>Teratogenicity LOAEL could not be determined since there were no treatment-related findings NOAEL = 150 mg/kg bw/day</p>	<p>30 mg/kg bw/day: No treatment-related effects</p> <p>75 mg/kg bw/day: Alopecia; scant feces; lower body weight and body weight gain; lower corrected body weight and body weight gain; lower gravid uterus weight</p> <p>150 mg/kg bw/day: Alopecia; scant feces; thin appearance; red vaginal discharge; lower body weight and body weight gain; lower corrected body weight and body weight gain; lower gravid uterus weight</p> <p>30 mg/kg bw/day: No treatment-related effect</p> <p>75 mg/kg bw/day: Decreased number of live fetuses; increased post-implantation loss; increase in partially or unossified sternbrae</p> <p>150 mg/kg bw/day: Decreased number of live fetuses; increased post-implantation loss; increase in resorptions; lower fetal body weight; increase in rudimentary 14th ribs; increase in partially or unossified sternbrae; increase in partially or unossified pubes</p> <p>No treatment-related teratogenic effects were noted at any dose level tested</p>

STUDY	SPECIES/STRAIN AND DOSES	NOAEL and LOAEL mg/kg bw/day	TARGET ORGAN/ SIGNIFICANT EFFECTS/COMMENTS
Teratogenicity oral gavage	Female rabbits—NZW, 20/group; 0, 10, 30, and 60 mg/kg bw/day.	<p>Maternal toxicity LOAEL = 30 mg/kg bw/day NOAEL = 10 mg/kg bw/day</p> <p>Developmental toxicity LOAEL = 60 mg/kg bw/day NOAEL = 30 mg/kg bw/day</p> <p>Teratogenicity LOAEL could not be determined since there were no treatment-related findings NOAEL = 30 mg/kg bw/day</p>	<p>10 mg/kg bw/day: No treatment-related effects</p> <p>30 mg/kg bw/day: Decreased food intake; anorexia; soft or scant feces</p> <p>60 mg/kg bw/day: Mortality; loss in body weight; decreased food intake; abortions; anorexia and soft or scant to no feces; red discharge</p> <p>10 and 30 mg/kg bw/day: No treatment-related effects</p> <p>60 mg/kg bw/day: Abortions and total resorption of litters (increased post-implantation loss)</p> <p>No treatment-related teratogenic effects were noted at any dose level tested.</p> <p>NOTE: A meaningful evaluation of soft tissue, visceral, or skeletal effects could not be conducted in the 60 mg/kg bw/day group (maternally toxic dose level) since only one litter was produced.</p>
MUTAGENICITY—TECHNICAL			
STUDY	SPECIES/STRAIN or CELL TYPE	DOSES EMPLOYED	SIGNIFICANT EFFECTS/COMMENTS
<i>Bacillus subtilis</i> , mammalian activation recombination repair assay	<i>B. subtilis</i> , strains H17 (rec+) and M45 (rec-)	625, 1250, 2500, 5 000, 10 000, and 50 000 µg/40 µL/plate, ± S9	Negative
<i>S. typhimurium</i> , mammalian activation gene mutation assay	<i>S. typhimurium</i> —TA 98, TA 100, TA 1535 and TA 1537	Assay 1: 50, 200, 500, 2000, and 5000 µg/plate, ± S9 Assay 2: 30, 50, 90, 160, and 300 µg/plate for strains TA100, TA1535, and TA1537, ± S9	Negative
<i>S. typhimurium</i> , mammalian activation gene mutation assay	<i>S. typhimurium</i> —TA 98, TA 100, TA 1535, and TA 1537	Assay 1: 20, 50, 200, 500, and 2000 µg/plate, ± S9 Assay 2: 160, 300, 500, 900, and 1600 µg/plate, + S9; 30, 50, 90, 160, and 300 µg/plate, -S9	Negative

STUDY	SPECIES/STRAIN or CELL TYPE	DOSES EMPLOYED	SIGNIFICANT EFFECTS/COMMENTS
<i>S. typhimurium</i> , mammalian activation gene mutation assay	<i>S. typhimurium</i> —TA 98, TA 100, TA 1535 and TA 1537	Assay 1: 50, 200, 500, 2000, and 5000 µg/plate, ± S9 Assay 2: 30, 50, 90, 160, and 300 µg/plate for strains TA1535 and TA1537, ± S9. 0.2, 0.5, 2, 5, and 20 µg/plate for strain TA98, ± S9 160, 300, 500, 900, and 1600 µg/plate for strain TA100, ± S9	Negative.
<i>E. coli</i> , mammalian activation gene mutation assay	<i>Escherichia coli</i> , strain WP2uvrA	156.25, 312.5, 625, 1250, 2500 and 5000 µg/plate, ± S9	Negative
Gene mutation assay	Chinese Hamster Ovary (CHO) cells	Assay 1: 10, 20, 30, 40, and 50 µg/mL, -S9 10, 35, 45, and 60 µg/mL, + S9 Assay 2: 15, 20, 25, 30, 35, and 40 µg/mL, -S9 30, 40, 45, 50, 55, and 60 µg/mL, +S9	Negative
Gene mutation assay	CHO cells	0, 3, 5, 10, 20, and 30 µg/mL, ± S9	Negative
Micronucleus assay, in vivo	Rat bone marrow cells	0 (vehicle control), 0.25, 1.25, and 2.5 g/kg bw, 15 rats/sex/group	Negative
Unscheduled DNA Synthesis	Rat Primary Hepatocytes	2.5, 5.0, 7.5, 10.0, 12.5, and 15.0 µg/mL	Negative
MUTAGENICITY—RH-11929			
<i>S. typhimurium</i> , mammalian activation gene mutation assay	<i>S. typhimurium</i> —TA 98, TA 100, TA 1535, and TA 1537	0, 156.25, 312.5, 625, 1250, 2500, and 5000 µg/plate, ± S9	Negative
MUTAGENICITY—RH-11930			
<i>S. typhimurium</i> , mammalian activation gene mutation assay	<i>S. typhimurium</i> —TA 98, TA 100, TA 1535 and TA 1537	0, 31.25, 62.5, 125, 250, 500, and 1000 µg/plate for strains TA100, TA1535, and TA 1537, ± S9 0, 156.25, 312.5, 625, 1250, 2500, and 5000 µg/plate for strain TA98, ± S9	Negative

STUDY	SPECIES/STRAIN or CELL TYPE	DOSES EMPLOYED	SIGNIFICANT EFFECTS/COMMENTS
SPECIAL STUDIES—TECHNICAL			
Hepatic effects, 4-week feeding study	<p>Female mice—CD-1, 10/sex/group; 0, 20, 60, 180, and 1300 ppm (equal to 0, 5.2, 13.6, 47.4, and 323.6 mg/kg bw/day)</p> <p>Positive control phenobarbital, 230.0 mg/kg bw/day</p> <p>Male rats—CrI:CD BR, 10/sex/group, 0 or 1600 ppm (equal to 0 and 130.0 mg/kg bw/day)</p> <p>Positive control phenobarbital, 86.9 mg/kg bw/day</p>	<p>Liver effects: LOAEL = 180 ppm (equal to 47.4 mg/kg bw/day) NOAEL = 60 ppm (equal to 13.6 mg/kg bw/day)</p>	<p>Mice: 5.2 and 13.6 mg/kg bw/day: No treatment-related effects 47.4 mg/kg bw/day: ↑ cytochrome P₄₅₀ (CYP2B); ↑ PROD activity. 323.6 mg/kg bw/day and phenobarbital: Increased liver weight; liver histopathology; hepatocyte proliferation (week 1 only); ↑ cytochrome P₄₅₀ (CYP2B); ↑ cytochrome b₆; ↑ PROD activity</p> <p>Rats: 130.0 mg/kg bw/day and phenobarbital: Increased liver weight; liver histopathology; ↑ cytochrome P₄₅₀ (CYP2B); ↑ cytochrome b₆; ↑ PROD activity.</p> <p>After a 6-week recovery period: Complete reversibility for all noted effects in mice and rats related to treatment with RH-7592 and phenobarbital</p>
Thyroid function and hepatic clearance of thyroxine, 13-week feeding study	<p>Male rats—CrI: CD BR, 10–20/group; 0, 8, 800, 1600, and 3200 ppm (approximately* equal to 0, 0.6, 57.9, 115.9, and 231.2 mg/kg bw/day). [* mg/kg bw/day determinations are approximate since values were calculated from data obtained from study weeks 1, 2, 3, 4, 5, 8, and 13 only.]</p>	<p>Thyroid function: LOAEL ≈ 57.9 mg/kg bw/day NOAEL ≈ 0.6 mg/kg bw/day</p>	<p>0.6 mg/kg bw/day: No treatment-related effects 57.9 mg/kg bw/day: Increased liver and thyroid weights; hyperplasia/hypertrophy of thyroid follicular cells; ↑ TSH (wk 4) 115.9 mg/kg bw/day: Decreased body weight gain; increased liver and thyroid weights; hyperplasia/hypertrophy of thyroid follicular cells; ↑ TSH (wk 4); ↓ T₄ (wk 13) 231.2 mg/kg bw/day: Decreased body weight gain; increased liver and thyroid weights; hyperplasia/hypertrophy of thyroid follicular cells; ↑ TSH (wk 4 and 13); ↓ T₄ (wk 4 and 13); ↓ rT₃ (wk 4); ↑ biliary excretion of T₄</p> <p>After a 9-week recovery period: Complete reversibility for all noted effects in rats related to treatment with RH-7592 and phenobarbital</p>

STUDY	SPECIES/STRAIN or CELL TYPE	DOSES EMPLOYED	SIGNIFICANT EFFECTS/COMMENTS
<p>Recommendation for ADI: 0.0128 mg/kg bw/day, based on the lowest NOAEL of 1.28 mg/kg bw/day in the mouse oncogenicity study, and using a 100-fold safety factor. This provides a margin of safety (MOS) of 500× for reproductive toxicity.</p> <p>Recommendation for ARfD: For females 13+ years: 0.10 mg/kg bw/day, based on the lowest NOAEL of 30 mg/kg bw/day in the rat and rabbit teratology studies, and using a 300-fold uncertainty factor. This is based on the standard uncertainty factor of 100 with an additional 3-fold uncertainty factor due to the severity of the toxicological endpoint, i.e., increased postimplantation loss and a decrease in the number of live fetuses/litter.</p> <p>There is no indication of increased susceptibility of rat or rabbit fetuses to in utero and/or postnatal exposure in the developmental and reproductive toxicity studies.</p> <p>Carcinogenicity: There was evidence of oncogenic/carcinogenic potential of fenbuconazole in rodents. For rats, there was an increased occurrence of thyroid follicular cell benign tumours and combined thyroid follicular cell benign and/or malignant tumours in males (28.87 mg/kg bw/day). For mice, this was based on the occurrence of an increased trend for malignant liver tumours in males (85.26 mg/kg bw/day) and an increase in benign liver tumours and combined benign and/or malignant liver tumours in females (208.84 mg/kg bw/day). The proposed mechanistic pathway for the thyroid tumours in rats was scientifically supported by sound mechanistic data, i.e., prolonged stimulation of the thyroid by TSH leads to chronic follicular hypertrophy/hyperplasia, progressing to follicular neoplasia of the thyroid. However, the proposed mechanistic pathway for the liver tumours in mice did not provide a convincing hypothesis. The in vitro and in vivo mutagenicity assays yielded negative results for genotoxic potential. It is therefore recommended that, for the purpose of risk characterization, a low-dose extrapolation model be applied for human risk (Q_1^*), based on the induction of liver carcinomas in male mice. The Q_1^* for fenbuconazole is 1.54×10^{-2} (mg/kg bw/day)⁻¹ in human equivalents.</p>			

Appendix II Food residue chemistry overview of metabolism studies and risk assessment

PARAMETER		PERTINENT INFORMATION																				
CHEMICAL		FENBUCONAZOLE																				
Crop	Formulation/type	Method/timing	Rate g a.i./ha	Number/ season	Maximum rate g a.i./ha	PHI (days)																
Apricots	Indar* 75 WSP Agricultural fungicide/water soluble packets	ground/at early red bud, at full bloom, and at petal-fall	105	7	735	0																
Cherries (sweet and sour)		ground/at early red bud, at full bloom, and at petal-fall																				
Nectarines		ground/at early red bud, at full bloom, and at petal-fall																				
Peaches		ground/at early red bud, at full bloom, at petal-fall, and at shuck split																				
Plums		ground/at early red bud, at white popcorn stage, at full bloom, and at petal-fall																				
LABEL RESTRICTIONS		Do not apply by air. Do not graze livestock in treated areas or feed cover crops grown in treated areas to livestock.																				
PHYSICOCHEMICAL PROPERTIES		Substance	Value																			
Water solubility, @ 20°C		Fenbuconazole (RH-7592)	3.8 ppm																			
Solvent solubility @ 25°C mg/mL			<table border="1"> <thead> <tr> <th>Solvent</th> <th>Solubility (g/0.1 L)</th> </tr> </thead> <tbody> <tr> <td>acetonitrile</td> <td>23.1</td> </tr> <tr> <td>Aromatic 200</td> <td>7.7</td> </tr> <tr> <td>cyclohexanone</td> <td>44.5</td> </tr> <tr> <td>ethyl acetate</td> <td>15.9</td> </tr> <tr> <td>ethyl alcohol</td> <td>3.9</td> </tr> <tr> <td>heptane</td> <td>0.1</td> </tr> <tr> <td>1-octanol</td> <td>1.3</td> </tr> </tbody> </table>				Solvent	Solubility (g/0.1 L)	acetonitrile	23.1	Aromatic 200	7.7	cyclohexanone	44.5	ethyl acetate	15.9	ethyl alcohol	3.9	heptane	0.1	1-octanol	1.3
Solvent	Solubility (g/0.1 L)																					
acetonitrile	23.1																					
Aromatic 200	7.7																					
cyclohexanone	44.5																					
ethyl acetate	15.9																					
ethyl alcohol	3.9																					
heptane	0.1																					
1-octanol	1.3																					
Octanol/water partition coefficient (K_{ow}) log K_{ow}		1700 ± 300 3.22 ± 0.08 (99.5% pure)																				
p K_a		N/A																				
Vapour pressure @ 25°C		0.37 × 10 ⁻⁷ mm HG																				
Relative density		N/A																				

<i>NATURE OF THE RESIDUE—ANIMAL</i>	Neither poultry nor livestock feed items are associated with the use of fenbuconazole on stone fruit. Therefore, no animal metabolism study is needed.												
<i>NATURE OF THE RESIDUE—PLANTS</i> <i>Radiolabelling positions</i> <i>Proposed Metabolic Pathway</i>	Peaches (Red Haven) Phenyl-labelled [¹⁴ C]RH-7592 and triazole-labelled [¹⁴ C]RH-7592 The metabolism of RH-7592 proceeded via three pathways. The first pathway was through an initial hydroxylation at benzylic carbon adjacent to the chlorophenyl ring, followed by ring closure to the iminolactone which readily hydrolysed to the lactone, RH-9129. The second pathway released free triazole, by oxidation of the carbon next to triazole ring, which led to the triazole conjugates, triazole alanine and triazole acetic acid. The third degradation pathway produced RH-4911 which led to the glucose conjugates, glucoside and malonyl glucoside of RH-4911.												
<i>Residue of Concern (ROC)</i>	The parent compound (RH-7592) and its lactone metabolite which exists in the form of two stereoisomers (RH-9129 and RH-9130)												
<i>RESIDUE ANALYTICAL METHOD</i>	Stone fruit MATRICES The residues of fenbuconazole and its metabolites are extracted in stone fruit using methanol. The extract is separated by filtration and then partitioned with 9.1% sodium chloride and methylene chloride. The elute is collected and evaporated to dryness, and residues are reconstituted in toluene:acetone (100:10, v:v). Further purification is achieved using silica gel and florisil column chromatography with toluene:acetone (100:30, v:v) as the eluent. The residue is collected, evaporated to dryness, redissolved in toluene:methanol (100:3, v:v) and then analysed by gas liquid chromatography using capillary column and thermionic specific detector optimized for nitrogen selectivity.												
<i>Data gathering method ID</i>	Martin, John J. (Revised by Burnett, Theodore F.) December 8, 1993. Revised Residue Analytical Method for Parent RH-7592 and its Lactone Metabolites RH-9129 and RH-9130 in Stone fruit. Rohm and Haas Co. Technical Report No. 34-90-47R. Unpublished.												
<i>Analytes</i>	RH-7592, RH-9129 and RH-9130												
<i>Instrument/detector</i>	Gas liquid chromatogram/thermionic specific detector optimized for nitrogen selectivity												
<i>Instrument parameters</i>	<table border="0"> <thead> <tr> <th>Temperature</th> <th>Flow of gases</th> <th>Bead Current</th> </tr> </thead> <tbody> <tr> <td>Column—245°C</td> <td>Air—175 mL/min</td> <td>3.3 amps</td> </tr> <tr> <td>Injector—265°C</td> <td>Hydrogen—4.5 mL/min</td> <td>(varies)</td> </tr> <tr> <td>Detector—300°C</td> <td>Helium—18 mL/min</td> <td></td> </tr> </tbody> </table>	Temperature	Flow of gases	Bead Current	Column—245°C	Air—175 mL/min	3.3 amps	Injector—265°C	Hydrogen—4.5 mL/min	(varies)	Detector—300°C	Helium—18 mL/min	
Temperature	Flow of gases	Bead Current											
Column—245°C	Air—175 mL/min	3.3 amps											
Injector—265°C	Hydrogen—4.5 mL/min	(varies)											
Detector—300°C	Helium—18 mL/min												
<i>Column</i>	SPB-608 (0.53 mm ID capillary column)												
<i>Standardization method</i>	An external standard method was used to establish retention time, response and calibration of each analyte.												
<i>Stability of primary and/or secondary standard solutions</i>	The standard solutions were refrigerated between analyses over a period ranging from two to fourteen weeks. No pattern of decline in detector response with time was noted during the storage and use of the standard solutions. The standard solutions appeared to be stable during this time.												
<i>Retention times</i>	4.18 minutes (RH-7592); 5.24 min. (RH-9129); 5.82 min. (RH-9130)												

<i>Limit of detection (LOD)</i>	The LOD was 0.01 ppm* for all analytes.
<i>Limit of quantitation (LOQ)</i>	The LOQ was established at 0.05 ppm* for all analytes.
<i>Repeatability/precision</i>	The analytical method obtained acceptable results by an independent laboratory validation (ILV) method for the determination of residues of RH-7592, RH-9129, and RH-9130 in stone fruit.
<i>Reproducibility</i>	The result of ILV indicated that analytical method (TR 34-90-47R) was reproducible.
<i>Linearity</i>	The method/detector response was linear (correlation coefficient, $r > 0.999$) within the range of 0.05–1.0 ppm for RH-7592, RH-9129, and RH-9130 in stone fruit.
<i>Specificity</i>	The chromatographic peaks were well defined and symmetrical with no apparent carryover to the following chromatograms in the area of analytical interest for both control and spiked samples.
<i>Multi-residue method</i>	Existing multi-residue methods of analysis that are currently in common usage were not found to be suitable for the determination of fenbuconazole in apricot, cherries, nectarine, peaches, plum, and prune
<i>STORAGE STABILITY DATA</i>	The residues of RH-7592, RH-9129, and RH-9130 were stable at approximately -10°C in stone fruit for more than 54.5 months. The residues were stable for storage duration of the samples from metabolism, supervised residue trials, processing, and analytical methodology studies.

<i>CROP FIELD TRIALS</i>	<p>Supervised crop field trials were conducted in the U.S. (zones 1, 1A, 2, 4, 5, 5A, 10, 11 and 12) applying five to ten times fenbuconazole (2F or Indar 75 WP) at a single rate of 56–224 g a.i./ha, up to 2.2 kg a.i./ha/season, on apricots, cherries, peaches, plums, and prunes. The treated crops were harvested at intervals between zero and twenty-one days after the final application. The proposed label rate recommends seven applications at a rate of 105 g a.i./ha for a total season rate of 735 g a.i./ha.</p> <p>The results from four trials conducted in the U.S. (zones 10 and 11) on apricots, applying six times at a rate of 140 g a.i./ha for a total of 840 g a.i./ha/season (1.14× the proposed label rate), indicated that the maximum combined residues of fenbuconazole and its metabolites in apricot, harvested at day zero following the last application, were 0.268 ppm. An MRL of 0.3 ppm is recommended to cover the expected residues of fenbuconazole (RH-7592) and its lactone metabolites (RH-9129 and RH-9130) from the proposed use of Indar 75 WSP on apricots in Canada.</p> <p>The results from nineteen trials conducted in the U.S. (zones 1, 5A and 11) on cherries, applying six times at a rate 112–140 g a.i./ha for a total of 672–1344 g a.i./ha/season (0.9–1.82× the proposed label rate), indicated that the maximum combined residues of fenbuconazole and its metabolites in cherries, harvested zero, three and seven days following the last application, were 0.749 ppm. An MRL of 0.8 ppm is recommended to cover the expected residues of fenbuconazole and its lactone metabolites from the proposed use of Indar 75 WSP on cherries in Canada.</p> <p>The results from eight trials conducted in the U.S. (zones 1, 2, 10 and 11) on peaches, applying seven to ten times at a rate of 112–224 g a.i./ha for a total of 784–1008 g a.i./ha/season (1.1–1.4 × the proposed label rate), indicated that the maximum combined residues of fenbuconazole and its metabolites in peaches, harvested on day zero following the last application, were 0.5 ppm. An MRL of 0.5 ppm is recommended to cover the expected residues of fenbuconazole and its lactone metabolites from the proposed use of fenbuconazole on peaches in Canada. The residue trials submitted for peaches are accepted as bridging data to support the use of Indar 75 WSP on nectarines. An MRL of 0.5 ppm is also recommended to cover the expected residues of fenbuconazole and its lactone metabolites from the proposed use of Indar 75 WSP on nectarines in Canada.</p> <p>The results from eight trials conducted in the U.S. (zones 1, 5A, 10, 11 and 12) on plum and prune, applying six to twelve times at a rate of 56–224 g a.i./ha for a total of 672–1008 g a.i./ha/season (0.9–1.4 × the proposed label rate), indicated that the maximum combined residues of fenbuconazole and its metabolites in plum and prune, harvested on day zero following the last application, were 0.08 ppm. An MRL of 0.1 ppm is recommended to cover the expected residues of fenbuconazole and its lactone metabolites from the proposed use of fenbuconazole on plums and fresh prunes in Canada.</p>
RESIDUE DECLINE	<p>There was some residue decline in peach after seven days of application. However, this will not affect the level of residue in stone fruit at harvest, since the proposed label suggests that Indar 75 WSP may be applied up to the day of harvest.</p>

PROCESSED FOOD	The fresh plums were processed into dry prunes. A comparison of the residues in the RAC with those in processed fruit demonstrated concentration in dry prunes. The residues of fenbuconazole and its lactone metabolites in dry prunes will be covered under the recommended MRL of 0.5 ppm.
DAIRY CATTLE FEEDING	Neither poultry nor livestock feed items are associated with the use of fenbuconazole on stone fruit. It is expected that no residue of fenbuconazole will be present in poultry and livestock food commodities as a result of this use.
PROPOSED MRLs	Total combined residues of RH-7592, RH-9129, and RH-9130 in/on apricot (0.3 ppm); cherry (0.8 ppm); peach (0.5 ppm); nectarine (0.5 ppm); plum (0.1 ppm); fresh prune (0.1 ppm); dry prune (0.5 ppm)
U.S. TOLERANCES	Residues of RH-7592, RH-9129, and RH-9130 in/on banana (0.3 ppm); blueberry (1.0 ppm); stone fruit crop group (except plums and prune) (2.0 ppm); grapefruit (0.5 ppm), grapefruit oil (35 ppm); grapefruit dried pulp (4.0); pecan (0.1 ppm)
CODEX MRLs	Residues of RH-7592, RH-9129, and RH-9130 in/on banana (0.05 ppm); cherries (1.0 ppm); cucumber (0.2 ppm); grapes (1.0 ppm); melons (except watermelon) (0.2 ppm); pecan (0.05 ppm); pomefruit (0.1 ppm); rye (0.1 ppm); summer squash (0.05 ppm); sunflower seed (0.05 ppm); wheat (0.1 ppm); wheat straw and fodder, dry (3.0 ppm)
DIETARY RISK ASSESSMENT (DRA) DEEM™ Version 7.72 1994–1998 Continuing Survey of Food Intake for Individuals	For chronic risk from dietary exposure to the residues of fenbuconazole and its metabolites from food and water, the potential daily intake (PDI) was less than 3% of the acceptable daily intake (ADI) for all population subgroups including infants, children, adults and seniors. For acute dietary intake at the 95 th percentile, exposure to the residues of fenbuconazole and its metabolites represented 1.87% of the ARfD for women 13+ years of age. The lifetime cancer risk from dietary exposure to the residues of fenbuconazole and its metabolites from food and water was estimated to be 1.72e-06 for all infants (<1 year) and children 1–6 years, and <8e-07 for the rest of the subgroups. It is expected that further refinement would result in a lifetime risk less than the level of concern of 1.00e-06.

* Analytical Chemistry Branch of U.S. EPA conducted method validation tests and noted that the sensitivity of the method under optimum operating conditions is 0.001 ppm for RH-7592 and 0.002 ppm for RH-9130 and RH-9129. The sensitivity may be increased by adjusting the current which is applied to the thermionic detector bead.

Appendix III Food residue chemistry integrated summary table

FENBUCONAZOLE		
PLANT STUDIES		
CROPS (N=1)	Peach	
ROC FOR MONITORING	RH-7592 (fenbuconazole) and its lactone metabolites, RH-9129 and RH-9130	
ROC FOR RISK ASSESSMENT	RH-7592 (fenbuconazole) and its lactone metabolites, RH-9129 and RH-9130	
DIETARY RISK FROM FOOD AND WATER		
Chronic non-cancer dietary risk ADI = 0.0128 mg/kg bw	POPULATION	ESTIMATED RISK (% of ADI)
		Median residue values; % U.S. import; % crop treated; processing factors; estimated environmental concentration for drinking water
	All infants <1 yr old	2.8
	Children 1 to 6 years	1.8
	Children 7 to 12 years	0.8
	Females 13+ years	0.5
	Males 13+ years	0.4
	Seniors 55+ years	0.6
Total Population	0.7	
Chronic cancer dietary risk Q ₁ * = 0.0154 mg/kg bw	POPULATION	ESTIMATED LIFETIME RISK
		Median/anticipated residue values; % U.S. import; % crop treated; processing factors; estimated environmental concentration for drinking water
	All infants <1 yr old	1.72×10^{-6}
	Children 1 to 6	1.72×10^{-6}
	Children 7 to 12	7.89×10^{-7}
	Females 13+	4.29×10^{-7}
	Males 13+	4.32×10^{-7}
	Seniors 55+	6.25×10^{-7}
Total Population	6.26×10^{-7}	

Acute dietary exposure analysis, 95th percentile ARfD = 0.01 mg/kg bw	POPULATION	ESTIMATED RISK (% of ARfD)
		Median residue values; % U.S. import; % crop treated; processing factors; estimated environmental concentration for drinking water
	Females 13+	1.87

Appendix IV Environmental assessment

Table 1 Fate and behaviour in the terrestrial environment

Property	Test substance	Value	Comments
Abiotic transformation			
Phototransformation on soil	RH-7592	DT ₅₀ : 79 d	Not an important route of transformation
Biotransformation			
Biotransformation in aerobic soil	RH-7592	DT ₅₀ : 285 d (silty clay loam) DT ₅₀ : 367 d (sandy loam)	This is a route of transformation, albeit a slow process. Fenbuconazole is persistent in soil under aerobic conditions.
Biotransformation in anaerobic soil	RH-7592	DT ₅₀ : 451 d (silty clay loam) DT ₅₀ : 655 d (sandy loam)	This is a route of transformation, albeit a very slow process. Fenbuconazole is persistent in soil under anaerobic conditions.

Property	Test substance	Value	Comments
Mobility			
Adsorption/desorption in soil		Adsorption K_d (mL/g): clay: 5.1 sand: 7.6 silty clay loam: 20 loam: 75 sandy loam: 115 Adsorption K_{oc} (mL/g): clay: 2200 sand: 2600 silty clay loam: 2900 loam: 5400 sandy loam: 9000	The degree of adsorption is associated with the percentage of organic matter in soil. Fenbuconazole was slightly mobile in soils with low organic carbon content (generally $\leq 1\%$) and relatively immobile in soils with higher levels of organic carbon. Desorption values were not accepted. In the soils tested, fenbuconazole was immobile in loam and sandy loam with slight to low mobility in clay, sand, and silty clay loam.
Soil leaching—aged soil		K_{oc} : ≥ 3445 mL/g (sandy loam)	Fenbuconazole has a slight potential to leach in sandy loam soil.
Field studies			
Field dissipation		DT ₅₀ for Ecoregion 9.2 (Temperate Prairies): Unable to determine to >364 d DT ₅₀ for Ecoregion 6.2 (Western Cordillera): 198 d to >364 d	Persistent under field conditions.
Field leaching	No data	—	No data

Table 2 Fate and behaviour in the aquatic environment

Property	Test material	Value	Comments
Abiotic transformation			
Hydrolysis	RH-7592	pH 5: 2210 d pH 7: 3740 d pH 9: 1340 d	Does not hydrolyze
Phototransformation in water	RH-7592	1280 d	Not a route of transformation
Biotransformation			
Biotransformation in aerobic water	Not applicable	No value	Fenbuconazole rapidly partitions to sediment; therefore, this study is not applicable.
Biotransformation in aerobic water/sediment systems	RH-7592	River water: 3.4-4.3 d River sediment: No value River system: 906 to >1000 d (loamy sand) Pond water: 1.2 d Pond sediment: No value Pond system: 442 to >1000 d (loam)	Not an important route of transformation. The fate of fenbuconazole in aerobic water/sediment systems is partitioning to sediment. Fenbuconazole was non-persistent in water and persistent in sediment.
Biotransformation in anaerobic water/sediment systems	Not applicable	—	A request for a waiver was granted as the aquatic biotransformation behaviour is described by other studies.
Partitioning			
Adsorption/desorption in sediment	Not applicable	—	No study was submitted. Fenbuconazole partitions to sediment where it is persistent.

Property	Test material	Value	Comments
Field studies			
Field dissipation	Not applicable	—	No aquatic field study was submitted.

Table 3 Effects on terrestrial organisms

Organism	Exposure	Test substance	Endpoint value	Degree of toxicity ^a
Invertebrates				
Earthworm	Acute	RH-7592	LC ₅₀ : >98 mg a.i./kg dw	Non-toxic up to 98 mg a.i./kg dw
Bee	Oral	Not applicable	—	No data
	Contact	RH-7592	LD ₅₀ : >292 µg a.i./bee NOEL not reported	Relatively non-toxic
	Brood/hive	Not applicable	—	No data
Predatory arthropod	Contact	Not applicable	—	No data
Parasitic arthropod	Contact	Not applicable	—	No data
Birds				
Bobwhite quail	Acute	RH-7592	LD ₅₀ : >2150 mg a.i./kg bw NOEL: 1470 mg a.i./kg bw	Practically non-toxic
	Dietary	RH-7592	LC ₅₀ : 4954 mg a.i./kg diet NOEC: 625 mg a.i./kg diet	Slightly toxic
	Reproduction	RH-7592	LOEC: 600 mg a.i./kg diet NOEC: 150 mg a.i./kg diet	—
Mallard duck	Acute	Not applicable	—	No data
	Dietary	RH-7592	LC ₅₀ : 2013 mg a.i./kg diet NOEC: 312 mg a.i./kg diet	Slightly toxic
	Reproduction	RH-7592	LOEC: 600 mg a.i./kg diet NOEC: 150 mg a.i./kg diet	—

Organism	Exposure	Test substance	Endpoint value	Degree of toxicity ^a
Mammals				
Refer to Section 3.				
Vascular plants				
Vascular plant	Seedling emergence	Not applicable	—	No data
	Vegetative vigour	Not applicable	—	No data
	Greenhouse study	RH-7592	NOEC: 750 g a.i./ha EC ₂₅ : >750 g a.i./ha	No phytotoxic “injury” was observed

^a Atkins et al. (1981) for bees and U.S. EPA classification for others, where applicable

Table 4 Effects on aquatic organisms

Organism	Exposure	Test substance	Endpoint value	Degree of toxicity ^a
Freshwater species				
<i>Daphnia magna</i>	Acute	RH-7592	EC ₅₀ : 2.3 mg a.i./L NOEC: 0.78 mg a.i./L	Moderately toxic
	Chronic	RH-7592	LOEC: 0.15 mg a.i./L NOEC: 0.078 mg a.i./L	—
Chironomid	Acute	Not applicable	—	No data
	Subchronic	RH-7592	EC ₅₀ for water: 0.86 mg a.i./L EC ₅₀ for sediment: 12.7 mg a.i./kg sediment NOEC for water: 0.74 mg a.i./L water NOEC for sediment: 9.1 mg a.i./kg sediment	—
Rainbow trout	Acute	RH-7592	LC ₅₀ : 1.4 mg a.i./L NOEL: 0.7 mg a.i./L	Moderately toxic
Bluegill sunfish	Acute	RH-7592	LC ₅₀ : 0.68 mg a.i./L NOEC: 0.42 mg a.i./L	Highly toxic
Fathead minnow	Chronic	RH-7592	Early life stage study NOEC: 0.082 mg a.i./L	—

Organism	Exposure	Test substance	Endpoint value	Degree of toxicity ^a
			LOEC: 0.16 mg a.i./L Whole life-cycle study NOEC: 0.027 mg a.i./L LOEC: 0.045 mg a.i./L	
Freshwater algae	Acute	RH-7592	<i>Selenastrum capricornutum</i> Printz (freshwater green algae) NOEC (120 h): 0.27 mg a.i./L EC ₂₅ (120 h): 0.39 mg a.i./L EC ₅₀ (120 h): 0.48 mg a.i./L	—
Vascular plant	Dissolved	Not applicable	—	No data
	Over-spray	Not applicable	—	No data
Marine species				
Crustacean	Acute	RH-7592	Mysid shrimp <i>Mysidopsis bahia</i> LC ₅₀ : 0.63 mg a.i./L NOEC: 0.16 mg a.i./L	Highly toxic
	Chronic	Not applicable	—	No data
Mollusk	Acute	RH-7592	Eastern Oyster, <i>Crassostrea virginica</i> NOEC: 0.53 mg a.i./L LOEC: 0.69 mg a.i./L	Highly toxic
	Chronic	Not applicable	—	No data
Fish	Acute	RH-7592	LC ₅₀ : 1.8 mg a.i./L NOEC: 0.89 mg a.i./L	Moderately toxic
	Salinity challenge	Not applicable	—	No data
Marine alga	Acute	Not applicable	—	No data

^aU.S. EPA classification, where applicable

Table 5 Risk to terrestrial organisms

Organism	Exposure	Endpoint value	EEC	MOS	Risk
Invertebrates					
Earthworm	Acute	98 mg a.i./kg soil	0.29 mg a.i./kg soil	340	Negligible risk
Bee	Oral	No data	N/A	N/A	No data
	Contact	327 kg a.i./ha	0.735 kg a.i./ha	445	Negligible risk
	Brood/hive	No data	N/A	N/A	No data
Predatory arthropod	Contact	Data required	N/A	N/A	Unknown
Parasitic arthropod	Contact	Data required	N/A	N/A	Unknown
Birds					
Bobwhite quail	Acute	1470 mg a.i./kg bw	53.5 mg a.i./kg dw	374 days*	Negligible risk
	Dietary	625 mg a.i./kg dw	53.5 mg a.i./kg dw	12	Negligible risk
	Reproduction	150 mg a.i./kg dw	53.5 mg a.i./kg dw	2.8	Low risk
Mallard duck	Acute	No data	N/A	N/A	No data
	Dietary	312 mg a.i./kg dw	9.5 mg a.i./kg dw	33	Negligible risk
	Reproduction	150 mg a.i./kg dw	9.5 mg a.i./kg dw	16	Negligible risk
Mammals					
Rat	Acute	400 mg a.i./kg bw	50 mg a.i./kg dw	55 days [†]	Negligible risk
	Dietary	20 mg a.i./kg dw	50 mg a.i./kg dw	0.4	Moderate risk
	Reproduction	80 mg a.i./kg dw	50 mg a.i./kg dw	1.6	Low risk
Mouse	Acute	500 mg a.i./kg bw	54 mg a.i./kg dw	36 days [‡]	Negligible risk
	Dietary	10 mg a.i./kg dw	54 mg a.i./kg dw	0.19	Moderate risk
	Reproduction	Not applicable	N/A	N/A	No data

Organism	Exposure	Endpoint value	EEC	MOS	Risk
Vascular plants					
Vascular plant	Screening tests	750 g a.i./hg	735 g a.i./ha	>1	Expected to be negligible risk

* For bobwhite quail acute oral toxicity (DACO 9.6.2.1), food consumption (FC) was 0.015 kg dw/ind/day, body weight per individual (BWI) was 0.204 bw/ind. Under Wild Bird Scenario 2, the EEC was 53.5 mg a.i./kg dw. Therefore, the daily intake (DI = FC × EEC) was 0.80 mg a.i./ind/day. The NOEL_(ind) (= NOEL × BWI) was 300 mg a.i./ind. The number of days for a wild population to reach the NOEL in the laboratory population was calculated as NOEL_(ind)/DI (= 374 d).

† For rat acute oral toxicity, the most sensitive rats were males in the end-use product study. Food consumption (FC) was 0.028 kg dw/ind/day, body weight per individual (BWI) was 0.192 kg bw/ind. Under Wild Mammal Scenario 2, the EEC was 50 mg a.i./kg dw. Therefore, the daily intake (DI = FC × EEC) was 1.4 mg a.i./ind/day. The NOEL_(ind) (= NOEL × BWI) was 76.8 mg a.i./ind. The number of days for a wild population to reach the NOEL in the laboratory population was calculated as NOEL_(ind)/DI (=55 d).

‡ For mouse acute oral toxicity, the most sensitive mice were females. Food consumption (FC) was 0.006 kg dw/ind/day, body weight per individual (BWI) was 0.023 kg bw/ind. Under Wild Mammal Scenario 2, the EEC was 54 mg a.i./kg dw. Therefore, the daily intake (DI = FC × EEC) was 0.32 mg a.i./ind/day. The NOEL_(ind) (= NOEL × BWI) was 11.5 mg a.i./ind. The number of days for a wild population to reach the NOEL in the laboratory population was calculated as NOEL_(ind)/DI (= 36 d).

Table 6 Risk to aquatic organisms

Organism	Exposure	Endpoint value	EEC	MOS	Risk
Freshwater species					
<i>Daphnia magna</i>	Acute	0.78 mg a.i./L	0.05 mg a.i./L	16	Negligible risk
	Chronic	0.078 mg a.i./L	—	N/A	No classification
Chironomid	Subchronic	0.74 mg a.i./L	Not determined	—	Not determined
Rainbow trout	Acute	0.7 mg a.i./L	0.05 mg a.i./L	14	Negligible risk
	Chronic	No data	N/A	N/A	No data
Bluegill sunfish	Acute	0.42 mg a.i./L	0.05 mg a.i./L	8.3	Low risk
	Chronic	No data	N/A	N/A	No data
Fathead minnow	Chronic (early-life stage toxicity)	0.082 mg a.i./L	0.05 mg a.i./L	1.6	Low risk

Organism	Exposure	Endpoint value	EEC	MOS	Risk
	Chronic (whole life-cycle toxicity)	0.027 mg a.i./L	N/A	N/A	Endpoints affected included time to first spawn, number of eggs produced, survival of parents and offspring, and number of eggs per spawn.
Freshwater algae	Acute	0.27 mg a.i./L	0.05 mg a.i./L	5.4	Low risk
Vascular plant	Dissolved	Incomplete data	N/A	N/A	Not determined
	Over-spray	Incomplete data	N/A	N/A	Not determined
Marine species					
Crustacean	Acute	0.16 mg a.i./L	0.05 mg a.i./L	3.2	Low risk (Result should be interpreted with caution for benthic species.)
	Chronic	No data	N/A	N/A	No data
Mollusk	Acute	0.53 mg a.i./L	0.05 mg a.i./L	11	Negligible risk (Result should be interpreted with caution for benthic species.)
	Chronic	No data	N/A	N/A	No data
Sheepshead minnow	Acute	0.89 mg a.i./L	0.05 mg a.i./L	19	Negligible risk
Salmonid	Acute	No data	N/A	N/A	No data
	Salinity challenge	No data	N/A	N/A	No data
Marine algae	Acute	No data	N/A	N/A	No data