

## **GEOPROCESS FILE SUMMARY REPORT**

### **WOLF LAKE MAP AREA N.T.S. 105B**

#### **INTRODUCTION**

The GEOPROCESS FILE is a compilation of information and knowledge on geological processes and terrain hazards, including mass movement processes, permafrost, flooding risks, faults, seismic activity and recent volcanism, etc. Please refer to the GEOPROCESS FILE Introduction and User's Guide for more in-depth information on how the maps were developed, which other GEOPROCESS FILE maps are available, how to utilize this inventory and how to interpret the legend. Special interest should be taken in the detailed description of the terrain hazard map units. Appendices in the User's Guide include summary papers on the geological framework, permafrost distribution, and Quaternary geology in Yukon and a list of comprehensive GEOPROCESS FILE references.

This report includes a brief discussion of the scope and limitations of the GEOPROCESS FILE compilation maps and summaries followed by summaries of the bedrock geology, surficial geology and terrain hazards for this N.T.S. map area, and a list of references.

#### ***Geological Processes and Terrain Hazard Compilation Maps***

The GEOPROCESS FILE map units were drafted on the 1:250,000 topographic base maps through interpretation from bedrock geology maps, surficial geology maps and in some cases terrain hazard maps at various scales. The compilation maps have a confidence level reflecting the original source material. All materials used to produce the maps are listed in the references attached to each map. A file containing the documentation used to construct these maps is available at the Indian and Northern Affairs library in Whitehorse, Yukon. Areas for which no surficial geology or terrain hazard information is published were left blank. Summary reports on surficial geology and terrain hazards for these map sheets were written by extrapolating the data from adjacent map sheets or smaller scale maps. Information from small scale (e.g. 1:1,000,000) maps was used for the summary reports, but not redrafted onto the 1:250,000 GEOPROCESS FILE maps.

The GEOPROCESS FILE compilation maps are intended as a first cut planning tool; the legend on the maps describes the general aspects of terrain hazards (also see below) and associated geological processes. **These maps should never replace individual site investigations for planning of site specific features, such as buildings, roads, pits, etc.**

#### ***Bedrock Geology Summaries***

Each 1:250,000 N.T.S. map area is described according to morphogeological belts and terranes defined by Gabrielse *et al.* (1991) and Wheeler *et al.* (1991). Bedrock geology (including structure) and mineral occurrences are briefly described and taken largely from the referenced, most recent 1:250,000 geological map with additional contributions from Wheeler and McFeely (1991), and Yukon MINFILE (1993). A summary paper ("A Geological Framework for Yukon") in Appendix A of the Introduction and User's Guide provides a framework and context for each of the bedrock summaries.

The level of knowledge and understanding of Yukon geology is constantly evolving with more detailed mapping and development of geological models. Names, ages and terrane affinities of rock units on the most recent 1:250,000 geological maps may, in some cases, now be considered incorrect. Thus information contained within some of the bedrock geology summaries may be out of date. Although much of the information reflects the knowledge at the time that the source map was published, additional information has been inserted whenever possible to assist the user in merging the information with current geological maps, concepts and understanding. The age ranges for similar packages of rocks may also vary between map areas since the actual rocks, or at least the constraints on their age, may vary

between map areas.

## **BEDROCK GEOLOGY**

The Wolf Lake map area is entirely within the Omineca Belt. The mountainous terrain of the Cassiar Mountains occupies most of the southern part of the map area whereas much of the northern half is underlain by low-lying regions of the Wolf Lake, Nisutlin Plateau and Liard Plain. The divide between the Yukon and Mackenzie River drainages follows an irregular line through the middle of the map area.

There are four main stratigraphic/tectonic packages of rocks in the Wolf Lake map area -- Slide Mountain, Dorsey, Cassiar and Yukon-Tanana Terranes. Rocks in the southwestern half of the map area include two northwest-trending belts of Slide Mountain Terrane volcanic rocks that are separated by the sedimentary rocks of Dorsey Terrane. Slide Mountain Terrane rocks include 400-340 million year old greenstone, chlorite schist, quartzite, phyllite, slate, argillite, chert, quartz-albite-mica gneiss, actinolite schist, limestone and dolomite. Dorsey Terrane rocks include 400-320 million year old chert, hornfels, argillite, slate, phyllite, quartzite, limestone, skarn, dolomite and conglomerate.

The central block, between Wolf Lake and the Liard River valley is dominated by Cassiar Terrane pre-570 million year old biotite schist, quartzite, marble, gneiss, slate, phyllite, quartz grit, conglomerate, hornfels, limestone and dolomite that are separated from younger 540-420 million year old Road River Group (also Cassiar Terrane) phyllite, limestone, hornfels and skarn and minor 390-325 million year old Earn Group. Northeast of the Tintina Fault, Yukon-Tanana Terrane rocks are represented by biotite schist and gneiss.

Several, relatively small intrusions of 180 million year old diorite, granodiorite and quartz diorite occur in the southwestern part of the map area. Numerous very large, 100 million year old batholiths, and smaller plutons of biotite-quartz monzonite, granodiorite, and alaskite occur throughout the map area and are responsible for localized hornfels and doming of the metasedimentary rocks in the central Cassiar Terrane. There are also some small 50 million year old felsic intrusions in the southern part of the map area.

Several outcrops of recent columnar jointed, olivine basalt belong to the Tuya lavas and are 200,000-800,000 years old.

### ***Mineral Deposits and Occurrences***

The Wolf Lake map area contains 141 mineral occurrences of which 103 host known mineralization. The area is known for its numerous occurrences (33) of silver-rich lead veins, skarns and replacement deposits which are concentrated near Rancheria in the southern part of the map area. The Logan deposit hosts 12.3 million tonnes of 6.17% zinc and 26 grams per tonne silver, and the Hart property contains 97,000 tonnes of 1025 grams per tonne silver. The southwestern portion of the map area hosts numerous tin-tungsten-molybdenum skarns, veins and porphyry deposits associated with granitic rocks. The largest of these deposits is the Logtung porphyry which hosts 160 million tonnes of 0.12% tungsten oxide and 0.052% molybdenum oxide. Also of note are asbestos and topaz occurrences.

## **SURFICIAL GEOLOGY**

The main sources of information for the Wolf Lake area are a surficial geology map by Klassen (1982), and a report by Jackson (1994) which discusses the terrain hazards and surficial geology of adjoining map sheets.

The surface deposits of the Wolf Lake map sheet are associated with the most recent Cordilleran ice sheet (McConnell) which is believed to have covered south and central Yukon between 26,500 and 10,000 years ago. The western half of the map sheet was covered by the Cassiar lobe which

flowed west and northwest of the Cassiar Mountains. The Liard lobe which flowed eastward has left streamlined morainal deposits (drumlins) in the Twin Lake valley. The Cassiar Mountains occupy the central portion of the map and were covered by ice caps and cirque glaciers. The surface of these mountains now consists mainly of exposed bedrock, locally covered by thin colluvial or morainal deposits.

The northwest corner of the map area is dominantly covered by morainal deposits. Till, or more correctly diamicton of glacial origin is an unsorted mixture of coarse material ranging in size from pebble to boulder, with a matrix of clay, silt and sand. The general composition of the till matrix in adjoining map areas (Jackson, 1994) indicates a wide range of content of sand (20 to 70%), silt (20 to 80%) and commonly lower clay (5 to 30%). The low clay content is reflected by the low plasticity of the matrix. Morainal deposits can provide a stable base, if there is no permafrost present. Lenses of permafrost may occur locally on north facing slopes and at high elevations, where thick organic deposits overlay the Quaternary sediments.

The Liard River and Rancheria River valleys, as well as the major creeks such as Gravel, Ram, Cabin and Irving Creeks, are commonly benched and covered by glaciofluvial sand and gravel. These sediments of variable thickness and composition, usually provide stable surfaces. However, they locally contain undesirable lithologies (weak) for their potential use as aggregate. These well drained and coarse deposits are commonly capped by silt and fine sand blankets (0.3 to 1.5 m thick).

Glaciolacustrine sediments occur around Wolf Lake. They commonly contain massive ice bodies and are susceptible to retrogressive thaw slide and thermokarst degradation when disturbed either by river erosion, forest fires, or other changes in surface conditions.

The White River Ash (1,200 years B.P.) is found at the surface of most landforms, except on actively colluviated slopes, landslides, and very recent alluvial landforms. This tephra can be used in some cases as an indication of active slope wash, creep or aeolian activity.

## **TERRAIN HAZARDS**

Slope failures in steep bedrock represent the highest risk hazard in the area. Although not documented in any of the publications available and not present on the terrain hazard map, the potential for rock slide and avalanches should be kept in mind when an area is investigated.

### ***Seismicity***

There are seven recorded seismic events within the northeast part of the map area. All of the recorded events are 4.0 to 4.999 or less in magnitude.

### ***Mass Movement Processes***

A few small earth flows or landslides were mapped in the Cassiar Mountains, but no large slumps or slides were identified. Snow avalanches could occur on steep bedrock slopes and can entrain a large volume of boulders and debris. Development of any kind in close proximity to steep escarpments, ridges,

cirques and arrêtes prone to these hazards should be discouraged. Map units labelled cs are considered unstable.

### ***Permafrost***

This area lies within the discontinuous permafrost zone (Brown, 1967). Distribution of permafrost is expected to be sporadic (Heginbottom and Radburn, 1992). Permafrost is commonly present at high elevations in colluvial and morainal blankets covered by thick organic mats. Its presence is often indicated by solifluction lobes, stripes and sorted stone polygons. Ice content in morainal and colluvial deposits is estimated to be low to moderate, in the form of small lenses, veins and crystals. Ice content is probably absent from coarser, well-drained deposits such as glaciofluvial sand and gravels (eskers, kames, terraces), and gravelly to sandy fluvial deposits. Ice content should be highest in fine-grained, less permeable sediments such as silty fluvial terrace deposits, the lowermost part of colluvial and alluvial fans, and glaciolacustrine deposits. These landforms are commonly covered with thick moss and organic soils which increase the possibility of permafrost. Thermokarst collapse and thaw slides are possible hazards in fine-grained glaciolacustrine and fluvial sediments around Wolf Lake.

### ***Flooding and Other Risks***

Floods related to ice-jams, snow melt and summer rainstorms are possible hazards in lower reaches of most streams in the area. The steep portions of alluvial fans, in addition to the flooding risk, are also exposed to the additional possibility of mud flows and debris flows associated with rapid discharge increase.

## References

### Wolf Lake Map Area N.T.S. 105 B

**To be thorough, check the references for adjacent N.T.S. map sheets and the General Reference List (See Introduction and User's Guide).**

**Most of the following references should be available for viewing in the DIAND library on the third floor of the Elijah Smith building in Whitehorse. The library and call number of some internal government reports are listed.**

- Abbott, J.G., 1986, Epigenetic mineral deposits of the Ketzia-Seagull district, Yukon. *In:* Yukon Geology, Vol. 1, Exploration and Geological Services Division, Yukon, Indian and Northern Affairs Canada, p. 56-62.
- Abbott, J.G., 1985, Silver-bearing veins and replacement deposits of the Rancheria District. *In:* Yukon Exploration and Geology 1983, Exploration and Geological Services Division, Yukon, Indian and Northern Affairs Canada, p. 34-41.
- Amukun, S.E. and Lowey, G.W., 1987, Geology of the Sab Lake (105B/7) and Meister Lake (105B/8) map-areas, Rancheria district, southeast Yukon, Canada. Exploration and Geological Services Division, Yukon, Indian and Northern Affairs Canada, Open-File Report 1987-1, (two 1:50,000 scale maps with marginal notes).
- Bremner, T. and Liverton, T., 1990, DAN. *In:* Yukon Exploration 1990; Exploration and Geological Services Division, Yukon, Indian and Northern Affairs Canada, p. 27-30.
- Bremner, T., 1990, MUNSON. *In:* Yukon Exploration 1990; Exploration and Geological Services Division, Yukon, Indian and Northern Affairs Canada, p. 59.
- Bremner, T., 1990, LOGAN. *In:* Yukon Exploration 1990; Exploration and Geological Services division, Yukon, Indian and Northern Affairs Canada, p. 40-41.
- Brown, R.J.E., 1967, Permafrost in Canada. Geological Survey of Canada, Map 1246A, (scale 1:7,603,200).
- Carlyle, L., 1995, Placer mining and exploration compilation (NTS 105A/B/C/D). Exploration and Geological Services Division, Yukon, Indian and Northern Affairs Canada, Canada/Yukon Economic Development Agreement, Open File 1995-10(G).
- Cathro, M.S., 1988, Gold and silver, lead deposits of the Ketzia River district, Yukon: Preliminary results of Field Work. *In:* Yukon Geology, Vol. 2, Exploration and Geological Services Division, Yukon, Indian and Northern Affairs Canada, p. 8-25.
- Fortescue, J.A.C., Gleeson, C.F., Kuehnbaum, R.M. and Martin, L., 1982. A practical approach to the interpretation of regional geochemical maps; an example using the Wolf Lake sheet, Yukon. Western Miner, Vol. 55, No. 2, p. 35-36.
- DIAND library**
- Gabrielse, H., Tempelman-Kluit, D.J., Blusson, S.L. and Campbell, R.B. (comp.), 1980, MacMillan River, Yukon - District of MacKenzie-Alaska (Sheet 105, 115). Geological Survey of Canada, Map 1398A (one 1:1,000,000)
- NTS 105, 115**

Gabrielse, H. and Yorath, C.J. (eds), 1991, Geology of the Cordilleran Orogen in Canada. Geological Survey of Canada, No. 4, 844 p.

**Contains summary of Yukon geology**

Geological Survey of Canada, 1986, Regional stream and water geochemical reconnaissance data, Yukon Territory. Geological Survey of Canada, Open File 563.

Germann, A., Friedrich, G., and Schattner, R., 1992, Ore mineralogy and formation conditions of vein and replacement-type Pb-Zn-Ag occurrences, Logan and YP properties, Rancheria District, Yukon. **In:** Yukon Geology, Vol. 3, Exploration and Geological Services Division, Yukon, Indian and Northern Affairs Canada, p. 37-44.

Hall, A. and Liverton, T., 1992, Trace ammonium in granites of the southern Yukon and its petrogenetic significance. **In:** Yukon Geology, Vol. 3, Exploration and Geological Services Division, Indian and Northern Affairs Canada, p. 45-51.

Heginbottom, J.A. and Radburn, L.K. (comp.), 1992, Permafrost and ground ice conditions of northwestern Canada. Geological Survey of Canada, Map 1691A, scale 1:1,000,000.

Indian and Northern Affairs, 1993, Abandoned Mines Assessment, 105B-01-1, Dale. DIAND Technical Services, Yukon, Indian and Northern Affairs Canada.

Indian and Northern Affairs, 1993, Abandoned Mines Assessment, 105B-01-2, Silver Seven. DIAND Technical Services, Yukon, Indian and Northern Affairs Canada.

Indian and Northern Affairs, 1993, Abandoned Mines Assessment, 105B-04-1, Logtung. DIAND Technical Services, Yukon, Indian and Northern Affairs Canada.

Indian and Northern Affairs, 1993, Abandoned Mines Assessment, 105B-04-2, Logjam. DIAND Technical Services, Yukon, Indian and Northern Affairs Canada.

Indian and Northern Affairs, 1993, Abandoned Mines Assessment, 105B-07-1, Hart. DIAND Technical Services, Yukon, Indian and Northern Affairs Canada.

Indian and Northern Affairs, 1993, Abandoned Mines Assessment, 105B-07-2, Silverhart. DIAND Technical Services, Yukon, Indian and Northern Affairs Canada.

Indian and Northern Affairs, 1993, Yukon MinFile 105B - Wolf Lake. Exploration and Geological Services Division, Yukon, Indian and Northern Affairs Canada.

Jackson, L.E., Jr. and MacKay, T.D., 1990, Glacial limits and ice-flow directions of the last Cordilleran ice sheet in Yukon Territory between 60 and 63 degrees north. Geological Survey of Canada, Open File 2329.

**NTS 95D, 105A, 105B, 105C, 105D, 115A, 115E, 115F, 115G, 115H, 115I, 115J, 106K, 106L, 115G, 115H**

Jackson, L.E., 1994, Terrain inventory and Quaternary history of the Pelly River area, Yukon Territory. Geological Survey of Canada, Memoir 437, 41 p.

**NTS 105J, 105K, 105F, 105G**

Klassen, R.W., 1978, A unique stratigraphic record of late Tertiary - Quaternary events in southeastern Yukon, Canadian Journal of Earth Sciences, Vol. 15, No. 11, p. 1884-1885.

Klassen, R.W., 1978, Surficial geology of Rancheria River, Meister River, Takhini River, Swift River and

Tagish, southern Yukon. Geological Survey of Canada, Open File 539 (1:100,000 scale maps).  
**NTS 104N/16, 104O/13,14 105A SW 105B/1,2,3 105C/1,2,3,4,5,6,7 105D/7,8,9,10,NW**

Klassen, R.W., 1982, Surficial geology; Wolf Lake, Yukon Territory. Geological Survey of Canada, Vol. 14-1982, 1 sheet, Surficial geology map, scale 1:250,000.

Klassen, R.W., 1987, The Tertiary-Pleistocene stratigraphy of the Liard plain, southeastern Yukon Territory. Geological Survey of Canada, Paper 86-17, 16 p.  
**NTS 105A, 105B**

Klassen, R.W., Thorsteinsson, E. and Hughes, O.L., 1978, Surficial geology and terrain evaluation, southern Yukon. *In*: Current Research, Part A, Geological Survey of Canada, No. 78-1A, p. 465.

Lowey, G.W. and Lowey, J.F., 1986, Geology of Spencer Creek (105B/1) and Daughney Lake (105B/2) map areas, Rancheria district, southeast Yukon, Canada. Exploration and Geological Services Division, Yukon, Indian and Northern Affairs Canada, Open File Report 1986-1 (text with two 1:50,000 scale maps).

Mulligan, R., 1969, Metallogeny of the region adjacent to the northern part of the Cassiar Batholith, Yukon Territory and British Columbia (parts of 104O, P and 105B). Geological Survey of Canada, Paper 68-70, 13 p.

Murphy, D.C., 1988, Geology of Gravel Creek (105B/10) and Irvine Lake (105B/11) map-areas, southeastern Yukon. Exploration and Geological Services Division, Yukon, Indian and Northern Affairs Canada, Open-File 1988-1 (two 1:50,000 maps with report), 61 p.

Poole, W.H., 1956, Geology of the Cassiar Mountains in the vicinity of the Yukon-B.C. boundary. Ph.D. thesis, Princeton University.

Poole, W.H., Roddick, J.A. and Green, L.H., 1960, Geology, Wolf Lake, Yukon Territory. Geological Survey of Canada, Preliminary Series Map 10-1960, scale 1:253,440.

Soroka, I. and Jack, M.E., 1982, Baseline study of the watershed near proposed Marbaco Mine, B.C. and Marbaco Mill, Yukon Territory. Department of Environment, Environmental Protection Service, Pacific Region, Yukon Branch, Regional Program Report 82-06, 63 p.  
**NTS 105B**  
**(Economic Development library)**  
**Call Number: TD 227 Y8 S67**

Wheeler, J.O., Brookfield, A.J., Gabrielse, H., Monger, J.W.H., Tipper, H.W. and Woodsworth, G.J., 1991, Terrane map of the Canadian Cordillera. Geological Survey of Canada, Map 1713.

Wheeler, J.O. and McFeely, P., 1991, Tectonic Assemblage map of the Canadian Cordillera and adjacent parts of the United States of America. Geological Survey of Canada. Map 1712A.