

GEOPROCESS FILE SUMMARY REPORT

WHITEHORSE MAP AREA N.T.S. 105D

INTRODUCTION

The GEOPROCESS FILE is a compilation of information and knowledge on geological processes and terrain hazards, including mass movement processes, permafrost, flooding risks, faults, seismic activity and recent volcanism, etc. Please refer to the GEOPROCESS FILE Introduction and User's Guide for more in-depth information on how the maps were developed, which other GEOPROCESS FILE maps are available, how to utilize this inventory and how to interpret the legend. Special interest should be taken in the detailed description of the terrain hazard map units. Appendices in the User's guide include summary papers on the geological framework, permafrost distribution, and Quaternary geology in Yukon and a list of comprehensive GEOPROCESS FILE references.

This report includes a brief discussion of the scope and limitations of the GEOPROCESS FILE compilation maps and summaries followed by summaries of the bedrock geology, surficial geology and terrain hazards for this N.T.S. map area, and a list of references.

Geological Processes and Terrain Hazard Compilation Maps

The GEOPROCESS FILE map units were drafted on the 1:250,000 topographic base maps through interpretation from bedrock geology maps, surficial geology maps and in some cases terrain hazard maps at various scales. The compilation maps have a confidence level reflecting the original source material. All materials used to produce the maps are listed in the references attached to each map. A file containing the documentation used to construct these maps is available at Exploration and Geological Services Division, Indian and Northern Affairs Canada in Whitehorse, Yukon. Areas for which no surficial geology or terrain hazard information is published were left blank. Summary reports on surficial geology and terrain hazards for these map sheets were written by extrapolating the data from adjacent map sheets or smaller scale maps. Information from small scale (e.g. 1:1,000,000) maps was used for the summary reports, but not redrafted onto the 1:250,000 GEOPROCESS FILE maps.

The GEOPROCESS FILE compilation maps are intended as a first cut planning tool; the legend on the maps describes the general aspects of terrain hazards (also see below) and associated geological processes. **These maps should never replace individual site investigations for planning of site specific features, such as buildings, roads, pits, etc.**

Bedrock Geology Summaries

Each 1:250,000 N.T.S. map area is described according to morphogeological belts and terranes defined by Gabrielse *et al.* (1991) and Wheeler *et al.* (1991). Bedrock geology, geological structures and mineral occurrences are briefly described and taken largely from the referenced, most recent 1:250,000 geological map with additional contributions from Wheeler and McFeely (1991), and Yukon MINFILE (1993). A summary paper ("A Geological Framework for Yukon") in Appendix A provides a framework and context for each of the bedrock summaries.

The level of knowledge and understanding of Yukon geology is constantly evolving with more detailed mapping and development of geological models. Names, ages and terrane affinities of rock units on the most recent 1:250,000 geological maps may, in some cases, now be considered incorrect. Thus information contained within some of the bedrock geology summaries may be out of date. Although much of the information reflects the knowledge at the time that the source map was published, additional

information has been inserted whenever possible to assist the user in merging the information with current geological maps, concepts and understanding. The age ranges for similar packages of rocks may also vary between map areas since the actual rocks, or at least the constraints on their age, may vary between map areas.

BEDROCK GEOLOGY (Wheeler 1961)

Most of the Whitehorse map area is in the Intermontane Belt except the southwestern portion which is in the Coast Belt. The Intermontane portion of the map area is characterized by rounded mountainous topography of the Teslin Plateau while the Coast Belt regions are rugged with much greater relief and outcrop exposure. The Yukon River valley and headwater lakes are the most notable physiographic features.

The Intermontane portion of the map area is underlain by the volcanic and sedimentary rocks of Stikinia and Cache Creek Terrane. Stikinia forms a northwest-trending belt that comprises the bulk of the map area and is composed of 230-200 million year old Lewes River Group basalt flows, andesite and associated pyroclastic rocks. These rocks are overlain by 200-160 million year old Whitehorse Trough Lewes River Group greywacke, siltstone, argillite, conglomerate, tuff and limestone, and Laberge Group greywacke, arkose, quartzite, conglomerate, siltstone, argillite and hornfels. Cache Creek Terrane rocks occur in the southeastern part of the map area and are faulted against rocks of Stikinia. Cache Creek rocks include 360-180 million year old greenstone flows, pyroclastic rocks, limestone, limestone breccia, chert and metamorphosed volcanic rocks containing numerous variably serpentized bodies of dunite, peridotite and pyroxenite. Tantalus Formation (100-60 million years old) arkose, siltstone, chert-pebble conglomerate argillite and coal occur as numerous small exposures along the western margin of Stikinia.

The Coast Belt portion of the map area is composed of ancient, metasedimentary rocks and more recent felsic plutonic rocks. The pre-550-320 million year old metasedimentary quartz-mica, quartz-chlorite, and mica schists, quartzite, micaceous quartzite, gneiss, marble and amphibolite used to be called the Yukon Group, but now are grouped as rocks of the Nisling and Nasina Assemblages of Yukon Tanana Terrane. The plutonic rocks consist of 180-50 million year old granite, granodiorite, quartz monzonite, quartz diorite, diorite and syenite of the Coast Plutonic Complex. Similar plutons also intrude Stikinia and Cache Creek Terrane rocks throughout the map area. Coarse-grained 55 million year old alaskite in the western part of the map area is crumbly weathering and shows well-defined joint surfaces. Volcanic rocks of the 95-70 million year old Hutshi Group (Mount Nansen and Carmacks groups) and 55 million year old Skukum Group include andesite, basalt, rhyolite, breccia and tuffs that are scattered through the map area. The youngest rocks in the map area are 2-9 million year old Miles Canyon columnar jointed olivine basalt and minor pyroclastic rocks which form numerous exposures, particularly near Whitehorse where they are known to overlie unconsolidated gravels.

Mineral Deposits and Occurrences

There are 196 mineral prospects listed in Yukon MINFILE for the Whitehorse map area. Most of these are fairly small, gold-silver veins of the Wheaton River area or copper-molybdenum skarns in Lewes River Group limestone adjacent to 115 million year old granodiorite plutons of the Whitehorse Copper Belt. Greater than 123,000 tonnes of copper, 90 tonnes of silver and 7 tonnes of gold were extracted from over 10 million tonnes of ore from seven deposits within the copper belt. Gold-silver mines included the Mount Skukum mine which produced 2,622 kilograms of gold from 201,461 tonnes; the Venus mine on Montana Mountain which produced 383 kilograms of gold, 12 tonnes of silver, 2,440 tonnes of lead and 1,650 tonnes of zinc from 66,000 tonnes of ore; and the Arctic Caribou deposit, also on Montana Mountain, which produced 543 kilograms of gold and 18 tonnes of silver from 53,500 tonnes of ore.

Reserves are reported for the following mineral deposits:

| Deposit | Tonnes | Commodity |
|----------------|---------------|------------------|
| Becker-Cochran | 127,273 | 4% antimony |

| | | |
|--|-----------|---|
| Arctic Caribou | 74,575 | 13.4g/t Au, 350 g/t Ag |
| Mount Skukum | 100,000 | 14.7 g/t Au |
| Skukum Creek | 800,150 | 7.6 g/t Au, 275 g/t Ag |
| Venus | 100,000 | 8.37 g/t Au, 247 g/t Ag 2.1% lead, 1.4% zinc |
| Whitehorse Copper (in several deposits) | 3,000,000 | 1% Cu, 0.06% molybdenum 0.1 g/t Au, 3.7 g/t Ag |

The area also hosts copper-gold, antimony and lead-silver veins as well as uranium and asbestos occurrences. Numerous coal occurrences are associated with Tantalus Formation strata southwest of Whitehorse including reserves of 500,000 tonnes at the Whitehorse Coal deposit.

Limited placer mining has been carried out on Sheldon Creek which drains into the Teslin River. A few attempts at placer mining have taken place on the Wheaton River.

SURFICIAL GEOLOGY

The Whitehorse map area lies within the limits of the McConnell Glaciation (Bostock, 1966). Subsequent to the maximum extent of McConnell ice, deglaciation produced disrupted drainage systems and large glacial lakes as a result of a complex assemblage of ice lobes restricted to valley bottoms which were controlled by the local topography. In the Whitehorse map area, representative sequences of Quaternary deposits are found in many major valleys such as the Yukon River valley, the Takhini River valley, and the M'Clintock River valley. Typically, morainic and colluvial blankets overlying the bedrock are found at high elevations above valley bottoms. Glaciofluvial sand and gravel terraces flank the valley sides and pitted or hummocky deposits of sand and gravel deposits line the bottom of some valleys. In the Takhini River and the Tagish River valleys, glacial lake Champagne deposited up to 75 metres of silt and clay. These fine grained, soft sediments are easily eroded and are susceptible to slumping, thermokarsting and frost heaving. Strandlines (beach deposits) can also be seen on valley sides at average elevations of 2500 ft above sea level.

The major sources of information for the compilation of terrain hazards and processes map include: 1. the Southern Lakes ecological land classification mapping (Morison and McKenna, 1981), and 2. the surficial geology maps by Morison and Klassen (1991) and various soil surveys by Mougeot and Smith (1992) and Rostad, *et al.* (1977).

TERRAIN HAZARDS

Geological processes within the Whitehorse map area which present the most immediate hazard to human activity are related to extreme topography, primarily in the Coast Mountains in the southwestern portion of the map area. Steep ridges, debris filled cirques, remnants of alpine glaciers, and steep precipitous walls are common.

Seismicity

One magnitude 4-5 earthquake and three magnitude 2-3 earthquake epicentres have been recorded in the southern portion of the map area. No evidence of landslide activity or fault movement related to these events has been noted. There is, however, concern over the presence of clay and silt substratum under the City of Whitehorse, since they are a good transmitter of S-waves. The effects of large magnitude earthquakes in the Gulf of Alaska are, as a result, commonly felt in the City of Whitehorse.

Mass Movement Processes

Rapid mass movement hazards include slope failures, avalanches and rockfalls. These active processes are considered as severe hazards and should thus be considered in development activities in the Whitehorse map area. Periglacial processes present, such as solifluction, nivation and thermokarst, may impact the stability of slopes covered by colluvial and morainal deposits, as well as the stability of river banks cut into silty sediments.

Snow slides affect the south Klondike Highway just north of the British Columbia-Yukon border; these are monitored on a regular basis and the highway is closed when hazards reach unacceptable levels.

Permafrost

The Whitehorse map sheet is part of the scattered permafrost zone (Brown, 1978). Generalized comments about permafrost distribution can be inferred from information available in adjoining map sheets (Heginbottom and Radburn, 1992). Permafrost is probably more common at high elevations, in morainal and colluvial deposits with visible ice, and nivation and cryoplanation features. At lower elevations, alluvial and glaciolacustrine sediments may have less extensive permafrost but higher ice content. Thermokarsting can develop in fine-grained glaciolacustrine sediments and fine-grained alluvial sediments in areas such as the Takhini valley (Rampton, 1972; Klassen, 1979).

Flooding Risks

Flooding in the Whitehorse map area is caused by unusually high precipitation, snowmelt runoff or ice jams during break up time. Ice jams commonly occur along the Yukon River, below Lake Laberge (Underwood McLelland, 1983). The Marwell area, located at the north end of the City of Whitehorse, is often flooded during late fall and early spring because of ice dams on the Yukon River. Minor flooding may occur at the mouth of the Takhini River into the Yukon River.

Other

Areas underlain by loosely consolidated boulder tills containing large granitic boulders can produce significant levels of radon gas in houses. This problem is easily overcome by using proper construction and ventilation techniques.

References

Whitehorse Map Area N.T.S. 105D

Note: To be thorough, check the references for adjacent N.T.S. map sheets and the General Reference List.

Most of the following references should be available for viewing in the DIAND library on the third floor of the Elijah Smith building in Whitehorse. The library and call number of some internal government reports are listed.

Bostock, H.S., 1966, Notes on glaciation in central Yukon Territory. Geological Survey of Canada, Paper 65-36, 18 p.

NTS 105N/E, 115N/W

Bremner, T., 1988, Geology of the Whitehorse coal deposit. *In:* Yukon Geology, Vol. 2, Exploration and Geological Services Division, Yukon, Indian and Northern Affairs Canada, p. 1-7.

Brooker, E.W. and Associates, 1971, Soil and Foundation Investigation of Whitehorse Escarpment. Department of Transportation, City of Whitehorse.

***Whitehorse Public Library
Call No. 551.302 Elm YA***

Brown, R.J.E., 1978, Permafrost: Plate 32, Hydrological Atlas of Canada, Fisheries and Environment Canada, Ottawa. 34 plates.

Bultman, T.R., 1979, Geology and tectonic history of the Whitehorse Trough west of Atlin, British Columbia. Unpublished Ph.D thesis, Yale University.

Burn, C.R., 1993, Permafrost degradation following forest fire, Takhini Valley, southern Yukon Territory, Ford, D., McCann, B. and Vajoczki, S. (compilers), *In:* Program with abstracts, 3rd International Geomorphology Conference, p. 116.

Dobrowolsky, H. and Ingram, R., 1993, A History of the Whitehorse Copper Belt. Exploration and Geological Services Division, Yukon, Indian and Northern Affairs Canada, Canada/Yukon Economic Development Agreement, Information Open File 1993-1(I).

Doherty, R.A. and Hart, C.J.R., 1988, Preliminary geology of Fenwick Creek (105D/3) and Alligator Lake (105D/6) map-areas. Exploration and Geological Services Division, Yukon, Indian and Northern Affairs Canada, Open File 1988-2.

EBA Engineering Consultants Ltd., 1988, Geotechnical evaluation, Watson River homestead subdivision, Carcross, Yukon. Contract report to Community and Transportation Services, Yukon Government. EBA Engineering Consultants, Ltd., Whitehorse, Yukon, 21 p.

Eiche, G.E., 1985, Petrology of Quaternary Alkaline Lavas from the Alligator Lake Volcanic Complex, Yukon Territory. Unpublished M.Sc. thesis, McGill University.

Gabrielse, H., Tempelman-Kluit, D.J., Blusson, S.L. and Campbell, R.B. (comp.), 1980, MacMillan River, Yukon - District of MacKenzie-Alaska (Sheet 105, 115). Geological Survey of Canada, Map 1398A (one 1:1,000,000 map).

NTS 105, 115

Gabrielse, H. and Yorath, C.J. (eds), 1991, Geology of the Cordilleran Orogen in Canada. Geological Survey of Canada, No. 4, 844 p.

Contains summary of Yukon geology

Geological Survey of Canada, 1985, Regional Stream Sediment and Water Geochemical Reconnaissance Data, NTS 105D, Open File 1218.

Hart, C.J.R., 1993, Geologic map of Thirty-seven Mile Creek map area, southern Yukon Territory. Exploration and Geological Services Division, Yukon, Indian and Northern Affairs Canada, Canada/Yukon Economic Development Agreement, Geoscience Open File 1993-4(G), (1:50,000 scale map).

Hart, C.J.R. and Brent, D., 1993, Preliminary geology of the Thirty-seven Mile Creek map sheet (105D/13). *In:* Yukon Exploration and Geology - 1992, Exploration and Geological Services Division, Yukon, Indian and Northern Affairs Canada, p. 39-48.

Hart, C.J.R. and Hunt J.A., 1994a, Geology of the Joe Mountain map area (105D/15), southern Yukon Territory. *In:* Yukon Exploration and Geology - 1993, Exploration and Geological Services Division, Yukon, Indian and Northern Affairs Canada, p. 47-66.

Hart, C.J.R. and Hunt, J.A., 1994b, Geological map of Joe Mountain map area, Yukon (105D/15). Exploration and Geological Services Division, Yukon, Indian and Northern Affairs, Canada, Canada/Yukon Economic Development Agreement, Geoscience Open-File 1994-4(G), (1:50,000 scale map).

Hart, C.J.R. and Hunt, J.A., 1995a, Geology of the Mount M'Clintock map area (105D/16), southern Yukon Territory. *In:* Yukon Exploration and Geology, 1994, Exploration and Geological Services Division, Yukon, Indian and Northern Affairs Canada, p. 87-104.

Hart, C.J.R. and Hunt, J.A., 1995b, Geology of the Mount M'Clintock map area (105D/16), southern Yukon Territory. Exploration and Geological Services Division, Yukon, Indian and Northern Affairs, Canada, Canada/Yukon Economic Development Agreement, Geoscience Open -File 1995-4(G), (scale 1:50,000).

Hart, C.J.R. and Pelletier, K.S., 1989a, Geology of Carcross (105D/2) and part of Robinson (105D/7) map areas. Exploration and Geological Services Division, Yukon, Indian and Northern Affairs Canada, Open File 1989-1, 84 p.

Hart C.J.R. and Pelletier, K.S., 1989b, Geology of Whitehorse (105D/11) map area. Exploration and Geological Services Division, Yukon, Indian and Northern Affairs Canada, Open File 1989-2, 51 p.

Hart, C.J.R. and Radloff, J.K., 1990, Geology of Whitehorse, Alligator Lake, Fenwick Creek, Carcross and part of Robinson map areas (105D/11, 6, 3, 2 and 7). Exploration and Geological Services Division, Yukon, Indian and Northern Affairs Canada, Canada/Yukon Economic Development Agreement, Geoscience Open File 1990-4, 113 p.

Heginbottom, J.A. and Radburn, L.K. (comp.), 1992, Permafrost and ground ice conditions of northwestern Canada. Geological Survey of Canada, Map 1691A, scale 1:1,000,000.

Hunt, J.A., 1989, Thermal maturation and source rock potential of the Tantalus Formation, Whitehorse area, southern Yukon Territory. Unpublished B.Sc. thesis, University of British Columbia.

Hunt, J.A. and Hart, C.J.R., 1994, Thermal maturation and hydrocarbon source rock potential of the Tantalus Formation coals in the Whitehorse area, Yukon Territory. *In:* Yukon Exploration and

Geology - 1993, Exploration and Geological Services Division, Yukon, Indian and Northern Affairs Canada, p. 67-77.

Indian and Northern Affairs, 1993, Abandoned Mines Assessment, 105D-02-1, Venus. DIAND Technical Services, Yukon, Indian and Northern Affairs Canada.

Indian and Northern Affairs, 1993, Abandoned Mines Assessment, 105D-02-2, Big Thing. DIAND Technical Services, Yukon, Indian and Northern Affairs Canada.

Indian and Northern Affairs, 1993, Abandoned Mines Assessment, 105D-02-3, Arctic Caribou. DIAND Technical Services, Yukon, Indian and Northern Affairs Canada.

Indian and Northern Affairs, 1993, Abandoned Mines Assessment, 105D-03-4, Tally-Ho. DIAND Technical Services, Yukon, Indian and Northern Affairs Canada.

Indian and Northern Affairs, 1993, Abandoned Mines Assessment, 105D-06-1, Wheaton Mountain. DIAND Technical Services, Yukon, Indian and Northern Affairs Canada.

Indian and Northern Affairs, 1993, Abandoned Mines Assessment, 105D-06-2, Gold Hill. DIAND Technical Services, Yukon, Indian and Northern Affairs Canada.

Indian and Northern Affairs, 1993, Abandoned Mines Assessment, 105D-06-3, Union. DIAND Technical Services, Indian and Northern Affairs Canada.

Indian and Northern Affairs, 1993, Abandoned Mines Assessment, 105D-06-4, Legal Tender. DIAND Technical Services, Indian and Northern Affairs Canada.

Indian and Northern Affairs, 1994, Yukon MINFILE 105D - Whitehorse. Exploration and Geological Services Division, Yukon, Indian and Northern Affairs Canada.

Jackson, L.E., Jr. and MacKay, T.D., 1990, Glacial limits and ice-flow directions of the last Cordilleran ice sheet in Yukon Territory between 60 and 63 degrees north. Geological Survey of Canada, Open File 2329.

NTS 95D, 105A, 105B, 105C, 105D, 115A, 115E, 115F, 115G, 115H, 115I, 115J, 106K, 106L, 115G, 115H

Klassen, R.W., 1978, Surficial geology of Rancheria River, Meister River, Takhini River, Swift River and Tagish, southern Yukon. Geological Survey of Canada, Open File 539, (1:100,000 scale maps).

NTS 104N/16 104O/13, 14 105A SW* 105B/1,2,3 105C/1,2,3,4,5,6,7 105D/7,8,9,10, NW

- Klassen, R.W., 1979, Thermokarst terrain near Whitehorse, Yukon Territory. Geological Survey of Canada, Paper 79-1A.
- Lambert, M.B., 1968. Wheaton River and Lindeman Lake, Yukon and British Columbia. Geological Survey of Canada, Paper 68-1.
- Lambert, M.B., 1974, The Bennett Lake cauldron subsidence complex, British Columbia and Yukon Territory. Geological Survey of Canada, Bulletin 227.
- Love, D.A., 1989, Geology of the epithermal Mount Skukum gold deposit, Yukon Territory. Geological Survey of Canada, Open File 2123.
- MacKay, G, Diment, R. and Falkiner, J., 1993, Whitehorse Copper Belt: A simplified technical history. Exploration and Geological Services Division, Yukon, Indian and Northern Affairs, Canada, Canada/Yukon Economic Development Agreement, Information Open File 1993-2(I).
- McDonald, B.W.R., 1986, Geology and the genesis of the Mount Skukum Tertiary epithermal gold-silver vein deposit, Southwestern Yukon. Unpublished M.Sc. thesis, University of British Columbia.
- McDonald, B.W.R. and Godwin, C.I., 1986, Geology of Main Zone at Mount Skukum Wheaton River area, southern Yukon. *In:* Yukon Geology Vol. 1, Exploration and Geological Services Division, Indian and Northern Affairs Canada.
- McDonald, B.W.R., Godwin, C.I. and Stewart, E.B., 1986, Exploration geology of the Mount Skukum epithermal gold deposit, southwestern Yukon. *In:* Yukon Geology Vol. 1, Exploration and Geological Services Division, Yukon, Indian and Northern Affairs Canada.
- McDonald, B.W.R., 1990, Geology and Genesis of the Mount Skukum Epithermal Gold-Silver Deposits, southwestern Yukon Territory. Exploration and Geological Services Division, Yukon, Indian and Northern Affairs, Canada, Bulletin 2, 65 p.
- Meinert, L.D., 1986, Gold in skarns of the Whitehorse Copper Belt. *In:* Yukon Geology - Vol. 1, Exploration and Geological Services Division, Yukon, Indian and Northern Affairs Canada.
- Melis Engineering Limited, 1993, Skukum Creek gold/silver deposit - Rainbow zone, bioleach and cyanidation gold/silver recovery, Phase 1 test program (for Wheaton River Minerals Limited). Exploration and Geological Services Division, Yukon, Indian and Northern Affairs Canada, Canada/Yukon Economic Development Agreement, Technology Open File 1993-10(T).
- Morison, S.R. and McKenna, K., 1981, Surficial Geology and Soils, Southern Lakes Study. Department of Renewable Resources, Yukon Government.
- Morison, S.R. and Klassen, R.W., 1991, Surficial Geology, Whitehorse, Yukon Territory. Geological Survey of Canada, Map 12-1990, (scale 1:250,000).
- Morrison, G.W., 1978, Granite rocks and associated mineral deposits of the Whitehorse map-area, Yukon. *In:* Mineral Industry Report 1975, Indian and Northern Affairs Canada.
- Morrison, G.W., 1979, Metallogenic Map, Whitehorse map-area, Yukon. Exploration and Geological Services Division, Indian and Northern Affairs Canada, Open File 1979-6.
- Morrison, G.W., 1981, Setting and origin of skarn deposits in the Whitehorse Copper Belt, Yukon. Unpublished Ph.D. thesis, University of Western Ontario.
- Morrison, G.W., Godwin, C.I. and Armstrong, R.L., 1981, Interpretation of isotopic ages and $^{87}\text{Sr}/^{86}\text{Sr}$ initial ratios for the plutonic rock in the Whitehorse Copper Belt, Yukon. Canadian Journal of Earth

Sciences, Vol. 16.

- Mougeot, C.M. and Smith, C.A.S., 1992, Soils of the Whitehorse Area, Takhini Valley, Yukon Territory. Agriculture Canada.
- Mougeot, C.M. and Smith, C.A.S., in press, Soils of the Whitehorse Area, Carcross Valley, Yukon Territory. Agriculture Canada.
- Murray, D.W. and Smith, C.A.S., 1988, City of Whitehorse Soil Survey: Takhini Valley area. Agriculture Development Series Report #5, Agriculture Canada, Land Resources Research Center, Box 2703, Whitehorse, Yukon.
- Pride, M.J. 1984, Interlayered sedimentary-volcanic sequence, Mount Skukum volcanic complex. *In:* Yukon Exploration and Geology 1993, Exploration and Geological Services Division, Yukon, Indian and Northern Affairs Canada.
- Pride, M.J. 1985, Preliminary geology map of Mount Skukum volcanic complex, Yukon Territory. Exploration and Geological Services Division, Yukon, Indian and Northern Affairs Canada, Open File 1985-2.
- Pride, M.J. 1986, Description of the Mt. Skukum volcanic complex in southern Yukon. *In:* Yukon Geology Vol. 1, Exploration and Geological Services Division, Yukon, Indian and Northern Affairs Canada.
- Public Works, 1969, Inventory-gravel and binder deposits: Mile 628 to Mile 898, Alaska Highway, Yukon Territory. Department of Public Works, Pacific Region, Government of Canada.
- Rampton, V.N. 1972, Surficial deposits, Alaska Highway, Whitehorse, to Alaska-Yukon boundary. Geological Survey of Canada, Paper 72-1.
- Roots, C.F., 1981, Geologic setting of gold-silver veins on Montana Mountain. *In:* Yukon Exploration and Geology 1979-80, Exploration and Geological Services Division, Indian and Northern Affairs Canada.
- Roots, C.F., 1982, Geology of the Montana Mountain area. Unpublished M.Sc. thesis, Carlton University.
- Rostad, H.P.W., Kozak, L.M. and Acton, D.F., 1977, Soil survey and land evaluation of the Yukon Territory. Department of Indian Affairs and Northern Development, Northern Environmental and Renewable Resources Branch, Land Management Division, Whitehorse, Yukon.
- Rucker, D.P., 1988, Fluid inclusion and ^{18}O study of the precious metal-bearing veins of the Wheaton River District, Yukon. Unpublished M.Sc. thesis, University of Alberta.
- Smith, M.J., 1982, Petrology and geology of high-level rhyolite intrusions of the Skukum area, Yukon Territory. *In:* Yukon Exploration and Geology 1981, Exploration and Geological Services Division, Yukon, Indian and Northern Affairs Canada.
- Smith, M.J., 1983, The Skukum volcanic complex: Geology and comparison to the Bennett Lake Cauldron Complex. *In:* Yukon Exploration and Geology 1982, Exploration and Geological Services Division, Yukon, Indian and Northern Affairs Canada.
- Tenney, D., 1981 The Whitehorse Copper Belt: Mining Exploration and Geology 1967-1980. Exploration and Geological Services Division, Yukon, Indian and Northern Affairs Canada, Bulletin 1, 29 p.
- Underwood McLellan Ltd., 1983, Yukon River Basin Flood Risk Study. Yukon River Basin Study, Hydrology Work Group Report No. 1, Environment Canada, Inland Waters Directorate, Government of Canada.

DIAND library
Call Number: GB1399.5 C3 U64
(from 65°N to Atlin Lake in B.C.)

- Von Gaza, P. and Eyton, J.R., 1988, Enhanced Landsat thematic mapper imagery for exploration geology in the Wheaton River district, southern Yukon. *In*: Yukon Geology, Vol. 2, Exploration and Geological Services Division, Yukon, Indian and Northern Affairs Canada.
- Walton, L.A., 1987, Geology and geochemistry of the Venus Au-Ag-Pb-Zn vein deposit Yukon Territory. Unpublished M.Sc. thesis, University of Alberta.
- Wamsley, M.E., 1981, Terrain classification and interpretive maps, Cowley Lake Area, Yukon Territory. Pedology Consultants. 1:5000 scale maps.
- Watson, P.H., 1984, The Whitehorse Copper Belt - a compilation, Yukon Territory. Exploration and Geological Services Division, Yukon, Indian and Northern Affairs Canada, Open File 1984-1.
- Wheeler, J.O., 1952, Geology and mineral deposits of the Whitehorse map-area, Yukon Territory. Geological Survey of Canada, Paper 52-30.
- Wheeler, J.O., 1961, Whitehorse map-area, Yukon Territory. Geological Survey of Canada, Memoir 312.
- Wheeler, J.O., Brookfield, A.J., Gabrielse, H., Monger, J.W.H., Tipper, H.W. and Woodsworth, G.J., 1991, Terrane map of the Canadian Cordillera. Geological Survey of Canada, Map 1713.
- Wheeler, J.O. and McFeely, P., 1991, Tectonic Assemblage map of the Canadian Cordillera and adjacent parts of the United States of America. Geological Survey of Canada, Map 1712A.
Includes all of Yukon
- Yukon Renewable Resources, 1990, Watson-Wheaton Resource Inventory. Parks, Resources and Regional Planning, Renewable Resources, Government of Yukon.