



Manitoba
Health



Manitoba Health Statistical Update on HIV/AIDS

1985 - June 2002

**Communicable
Disease Control Unit
Public Health**

MANITOBA HEALTH STATISTICAL UPDATE ON HIV/AIDS 1985 TO JUNE 2002


HIV – January 1, 1985 to June 30, 2002

Between January 1 and June 30, 2002, 32 newly diagnosed cases of HIV were reported in Manitoba; 18 males and 14 females, bringing the total number of cases to 947 since 1985 (Table 1, attached report). While females represent 19% of all HIV cases reported since 1985, 8% of HIV positive individuals were accounted for by females between 1985 and 1994 as compared to 29% between 1995 and June 2002 (Table 1, attached report). The majority of all new cases, both male and female, were between the ages of 20 and 39 years (Figure 2, attached report).

Between 1985 and June 2002, 83% (n=789) of all HIV cases reported (at the time of testing) were residents of Winnipeg, while 13% (n=125) of cases resided outside of Winnipeg (Figure 3, attached report). Of the total cases, 3% (n=29) of individuals were from out of province while less than 1% (n=4) of individuals reported missing or unknown geographic information.

With the exception of 2001, there has been a gradual but consistent increase in the percentage of cases residing outside of Winnipeg (see figure below) over recent years. This observation has important implications regarding the availability of HIV prevention and education resources outside of the major urban centre. Further, this finding encourages health care providers to continue to offer HIV testing and counseling.

**Percentage of HIV Positive Cases* by Region of Residence,
Manitoba, January 1985 - June 2002**



Note: Dotted lines reflect half a year of data.

*Cases residing out-of-province or of unknown residence (at the time of testing) are excluded from the denominator.

Self-Reported Ethnicity


As presented in the charts below, 41% (13/32 cases) of newly diagnosed cases of HIV between January and June 2002 were self-reported as Aboriginal at the time of follow-up, while 19% (6/32 cases) were self-reported as Caucasian. These numbers increase to 50% and 23% respectively when cases with unknown or missing ethnicity are excluded (n=6) due to incomplete or missing *Notification of HIV Infection* forms. When these groups are further examined by mode of transmission, the most commonly reported category for Aboriginals includes IDU¹ (10/13 cases; 77%). For Caucasians, the majority of individuals reported MSM² (3/6; 50%).

Between January 1999 and December 2001, the majority of new HIV cases self-reported as Aboriginal (75/195 cases, 38%) and Caucasian (66/195 cases, 34%). These values increase to 45% and 40%, respectively, when cases with missing or unknown ethnicity are excluded (n=29). Similar patterns regarding the predominant modes of transmission were observed between 1999 and 2001. For Aboriginals, the two most common transmission categories include IDU (41/75 cases; 55%) and heterosexual activity with person(s) at increased risk of HIV (23/75 cases; 31%). The most commonly reported categories for Caucasians include MSM (26/66 cases; 39%) and heterosexual activity with person(s) at increased risk of HIV (24/66 cases; 36%).


It is important to note that these data are self-reported and reflect individuals coming forward for testing. Misclassification may occur when the index case fails to self-identify, leading to under-representation. In addition, approximately 15% of HIV cases reported between January 1999 and June 2002 were incomplete due to missing or unknown information pertaining to ethnicity.

Despite these limitations, this information is important as it further characterizes at-risk populations to support targeted HIV prevention and planning initiatives. In addition, this information may be used to facilitate the allocation of resources for education and treatment programs within Regional Health Authorities, other health care jurisdictions and the province.

Percentage of New Positive HIV Cases by Ethnicity in Manitoba, January 2002 to June 2002 (N=32)



Percentage of New Positive HIV Cases by Ethnicity in Manitoba, January 1999 to December 2001 (N=195)



¹ Injection drug use.

² Men having sex with men.


HIV Transmission Patterns

Of the 14 females testing HIV positive between January and June 2002, the predominant modes of transmission after excluding those with no identified risk (n=1; 7%), were injection drug use (6/13 cases; 46%) and endemic³ (5/13 cases; 39%). Of the 18 males and excluding those with no identified risk (n=2; 11%), the predominant modes of transmission included IDU (7/16; 44%), MSM (4/16; 25%) and heterosexual activity with a person(s) at risk of HIV infection (4/16; 25%).

Table 4 (attached report) describes the mode of transmission for all HIV positive cases. When cases are reviewed from 1985 to June 2002, the most common transmission categories for females include heterosexual activity with person(s) at risk of HIV and IDU. For males, the primary modes of transmission include MSM, heterosexual activity with person(s) at risk of HIV and IDU (Figure 4, attached report).

In total, MSM, IDU and heterosexual activity with person(s) at risk of HIV represent roughly 90% of all HIV antibody positive individuals diagnosed between January 1985 and June 2002 (excluding cases with missing/unknown mode of transmission; n=45). There has been a substantial increase in IDU from 29% in 1999 to 41% in 2001 and 45% between January and June 2002 (refer to graph below). Examined over time, it is evident that the proportion of individuals reporting MSM has declined since 1985 and dropped substantially in 1995. However, the *number* of new cases reporting MSM has remained relatively constant over the last five years, with an average of 13 to 15 cases per year, with the exception of 2001 where there were only 9 cases reported as MSM. The proportion of cases reporting heterosexual activity with person(s) at increased risk of HIV has increased steadily since 1995 reaching a high of nearly 40% in 2000.

Risk Profile for HIV Positive Cases in Manitoba, January 1985 - June 2002



Note: Dotted lines reflect half a year of data.

*Cases with no identified risk (NIR) were excluded from the denominator.

MSM is men having sex with men.

IDU is injection drug use and includes MSM/IDU.

HET is heterosexual activity with person(s) at increased risk of HIV.

³ Endemic includes persons originating from or residing in countries with a high prevalence of HIV.


Heterosexual Contact with Person(s) at Risk of HIV

Between January and June 2002, heterosexual contact with person(s) at risk of HIV was reported as the primary mode of transmission for 6 cases. In 1999, 2000 and 2001 there were 26, 21, 19 cases, respectively.

In 1999, the majority of individuals reported heterosexual contact with an unknown partner (9/26 cases; 35%) and having heterosexual contact with multiple sex partners (8/26 cases; 31%). Similar patterns were observed in 2000 and 2001, where the majority of cases reported having heterosexual contact with multiple partners (9/21 cases; 43% and 9/19; 47%, respectively) and unknown partners (7/21; 33% and 6/19; 32%, respectively). See figure below.

Although these data reflect individuals coming forward for testing and are subject to over- or under-reporting, they are helpful in identifying current trends. This information is critical to support and direct planning and prevention services within and across health care jurisdictions in Manitoba. Finally, the increase in HIV among heterosexuals over recent years may suggest that HIV testing is becoming more acceptable among this population. As a result, it may be timely to increase the targeted promotion of HIV testing among this group. Further, it is important that health care professionals offer HIV testing and counseling to individuals infected with a bacterial sexually transmitted disease (STD) or to those named as a contact to someone infected with a STD. Both the Provincial AIDS Strategy and Provincial STD Strategy provide goals and objectives to facilitate program planning and prevention strategies to reduce the risk and transmission of HIV and STD.

**Heterosexual Contact with Person(s) at Risk of HIV,
Manitoba, January 1999 - June 2002**




Note: Number of responses may not add up to the total number of individuals reporting heterosexual contact since all categories within this variable that apply for each case are recorded.


Age of HIV Positive Cases among MSM Risk Profile

Between January 1995 and June 2002, there was an increase in the age of HIV positive cases reporting MSM as the primary mode of transmission as compared to the earlier time period of 1985-1994. This is particularly evident for those individuals 30 years of age and older, where there was an observed increase, from 58% to 80%. Alternatively, there has been a decrease in cases aged 20 and younger, from 42% during 1985 to 1994 to 20% for the period 1995 to June 2002.

Age of HIV Positive Cases with MSM Risk Profile in Manitoba, January 1995 - June 2002 (N=98)




Age of HIV Positive Cases with MSM Risk Profile in Manitoba, January 1985 - December 1994 (N=321)



Confirmatory HIV Antibody Testing Data

Cadham Provincial Laboratory conducts all HIV antibody testing in Manitoba. Data include those who are non-residents as well as those who did not have a Regional Health Authority recorded on the HIV test requisition. Of the laboratory testing requisitions, 140,400 tests were carried out between 1996 and 2001, ranging from 17,343 in 1996 to 26,579 in 2001. Data extracted from the laboratory's information system illustrate that the majority of those seeking testing in 2001 were between 20 and 39 years of age for both males (4,831/8,311 males; 58%) and females (13,017/18,268 females; 71%). This observation was also observed between 1996 and 2000 (refer to Appendix B). These data reflect the number of HIV antibody tests performed and may not necessarily reflect the total number of individuals who seek testing since an individual may be tested more than once.

Number of HIV Antibody Tests by Age and Gender, Manitoba, 2001




*Excludes bone marrow transplant patients, blood donors and where age was unknown (2% of all tests).

As illustrated in the graph below, the most common reason for testing between 1996 and 2001 was heterosexual contact with person(s) at risk of HIV. However, in recent years, other reasons for testing have become more common, such as prenatal HIV testing. Although homosexual contact, injection drug use and STD, make up a small component of the total reasons for testing, this information is important in identifying testing trends, which consequently may aid in directing appropriate testing and prevention programs. Testing data for these particular categories have remained stable over the last 6 years (see Appendix B).

Manitoba Health continues to focus on increasing the number of individuals seeking HIV antibody testing. By increasing these numbers, more individuals at risk of HIV can be reached. Those who seek testing but are not infected are educated on HIV and on how to prevent infection and those who seek testing and are infected can be educated on how to prevent transmission as well as being connected with care and treatment services more quickly. Currently, evaluation of the provincial non-nominal HIV testing system is under review by the HIV Testing Options Sub-Committee of the Manitoba Advisory Committee on Infectious Diseases. Testing options that are available in other provinces are being reviewed. These include anonymous and nominal HIV testing.

Most Common Reasons for HIV Testing and Year, Manitoba, 1996-2001



Source: Cadham Provincial Laboratory Information System.

Note: The graph does not represent number of individuals, rather it reflects reasons for testing as reported on the requisition.

Note: There has been no hierarchy applied; all reasons for testing as reported on the requisition have been included in the graph.

AIDS – January 1985 to June 30, 2002

Between January 1 and June 30, 2002, 4 new cases of AIDS were reported; all 4 cases were male. These reports bring the total number of cases to 202 since 1985 (Table 5, attached report). The number of reported AIDS cases has declined somewhat over recent years, due in part to early diagnosis and improved treatment of individuals with HIV infection. Seventy-nine percent of individuals reported with AIDS have died. However, delays in reporting of both cases and deaths make it difficult to determine precisely the incidence and mortality rate.

APPENDIX A

Reporting of HIV and AIDS in Manitoba

In Manitoba, HIV testing is non-nominal. A prescribed patient code is assigned when a physician completes the appropriate requisition. This code includes the last two letters of the mother's maiden name, the patient's year of birth, day of birth, gender, regional health authority (as defined by number) and first three characters of the patient's postal code. Prior to August 1998, the former Manitoba Health region (as defined by letter) was assigned to identify the patient's region of residence. As well, postal code was not included.

All HIV antibody testing is carried out at the Cadham Provincial Laboratory (CPL). Positive test results are subsequently reported to the Director of Communicable Disease Control as required by the *Diseases and Dead Bodies Regulation, Public Health Act*. It has been the practice of Communicable Disease Control (CDC) Unit to enter case information into the HIV Database *after* the physician (requesting the test) has verified the test result as a new or existing case. However, there have been delays in the completion of and return of the *Notification of HIV Infection Form* (Appendix B) by health care professionals. Consequently, all HIV positive test results are considered new cases unless otherwise advised by the appropriate health care professional. This practice will avoid the under-reporting of HIV in Manitoba, although, duplicate cases may be included. The CDC Unit continues to work with Regional Health Authorities towards a satisfactory resolution in this regard. A collaborative effort between the Winnipeg Regional Health Authority and the CDC Unit, Manitoba Health has decreased the number of outstanding *Notification of HIV Forms* for 1999 to present.

Twice a year, line-listed data from the HIV Database are extracted and forwarded to the Centre for Infectious Disease Prevention and Control, Health Canada in Ottawa for inclusion within the national report, *HIV and AIDS in Canada*. Although non-nominal, the prescribed patient code is stripped prior to release. Instead, a sequential case number is assigned.

Provincially and nationally, AIDS cases and deaths are reportable by physicians. A federal reporting form, the *AIDS Case Reporting Form*, is used for this purpose. New AIDS cases and deaths are reported to the Director of Communicable Disease Control and subsequently forwarded to the Centre for Infectious Disease Prevention and Control. The Centre works diligently with other provinces to ensure that there are no duplications in the counting of cases. The variations seen from previous reports with respect to the number of AIDS cases and deaths may be accounted for by delays in reporting as well as the fact that in Manitoba, the database is updated immediately once surveillance staff are notified that a particular case has been accounted for in another province.

APPENDIX B

Number of HIV Antibody Tests by Age and Gender, Manitoba, 1996-2001

GENDER	AGE GROUP	YEAR						TOTAL
		2001	2000	1999	1998	1997	1996	
Male	other*	319	312	138	35	42	48	894
	<15	140	123	144	135	131	119	792
	15-19	611	610	558	569	597	516	3461
	20-29	2728	2664	2684	2633	2544	2522	15775
	30-39	2103	2121	2072	2102	2027	1940	12365
	40-49	1311	1291	1310	1147	1114	988	7161
	50+	1099	974	1117	918	763	745	5616
	Total	8311	8095	8023	7539	7218	6830	46016
Female	other	90	70	63	30	23	32	308
	<15	238	242	238	162	243	173	1296
	15-19	2360	2268	2258	2269	1934	1419	12508
	20-29	7998	7552	7576	7144	6261	4453	40984
	30-39	5019	4782	4931	4679	4031	2663	26105
	40-49	1603	1495	1552	1281	1175	1078	8184
	50+	960	776	1207	720	641	647	4951
	Total	18268	17185	17825	16285	14308	10465	94336
Total	other	409	382	201	65	65	80	1202
	<15	378	365	382	297	374	292	2088
	15-19	2971	2878	2816	2838	2531	1935	15969
	20-29	10726	10216	10260	9777	8805	6975	56759
	30-39	7122	6903	7003	6781	6058	4603	38470
	40-49	2914	2786	2862	2428	2289	2066	15345
	50+	2059	1750	2324	1638	1404	1392	10567
	Total	26579	25280	25848	23824	21526	17343	140400

Source: Cadham Provincial Laboratory Information System.

* Other refers to bone marrow transplant and donor patients tested as well where age was unknown.

**HIV Antibody Testing Patterns by Reason for Testing and Year,
Manitoba, 1996-2001***

Reason for Testing	2001	2000	1999	1998	1997	1996	TOTAL
Heterosexual	10782	11258	11998	10283	6157	6230	56708
Patient Request	10063	9548	8994	5334	4027	3710	41676
Prenatal	8031	7182	7264	6702	5219	1590	35988
Other	6577	5856	6663	4618	4031	3925	31670
STD	875	909	802	747	597	385	4315
IDU	565	632	681	719	675	669	3941
Homosexual	426	474	430	435	468	510	2743
No Reason Indicated	324	372	353	217	164	142	1572
Bisexual	0	2	14	163	206	206	591
Total**	37643	36233	37199	29218	21544	17367	179204
Total Requisitions	26579	25280	25848	23824	21526	17343	140400

Source: Cadham Provincial Information System

*There has been no hierarchy applied; all reasons for testing as reported on the requisitions have been included in the table.

**This total equals the number of individuals who reported a specific reason for testing and is not equivalent to the total number of requisitions since there may or may not be more than one reason indicated on a requisition.

NOTE:

Heterosexual reflects HET, MSP, SBM, SHP, SIV, SPG, SPR, SUP.

Patient request denotes those who request test and no risk is identified (TRP).

Prenatal reflects those who were tested as part of prenatal care (PRE).

Other reflects ACP, BLT, CLR, GLI, GLO, HPL, HRP, OFF, OIE, OTH, SAS and 18% of the total reasons for testing reported between 1996 and 2001; roughly 18% per year.

STD reflects individuals who were tested because they had a sexually transmitted disease.

IDU reflects those who reported injected drug use (IVD, BII, HOI).

Homosexual reflects HOM, MSM, HOI

Bisexual reflects BIS, BII

For acronym definitions, refer to the table entitled, "Laboratory Coding for HIV Testing Requisition".

**Reasons for Testing (Transmission Categories) as
Defined on the HIV Testing Requisition**

GROUP	DESCRIPTION
ACP	accident/parental exposure
BII	bisexual/IVDA
BIS	bisexual/IVDA
BLT	blood/blood product recipient
CLR	clotting factor recipient
GLI	global/IVDA
GLO	global endemic
HET	heterosexual
HOI	homosexual/IVDA
HOM	homosexuals
HPL	hemophiliacs
HRP	high risk partner
IVD	IVDA
MSM	men who have sex with men
MSP	multiple sex partners
OFF	offspring
OIE	other injury exposure (splashes, bites)
OTH	other risks
PRE	prenatal
SAS	sexual assault
SBM	sex contact with bisexual male
SHP	sex with HIV positive person
SIV	sex with IVDA person
SPG	sex with person from global endemic area
SPR	sex with prostitute
STD	current sexually transmitted disease
SUP	sex with unknown partner
TRP	patient request - no risk identified
NO REASON	no reason indicated

APPENDIX C

NOTIFICATION OF HIV INFECTION (Form prescribed pursuant to subsection 43(2) The Public Health Act: P210)

DESIGNATED PATIENT CODE _____ PHYSICIAN NAME _____
 (As per CPL requisition: Last two initials of mother's maiden name; year of birth; day of birth; gender; RHA of residence code; 3-digit forward sortation postal code)

LABORATORY REQUISITION NUMBER _____ SPECIMEN DATE _____ / _____ / _____
 yyyy mm dd

PRINCIPAL REASON FOR TEST (ONE ONLY)

- Requested by patient (no risk identified)
- Risk factor present (asymptomatic)
- Symptomatic STD work-up
- Travel Insurance
- Prenatal
- Other (specify) _____

GENDER Male Female Trans-gender

If female, pregnant? Yes No
 Receiving anti-retroviral drug(s)? Yes No

M/S Unmarried Married/CL S/D/W

COUNTRY OF BIRTH

Canada Other _____
 If other, year of arrival in Canada _____

ETHNICITY

- Caucasian African/African-American
- Aboriginal Asian
- Other _____

If aboriginal, treaty status: Treaty Non-treaty
 Band number: _____

CLINICAL STATUS

Are HIV-related symptoms present? Yes No

Does the patient have AIDS? Yes No

PAST HISTORY

1) Previous HIV testing? Yes No Unknown
 If yes:
 Date of most recent negative test: _____
 Date of first positive test: _____

2) History of STD ever Yes No

3) STD in past 3 months Yes No

4) Previous blood or tissue donation Yes No

If yes, most recent date _____

Location _____

RISK INFORMATION

(Since 1978; check all client characteristics that apply)

	Yes	No	Unk.
1) <u>Has had sex with:</u>			
A male	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
A female	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
2) <u>Has had heterosexual sex with:</u>			
A bisexual partner	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
An unknown partner	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Multiple sex partners	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
A sex trade worker	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
A client of a sex trade worker (i.e. patient is a sex trade worker)	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
A person with known/suspected HIV	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
An injection drug user	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
A person from an HIV endemic area	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Another high risk partner	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
3) <u>Has used needles for recreational (non-medical) drug injection</u>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
4) <u>Has received blood or blood products</u>			
a) Prior to Nov. 1985	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b) After Nov. 1985	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
5) <u>Has received blood or blood products for treatment of a coagulation disorder</u>			
a) Prior to Nov. 1985	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
b) After Nov. 1985	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
6) <u>Has been exposed to HIV in an occupational setting (e.g. needlestick injury)</u>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
7) <u>Born to an HIV positive mother</u>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
8) <u>Born in or resident of an HIV-endemic country</u>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
9) <u>Has had:</u>			
<input type="checkbox"/> tattoo			
<input type="checkbox"/> body piercing			
<input type="checkbox"/> acupuncture			
<input type="checkbox"/> blood contact from bite, altercation, etc.			
10) <u>Other exposure which could have been source of HIV infection, specify</u> _____			
11) <u>No identifiable risk factor</u> <input type="checkbox"/>			

Interview for partners at risk to be done by: Physician Yes No Public Health Nurse Yes No

If by public health nurse, physician must first obtain informed consent from client. Has informed consent been obtained? Yes No

CONTACT INFORMATION ON PARTNERS TO BE FOLLOWED BY PUBLIC HEALTH:

Name _____ Home tel _____ Work tel _____
 Alias _____ Sex F M
 Address _____ Postal Code _____ Age/Birth date _____
 Occupation _____ Place of Employment/School _____
 Live-In Partner Single Other _____ Lives with Parents Informant Other
 Characteristics: Height _____ Wt. _____ Eye Colour _____ Hair _____ Complexion _____
 Sexual Exposure: (First) _____ To _____ (Last) Parenteral(First) _____ To _____ (Last)
 Notified: Yes Date _____ No By Whom _____



Manitoba Health Statistical Update on HIV/AIDS

1985 - JUN 2002

CDC Unit
Public Health Branch
Manitoba Health



MANITOBA HEALTH

Table 1. NUMBER OF INDIVIDUALS TESTING HIV ANTIBODY POSITIVE, 1985-2002*

Year	Male	Female	Total
1985	3	0	3
1986	70	0	70
1987	50	3	53
1988	37	4	41
1989	57	3	60
1990	40	3	43
1991	33	6	39
1992	39	6	45
1993	54	4	58
1994	50	7	57
1995	42	9	51
1996	37	16	53
1997	59	17	76
1998	53	18	71
1999	53	20	73
2000	38	19	57
2001	39	26	65
2002	18	14	32
Total	772	175	947

* to 06/30/2002

Figure 1. NUMBER OF INDIVIDUALS TESTING HIV ANTIBODY POSITIVE, 1985-2002*





Table 2. NUMBER OF INDIVIDUALS TESTING HIV ANTIBODY POSITIVE BY AGE AND GENDER, 2002*, 2001. CUMULATIVE 1985-2000 AND 1985-2002*

Age	Gender	2002*	2001	1985-2000	1985-2002*
		JAN-JUN	Total	Total	Total
<15	M	0	0	6	6
	F	1	0	4	5
15-19	M	1	0	14	15
	F	1	0	9	10
20-29	M	2	2	246	250
	F	5	11	64	80
30-39	M	9	21	283	313
	F	6	9	38	53
40-49	M	3	5	109	117
	F	0	4	13	17
50+	M	3	11	57	71
	F	1	2	7	10
Total	M	18	39	715	772
	F	14	26	135	175

* to 06/30/2002

Figure 2. NUMBER OF INDIVIDUALS TESTING HIV ANTIBODY POSITIVE BY AGE AND GENDER, 1985-2002*




* to 06/30/2002

Table 3. NUMBER OF INDIVIDUALS TESTING HIV ANTIBODY POSITIVE BY GEOGRAPHIC RESIDENCE AND GENDER, 2002*, 2001, CUMULATIVE 1985-2000 AND 1985-2002*

Geographic Residence	Gender	2002*	2001	1985-2000	1985-2002*
		JAN-JUN	Total	Total	Total
Winnipeg	M	16	33	594	643
	F	12	23	111	146
Brandon	M	0	0	1	1
	F	1	0	0	1
Marquette	M	0	0	1	1
Brandon, Marquette, S. Westman	M	0	0	18	18
	F	0	0	1	1
N. Eastman	M	0	1	1	2
	F	0	0	1	1
S. Eastman	M	0	0	1	1
N. Eastman, S. Eastman	M	0	0	11	11
	F	0	0	2	2
Interlake	M	0	2	24	26
	F	1	0	3	4
Central	M	2	1	26	29
	F	0	3	5	8
Parkland	M	0	0	5	5
	F	0	0	1	1
Norman	M	0	1	3	4
	F	0	0	1	1
Burntwood	M	0	0	6	6
	F	0	0	2	2
Unknown	M	0	1	2	3
	F	0	0	1	1
Out of Province	M	0	0	22	22
	F	0	0	7	7
Total	M	18	39	715	772
	F	14	26	135	175

* to 06/30/2002

Figure 3. PERCENTAGE OF HIV POSITIVE INFECTIONS IN MANITOBA BY REGION OF RESIDENCE AND GENDER, 1985-2002*



* to 06/30/2002

Table 4. NUMBER OF INDIVIDUALS TESTING HIV ANTIBODY POSITIVE BY TRANSMISSION CATEGORY AND GENDER, 2002*, 2001, CUMULATIVE 1985-2000 AND 1985-2002*

Transmission Category	Gender	2002*	2001	1985-2000	1985-2002*
		JAN-JUN	Total	Total	Total
MSM	M	4	9	406	419
MSM/IDU	M	0	1	46	47
Heterosexual'	M	4	9	105	118
	F	2	10	57	69
IDU	M	7	13	82	102
	F	6	10	47	64
Perinatal	M	0	0	2	2
	F	0	0	1	1
Recp B/B products	M	1	0	28	29
	F	0	2	6	8
Endemic''	M	0	2	20	22
	F	5	3	13	21
NIR	M	2	5	26	33
	F	1	1	10	12
Total	M	18	39	715	772
	F	14	26	135	175

MSM = men having sex with men

IDU = injection drug use

Recp B/B products = recipient of blood/blood product


NIR = No Identified Risk

' Heterosexual activity includes persons reporting heterosexual activity with person(s) at risk of HIV infection

'' Endemic includes persons originating from or residing in countries with a high prevalence of HIV

* to 06/30/2002

Figure 4. PERCENTAGE OF HIV POSITIVE INFECTIONS IN MANITOBA BY TRANSMISSION CATEGORY, 1985-2002*



* to 06/30/2002


Table 5. NUMBER OF REPORTED AIDS CASES AND DEATHS, 1985-2002*

Year	Cases Reported"	Deaths Reported"
1985	2	0
1986	14	5
1987	8	7
1988	5	5
1989	17	11
1990	11	9
1991	21	10
1992	16	13
1993	15	22
1994	12	17
1995	18	16
1996	9	7
1997	7	5
1998	11	5
1999	12	8
2000	12	11
2001	8	7
2002	4	2
Total	202	160

"Because of delays in reporting, the number of reported cases and deaths does not necessarily represent the number of cases diagnosed or deaths occurring during the period.

*to 06/30/2002

Figure 5. NUMBER OF REPORTED AIDS CASES AND DEATHS, 1985-2002*




*to 06/30/2002

Table 6. NUMBER OF REPORTED CASES OF AIDS BY AGE AND GENDER, 2002*, 2001, CUMULATIVE 1985-2000 AND 1985-2002*

Age	Gender	2002*	2001	1985-2000	1985-2002*
		JAN-JUN	Total	Total	Total
<15	M	0	0	1	1
	F	0	0	1	1
15-19	M	0	0	2	2
	F	0	0	0	0
20-29	M	0	1	25	26
	F	0	1	2	3
30-39	M	1	1	86	88
	F	0	0	6	6
40-49	M	3	2	35	40
	F	0	1	3	4
50+	M	0	2	27	29
	F	0	0	2	2
Total	M	4	6	176	186
	F	0	2	14	16

*to 06/30/2002

Figure 6. NUMBER OF REPORTED CASES OF AIDS BY AGE AND GENDER, 1985-2002*




*to 06/30/2002

Table 7. NUMBER OF REPORTED CASES OF AIDS BY GEOGRAPHIC RESIDENCE AND GENDER, 2002*, 2001, CUMULATIVE 1985-2000 AND 1985-2002*

Geographic Residence	Gender	2002*	2001	1985-2000	1985-2002*
		JAN-JUN	Total	Total	Total
Winnipeg	M	4	0	159	168
	F	0	1	13	14
Brandon	M	0	5	3	3
	F	0	0	0	0
Marquette	M	0	0	2	2
	F	0	0	0	0
S. Westman	M	0	0	1	1
	F	0	0	0	0
N. Eastman	M	0	0	0	0
	F	0	0	0	0
S. Eastman	M	0	0	1	1
	F	0	0	0	0
Interlake	M	0	0	3	3
	F	0	0	0	0
Central	M	0	0	0	0
	F	0	0	0	0
Parkland	M	0	1	2	3
	F	0	0	1	1
Norman	M	0	0	1	1
	F	0	0	0	0
Burntwood and Churchill	M	0	0	1	1
	F	0	1	0	1
Unknown	M	0	0	0	0
	F	0	0	0	0
Out of Province	M	0	0	3	3
	F	0	0	0	0
Total	M	4	6	176	186
	F	0	2	14	16

*to 06/30/2002

Figure 7. PERCENTAGE OF AIDS CASES IN MANITOBA BY REGION OF RESIDENCE AND GENDER, 1985-2002*



*to 06/30/2002

Table 8. NUMBER OF REPORTED CASES OF AIDS BY TRANSMISSION CATEGORY AND GENDER, 2002*, 2001, CUMULATIVE 1985-2000 AND 1985-2002*

Transmission Category	Gender	2002*	2001	1985-2000	1985-2002*
		JAN-JUN	Total	Total	Total
MSM	M	3	1	130	134
Heterosexual	M	0	3	24	27
	F	0	2	10	12
IDU	M	0	2	7	9
	F	0	0	1	1
Perinatal	M	0	0	1	1
	F	0	0	1	1
Recp B/B products	M	0	0	14	14
	F	0	0	2	2
Endemic"	M	0	0	0	0
	F	0	0	0	0
NIR	M	1	0	0	1
	F	0	0	0	0
Total	M	4	6	176	186
	F	0	2	14	16

*to 06/30/2002

MSM = men having sex with men

IDU = injection drug use


Recp B/B products = recipient of blood/blood product

NIR = no identified risk

"Heterosexual activity includes persons reporting heterosexual activity with person(s) at risk of HIV infection

"Endemic includes persons originating from or residing in countries with a high prevalence of HIV

Figure 8. PERCENTAGE OF AIDS CASES IN MANITOBA BY TRANSMISSION CATEGORY AND GENDER, 1985-2002*




*to 06/30/2002

Table 9. NUMBER OF REPORTED CASES OF AIDS BY PRIMARY DIAGNOSIS AND GENDER, 2002*, 2001, CUMULATIVE 1985-2000 AND 1985-2002*

Primary Diagnosis	Gender	2002*	2001	1985-2000	1985-2002*
		JAN-JUN	Total	Total	Total
PCP	M	2	2	65	69
	F	0	0	1	1
Kaposi's Sarcoma	M	0	1	23	24
	F	0	0	0	0
Tuberculosis	M	0	0	13	13
	F	0	0	0	0
HIV Encephalopathy	M	0	0	2	2
	F	0	1	0	1
Other Opportunistic Diseases	M	2	2	47	51
	F	0	1	10	11
Other Malignancy	M	0	1	7	8
	F	0	0	0	0
HIV Wasting	M	0	0	17	17
	F	0	0	3	3
Unknown	M	0	0	2	2
	F	0	0	0	0
Total	M	4	6	176	186
	F	0	2	14	16

PCP = pneumocystis carinii pneumonia
 *to 06/30/2002

Figure 9. PERCENTAGE OF AIDS CASES IN MANITOBA BY PRIMARY DIAGNOSIS, 1985-2002*



*to 06/30/2002