



**Canadian Community Epidemiology Network on Drug Use (CCENDU):  
National Mortality Report**

May, 2006

Published by the Canadian Centre on Substance Abuse (CCSA)

Indicator	<b>MORTALITY</b>			
Jurisdiction	<b>Canada</b>			
Source(s) included	<b>Mortality Statistics</b>	<b>2000</b>	<b>2001</b>	<b>2002</b>

---

**For further information, please write:**  
Canadian Centre on Substance Abuse (CCSA)  
Suite 300, 75 Albert St., Ottawa, ON K1P 5E7  
Tel.: (613) 235-4048; fax (613) 235-8101

Visit our website at [www.ccsa.ca](http://www.ccsa.ca)

**Ce document est également disponible en français sous le titre:**  
Réseau communautaire Canadien d'épidémiologie des toxicomanies (RCCET)-Rapport national sur la mortalité

## **About the Vital Statistics- Death Database**

**Source:** Statistics Canada. *Vital Statistics- Death Database: Detailed Information for 2003*.

Available at: <http://www.statcan.ca/english/freepub/84-208-XIE/84-208-XIE2004002.htm>.

### **Population**

The actual (survey) population of the death database is deaths of Canadian residents and non-residents in Canada, and deaths of Canadian residents in some American states.

### **General Notes**

Causes of death are tabulated according to the detailed categories of the *International Statistical Classification of Diseases and Related Health Problems* (ICD-10).

Data users must note that cause of death data coded to different revisions of the classification are not comparable [(e.g., ICD9 codes used prior to 2000 are not comparable to ICD10 codes used from 2000 to present)].

Canadian mortality data reflects the internationally defined underlying cause of death. Nature of injury codes are, by definition, not used for underlying cause of death. ICD9 nature of injury codes (N800-N999) were included in a supplementary table in the Statistics Canada causes of death publications from 1979-1999 however, ICD10 nature of injury codes (S00-T98) are not available from 2000 onward. As a result, the national tables presented on the CCENDU site for 2000, 2001, and 2002 show substantially fewer categories than the previous years for both illicit and alcohol related deaths.

For more information on The Vital Statistics Death Database see, *Vital Statistics- Death Database: Detailed Information for 2003*. Available at:  
<http://www.statcan.ca/english/freepub/84-208-XIE/84-208-XIE2004002.htm>

### Mortality Statistics 2000 - Canada

DIAGNOSIS	ICD10	Males	Females	Total
<b>2000- Illicit related deaths</b>				
Mental and behavioural disorders due to use of opioids, withdrawal state	F11.3	0	0	<b>0</b>
Mental and behavioural disorders due to use of opioids, withdrawal state with delirium	F11.4	0	0	<b>0</b>
Mental and behavioural disorders due to use of cannabinoids, withdrawal state	F12.3	0	0	<b>0</b>
Mental and behavioural disorders due to use of cannabinoids, withdrawal state with delirium	F12.4	0	0	<b>0</b>
Mental and behavioural disorders due to use of sedatives or hypnotics, withdrawal state	F13.3	0	0	<b>0</b>
Mental and behavioural disorders due to use of sedatives or hypnotics, withdrawal state with delirium	F13.4	0	0	<b>0</b>
Mental and behavioural disorders due to use of cocaine, withdrawal state	F14.3	0	0	<b>0</b>
Mental and behavioural disorders due to use of cocaine, withdrawal state with delirium	F14.4	0	0	<b>0</b>
Mental and behavioural disorders due to use of other stimulants, including caffeine, withdrawal state	F15.3	0	0	<b>0</b>
Mental and behavioural disorders due to use of other stimulants, including caffeine, withdrawal state with delirium	F15.4	0	0	<b>0</b>
Mental and behavioural disorders due to use of hallucinogens, withdrawal state	F16.3	0	0	<b>0</b>

**CCENDU-Canadian Community Epidemiology Network on Drug Use**  
**Indicator: Mortality - Canada**

---

Mental and behavioural disorders due to use of hallucinogens, withdrawal state with delirium		F16.4	0	0	<b>0</b>
Mental and behavioural disorders due to use of tobacco, withdrawal state		F17.3	0	0	<b>0</b>
Mental and behavioural disorders due to use of tobacco, withdrawal state with delirium		F17.4	0	0	<b>0</b>
Mental and behavioural disorders due to use of volatile solvents, withdrawal state		F18.3	0	0	<b>0</b>
Mental and behavioural disorders due to use of volatile solvents, withdrawal state with delirium		F18.4	0	0	<b>0</b>
Mental and behavioural disorders due to multiple drug use and use of psychoactive substances, withdrawal state		F19.3	0	0	<b>0</b>
Mental and behavioural disorders due multiple drug use and use of psychoactive substances, withdrawal state with delirium		F19.4	0	0	<b>0</b>
Mental and behavioural disorders due to use of cocaine, dependence syndrome		F14.2	2	0	<b>2</b>
Mental and behavioural disorders due to use of cannabinoids, dependence syndrome		F12.2	0	0	<b>0</b>
Mental and behavioural disorders due to use of cannabinoids, harmful use		F12.1	0	0	<b>0</b>
Mental and behavioural disorders due to use of hallucinogens, harmful use		F16.1	0	0	<b>0</b>
Mental and behavioural disorders due to use of cocaine, harmful use		F14.1	1	2	<b>3</b>
Fetus and newborn affected by maternal use of drugs of addiction		P04.4 (baby)	0	0	<b>0</b>
Intent self-poisoning by and exposure to nonopioid analgesics, antipyretics, and antirheumatics		X60	7	14	<b>21</b>

CCENDU-Canadian Community Epidemiology Network on Drug Use  
Indicator: Mortality - Canada

Intent self-poisoning by and exposure to antiepileptic, sedative-hypnotic, antiparkinsonism, and psychotropic drugs, NEC		X61	52	54	<b>106</b>
Intent self-poisoning by and exposure to narcotics and psychodysleptics, NEC		X62	40	16	<b>56</b>
Intent self-poisoning by and exposure to other drugs acting on autonomic nervous system		X63	4	2	<b>6</b>
Intent self-poisoning by other and unspecified drugs, medicaments and biological substances		X64	162	148	<b>310</b>
<b>Total Deaths Related to Illicit Drug Use</b>			<b>268</b>	<b>236</b>	<b>504</b>

DIAGNOSIS	ICD10	Males	Females	Total
<b>2000- (II)Licit related deaths</b>				
Mental and behavioural disorders due to use of opioids, dependence syndrome	F11.2	4	2	<b>6</b>
Mental and behavioural disorders due to use of other stimulants including caffeine, dependence syndrome	F15.2	0	0	<b>0</b>
Mental and behavioural disorders due to use of hallucinogens, dependence syndrome	F16.2	0	0	<b>0</b>
Mental and behavioural disorders due to multiple drug use and use of psychoactive substances, dependence syndrome	F19.2	5	7	<b>12</b>
Mental and behavioural disorders due to use of sedatives or hypnotics, harmful use	F13.1	1	0	<b>1</b>
Mental and behavioural disorders due to use of opioids, harmful use	F11.1	0	0	<b>0</b>
Mental and behavioural disorders due to use of other stimulants including caffeine, harmful use	F15.1	0	0	<b>0</b>

CCENDU-Canadian Community Epidemiology Network on Drug Use  
Indicator: Mortality - Canada

Mental and behavioural disorders due to multiple drug use and use of psychoactive substances, harmful use		F19.1	22	9	<b>31</b>
Mental disorders and diseases of the nervous system, complicating pregnancy, childbirth, and puerperium	This category includes alcohol related incidences. In other words, it is no longer exclusive to the licit/illicit drug classification.	O99.3 (mom)	0	0	<b>0</b>
Maternal care for (suspected) damage to fetus by drugs		O35.5 (mom)	0	0	<b>0</b>
Accidental poisoning by and exposure to nonopioid analgesics, antipyretics, antirheumatics		X40	9	16	<b>25</b>
Accidental poisoning by and exposure to antiepileptic, sedative-hypnotic, antiparkinsonism, psychotropic drugs, NEC		X41	38	26	<b>64</b>
Accidental poisoning by and exposure to narcotic psychodysleptics, NEC		X42	344	71	<b>415</b>
Accidental poisoning by and exposure to other drugs acting on the autonomic nervous system		X43	2	1	<b>3</b>
Accidental poisoning by other unspecified drugs, medicaments, and biological substances		X44	154	132	<b>286</b>
<b>Total Deaths Related to Illicit &amp; Licit Drug Use</b>			<b>579</b>	<b>264</b>	<b>843</b>

### Mortality Statistics 2000 – Canada (Cont.)

DIAGNOSIS	ICD10	Males	Females	Total
<b>2000- Alcohol related deaths</b>				
Mental and behavioural disorders due to use of alcohol, withdrawal state with delirium	F10.4	9	1	<b>10</b>
Mental and behavioural disorders due to use of alcohol, amnesic syndrome	F10.6	19	10	<b>29</b>
Mental and behavioural disorders due to use of alcohol, residual and late onset psychotic disorder	F10.7	37	9	<b>46</b>
Mental and behavioural disorders due to use of alcohol, psychotic disorder	F10.5	0	0	<b>0</b>
Mental and behavioural disorders due to use of alcohol, acute intoxication	F10.0	38	11	<b>49</b>
Mental and behavioural disorders due to use of alcohol, other mental and behavioural disorders	F10.8	0	0	<b>0</b>
Mental and behavioural disorders due to use of alcohol, unspecified mental and behavioural disorder	F10.9	17	5	<b>22</b>
Mental and behavioural disorders due to use of alcohol, dependence syndrome	F10.2	263	75	<b>338</b>
Mental and behavioural disorders due to use of alcohol, harmful use	F10.1	113	37	<b>150</b>
Alcoholic polyneuropathy	G62.1	0	0	<b>0</b>
Alcoholic cardiomyopathy	I42.6	58	9	<b>67</b>
Alcoholic gastritis	K29.2	2	3	<b>5</b>
Alcoholic fatty liver	K70.0	4	5	<b>9</b>
Alcoholic hepatitis	K70.1	44	23	<b>67</b>
Alcoholic cirrhosis of liver	K70.3	611	175	<b>786</b>
Alcoholic liver disease unspecified	K70.9	74	16	<b>90</b>
Hepatic fibrosis	K74.0	2	1	<b>3</b>

CCENDU-Canadian Community Epidemiology Network on Drug Use  
 Indicator: Mortality - Canada

Other and unspecified cirrhosis of the liver		K74.6	620	377	<b>997</b>
Liver disease unspecified		K76.9	21	19	<b>40</b>
Acute pancreatitis		K85	162	123	<b>285</b>
Alcohol-induced chronic pancreatitis		K86.0	18	5	<b>23</b>
Other chronic pancreatitis		K86.1	14	7	<b>21</b>
Finding of alcohol in blood		R78.0	0	0	<b>0</b>
Accidental poisoning by and exposure to alcohol		X45	50	20	<b>70</b>
Mental disorders and diseases of the nervous system, complicating pregnancy, childbirth, and puerperium	This category includes licit/illicit drug related incidences. In other words, it is no longer exclusive to the alcohol classification.	O99.3 (mom)	0	0	<b>0</b>
<b>Total Deaths Related to Alcohol</b>			<b>2176</b>	<b>931</b>	<b>3107</b>



### Mortality Statistics 2001 - Canada

DIAGNOSIS	ICD10	Males	Females	Total
<b>2001- Illicit related deaths</b>				
Mental and behavioural disorders due to use of opioids, withdrawal state	F11.3	1	0	1
Mental and behavioural disorders due to use of opioids, withdrawal state with delirium	F11.4	0	0	0
Mental and behavioural disorders due to use of cannabinoids, withdrawal state	F12.3	0	0	0
Mental and behavioural disorders due to use of cannabinoids, withdrawal state with delirium	F12.4	0	0	0
Mental and behavioural disorders due to use of sedatives or hypnotics, withdrawal state	F13.3	0	0	0
Mental and behavioural disorders due to use of sedatives or hypnotics, withdrawal state with delirium	F13.4	0	0	0
Mental and behavioural disorders due to use of cocaine, withdrawal state	F14.3	0	0	0
Mental and behavioural disorders due to use of cocaine, withdrawal state with delirium	F14.4	0	0	0
Mental and behavioural disorders due to use of other stimulants, including caffeine, withdrawal state	F15.3	0	0	0
Mental and behavioural disorders due to use of other stimulants, including caffeine, withdrawal state with delirium	F15.4	0	0	0
Mental and behavioural disorders due to use of hallucinogens, withdrawal state	F16.3	0	0	0

**CCENDU-Canadian Community Epidemiology Network on Drug Use**  
**Indicator: Mortality - Canada**

Mental and behavioural disorders due to use of hallucinogens, withdrawal state with delirium		F16.4	0	0	<b>0</b>
Mental and behavioural disorders due to use of tobacco, withdrawal state		F17.3	0	0	<b>0</b>
Mental and behavioural disorders due to use of tobacco, withdrawal state with delirium		F17.4	0	0	<b>0</b>
Mental and behavioural disorders due to use of volatile solvents, withdrawal state		F18.3	0	0	<b>0</b>
Mental and behavioural disorders due to use of volatile solvents, withdrawal state with delirium		F18.4	0	0	<b>0</b>
Mental and behavioural disorders due to multiple drug use and use of psychoactive substances, withdrawal state		F19.3	0	0	<b>0</b>
Mental and behavioural disorders due multiple drug use and use of psychoactive substances, withdrawal state with delirium		F19.4	0	0	<b>0</b>
Mental and behavioural disorders due to use of cocaine, dependence syndrome		F14.2	3	0	<b>3</b>
Mental and behavioural disorders due to use of cannabinoids, dependence syndrome		F12.2	0	0	<b>0</b>
Mental and behavioural disorders due to use of cannabinoids, harmful use		F12.1	0	0	<b>0</b>
Mental and behavioural disorders due to use of hallucinogens, harmful use		F16.1	0	0	<b>0</b>
Mental and behavioural disorders due to use of cocaine, harmful use		F14.1	3	1	<b>4</b>
Fetus and newborn affected by maternal use of drugs of addiction		P04.4 (baby)	0	0	<b>0</b>
Intent self-poisoning by and exposure to nonopioid analgesics, antipyretics, and antirheumatics		X60	12	12	<b>24</b>

**CCENDU-Canadian Community Epidemiology Network on Drug Use**  
**Indicator: Mortality - Canada**

Intent self-poisoning by and exposure to antiepileptic, sedative-hypnotic, antiparkinsonism, and psychotropic drugs, NEC	X61	47	61	<b>108</b>
Intent self-poisoning by and exposure to narcotics and psychodysleptics, NEC	X62	40	26	<b>66</b>
Intent self-poisoning by and exposure to other drugs acting on autonomic nervous system	X63	3	3	<b>6</b>
Intent self-poisoning by other and unspecified drugs, medicaments and biological substances	X64	163	187	<b>350</b>
<b>Total Deaths Related to Illicit Drug Use</b>		<b>272</b>	<b>290</b>	<b>562</b>

<b>DIAGNOSIS</b>	<b>ICD10</b>	<b>Males</b>	<b>Females</b>	<b>Total</b>
<b>2001- (II)Licit related deaths</b>				
Mental and behavioural disorders due to use of opioids, dependence syndrome	F11.2	1	0	<b>1</b>
Mental and behavioural disorders due to use of other stimulants including caffeine, dependence syndrome	F15.2	0	0	<b>0</b>
Mental and behavioural disorders due to use of hallucinogens, dependence syndrome	F16.2	0	0	<b>0</b>
Mental and behavioural disorders due to multiple drug use and use of psychoactive substances, dependence syndrome	F19.2	9	9	<b>18</b>
Mental and behavioural disorders due to use of sedatives or hypnotics, harmful use	F13.1	0	0	<b>0</b>
Mental and behavioural disorders due to use of opioids, harmful use	F11.1	1	2	<b>3</b>

CCENDU-Canadian Community Epidemiology Network on Drug Use  
Indicator: Mortality - Canada

Mental and behavioural disorders due to use of other stimulants including caffeine, harmful use		F15.1	0	0	<b>0</b>
Mental and behavioural disorders due to multiple drug use and use of psychoactive substances, harmful use		F19.1	12	8	<b>20</b>
Mental disorders and diseases of the nervous system, complicating pregnancy, childbirth, and puerperium	This category includes alcohol related incidences. In other words, it is no longer exclusive to the licit/illicit drug classification.	O99.3 (mom)	0	0	<b>0</b>
Maternal care for (suspected) damage to fetus by drugs		O35.5 (mom)	0	0	<b>0</b>
Accidental poisoning by and exposure to nonopioid analgesics, antipyretics, antirheumatics		X40	10	18	<b>28</b>
Accidental poisoning by and exposure to antiepileptic, sedative-hypnotic, antiparkinsonism, psychotropic drugs, NEC		X41	34	26	<b>60</b>
Accidental poisoning by and exposure to narcotic psychodysleptics, NEC		X42	350	89	<b>439</b>
Accidental poisoning by and exposure to other drugs acting on the autonomic nervous system		X43	0	4	<b>4</b>
Accidental poisoning by other unspecified drugs, medicaments, and biological substances		X44	162	107	<b>269</b>
<b>Total Deaths Related to Illicit &amp; Licit Drug Use</b>			<b>579</b>	<b>263</b>	<b>842</b>

**Mortality Statistics 2001 – Canada (Cont.)**

DIAGNOSIS	ICD10	Males	Females	Total
<b>2001- Alcohol related deaths</b>				
Mental and behavioural disorders due to use of alcohol, withdrawal state with delirium	F10.4	7	1	<b>8</b>
Mental and behavioural disorders due to use of alcohol, amnesic syndrome	F10.6	20	2	<b>22</b>
Mental and behavioural disorders due to use of alcohol, residual and late onset psychotic disorder	F10.7	38	13	<b>51</b>
Mental and behavioural disorders due to use of alcohol, psychotic disorder	F10.5	0	1	<b>1</b>
Mental and behavioural disorders due to use of alcohol, acute intoxication	F10.0	54	18	<b>72</b>
Mental and behavioural disorders due to use of alcohol, other mental and behavioural disorders	F10.8	0	0	<b>0</b>
Mental and behavioural disorders due to use of alcohol, unspecified mental and behavioural disorder	F10.9	21	5	<b>26</b>
Mental and behavioural disorders due to use of alcohol, dependence syndrome	F10.2	267	71	<b>338</b>
Mental and behavioural disorders due to use of alcohol, harmful use	F10.1	123	36	<b>159</b>
Alcoholic polyneuropathy	G62.1	3	0	<b>3</b>
Alcoholic cardiomyopathy	I42.6	56	10	<b>66</b>
Alcoholic gastritis	K29.2	4	0	<b>4</b>
Alcoholic fatty liver	K70.0	3	6	<b>9</b>
Alcoholic hepatitis	K70.1	47	25	<b>72</b>
Alcoholic cirrhosis of liver	K70.3	614	185	<b>799</b>
Alcoholic liver disease unspecified	K70.9	61	20	<b>81</b>
Hepatic fibrosis	K74.0	2	2	<b>4</b>

**CCENDU-Canadian Community Epidemiology Network on Drug Use**  
**Indicator: Mortality - Canada**

Other and unspecified cirrhosis of the liver		K74.6	626	424	<b>1050</b>
Liver disease unspecified		K76.9	35	20	<b>55</b>
Acute pancreatitis		K85	145	134	<b>279</b>
Alcohol-induced chronic pancreatitis		K86.0	20	3	<b>23</b>
Other chronic pancreatitis		K86.1	11	9	<b>20</b>
Finding of alcohol in blood		R78.0	0	0	<b>0</b>
Accidental poisoning by and exposure to alcohol		X45	58	30	<b>88</b>
Mental disorders and diseases of the nervous system, complicating pregnancy, childbirth, and puerperium	This category includes licit/illicit drug related incidences. In other words, it is no longer exclusive to the alcohol classification.	O99.3 (mom)	0	0	<b>0</b>
<b>Total Deaths Related to Alcohol</b>			<b>2215</b>	<b>1015</b>	<b>3230</b>

### Mortality Statistics 2002 – Canada

DIAGNOSIS	ICD10	Males	Females	Total
<b>2002- Illicit related deaths</b>				
Mental and behavioural disorders due to use of opioids, withdrawal state	F11.3	0	0	0
Mental and behavioural disorders due to use of opioids, withdrawal state with delirium	F11.4	0	0	0
Mental and behavioural disorders due to use of cannabinoids, withdrawal state	F12.3	0	0	0
Mental and behavioural disorders due to use of cannabinoids, withdrawal state with delirium	F12.4	0	0	0
Mental and behavioural disorders due to use of sedatives or hypnotics, withdrawal state	F13.3	0	0	0
Mental and behavioural disorders due to use of sedatives or hypnotics, withdrawal state with delirium	F13.4	0	0	0
Mental and behavioural disorders due to use of cocaine, withdrawal state	F14.3	0	0	0
Mental and behavioural disorders due to use of cocaine, withdrawal state with delirium	F14.4	0	0	0
Mental and behavioural disorders due to use of other stimulants, including caffeine, withdrawal state	F15.3	0	0	0
Mental and behavioural disorders due to use of other stimulants, including caffeine, withdrawal state with delirium	F15.4	0	0	0

**CCENDU-Canadian Community Epidemiology Network on Drug Use**  
**Indicator: Mortality - Canada**

Mental and behavioural disorders due to use of hallucinogens, withdrawal state		F16.3	0	0	<b>0</b>
Mental and behavioural disorders due to use of hallucinogens, withdrawal state with delirium		F16.4	0	0	<b>0</b>
Mental and behavioural disorders due to use of tobacco, withdrawal state		F17.3	0	0	<b>0</b>
Mental and behavioural disorders due to use of tobacco, withdrawal state with delirium		F17.4	0	0	<b>0</b>
Mental and behavioural disorders due to use of volatile solvents, withdrawal state		F18.3	0	0	<b>0</b>
Mental and behavioural disorders due to use of volatile solvents, withdrawal state with delirium		F18.4	0	0	<b>0</b>
Mental and behavioural disorders due to multiple drug use and use of psychoactive substances, withdrawal state		F19.3	0	0	<b>0</b>
Mental and behavioural disorders due multiple drug use and use of psychoactive substances, withdrawal state with delirium		F19.4	0	0	<b>0</b>
Mental and behavioural disorders due to use of cocaine, dependence syndrome		F14.2	3	1	<b>4</b>
Mental and behavioural disorders due to use of cannabinoids, dependence syndrome		F12.2	0	0	<b>0</b>
Mental and behavioural disorders due to use of cannabinoids, harmful use		F12.1	0	0	<b>0</b>
Mental and behavioural disorders due to use of hallucinogens, harmful use		F16.1	0	0	<b>0</b>
Mental and behavioural disorders due to use of cocaine, harmful use		F14.1	2	0	<b>2</b>
Fetus and newborn affected by meternal use of drugs of addicton		P04.4 (baby)	0	0	<b>0</b>



**CCENDU-Canadian Community Epidemiology Network on Drug Use**  
**Indicator: Mortality - Canada**

Intent self-poisoning by and exposure to nonopioid analgesics, antipyretics, and antirheumatics		X60	13	12	<b>25</b>
Intent self-poisoning by and exposure to antiepileptic, sedative-hypnotic, antiparkinsonism, and psychotropic drugs, NEC		X61	58	66	<b>124</b>
Intent self-poisoning by and exposure to narcotics and psychodysleptics, NEC		X62	44	25	<b>69</b>
Intent self-poisoning by and exposure to other drugs acting on autonomic nervous system		X63	4	3	<b>7</b>
Intent self-poisoning by other and unspecified drugs, medicaments and biological substances		X64	142	167	<b>309</b>
<b>Total Deaths Related to Illicit Drug Use</b>			<b>266</b>	<b>274</b>	<b>540</b>

<b>DIAGNOSIS</b>	<b>ICD10</b>	<b>Males</b>	<b>Females</b>	<b>Total</b>
<b>2002- (il)Licit related deaths</b>				
Mental and behavioural disorders due to use of opioids, dependence syndrome	F11.2	1	0	<b>1</b>
Mental and behavioural disorders due to use of other stimulants including caffeine, dependence syndrome	F15.2	0	0	<b>0</b>
Mental and behavioural disorders due to use of hallucinogens, dependence syndrome	F16.2	0	0	<b>0</b>
Mental and behavioural disorders due to multiple drug use and use of psychoactive substances, dependence syndrome	F19.2	19	15	<b>34</b>
Mental and behavioural disorders due to use of sedatives or hypnotics, harmful use	F13.1	0	0	<b>0</b>

**CCENDU-Canadian Community Epidemiology Network on Drug Use**  
**Indicator: Mortality - Canada**

Mental and behavioural disorders due to use of opioids, harmful use		F11.1	0	0	<b>0</b>
Mental and behavioural disorders due to use of other stimulants including caffeine, harmful use		F15.1	0	0	<b>0</b>
Mental and behavioural disorders due to multiple drug use and use of psychoactive substances, harmful use		F19.1	15	9	<b>24</b>
Mental disorders and diseases of the nervous system, complicating pregnancy, childbirth, and puerperium	This category includes alcohol related incidences. In other words, it is no longer exclusive to the licit/illicit drug classification.	O99.3 (mom)	0	1	<b>1</b>
Maternal care for (suspected) damage to fetus by drugs		O35.5 (mom)	0	0	<b>0</b>
Accidental poisoning by and exposure to nonopioid analgesics, antipyretics, antirheumatics		X40	15	16	<b>31</b>
Accidental poisoning by and exposure to antiepileptic, sedative-hypnotic, antiparkinsonism, psychotropic drugs, NEC		X41	44	20	<b>64</b>
Accidental poisoning by and exposure to narcotic psychodysleptics, NEC		X42	309	107	<b>416</b>
Accidental poisoning by and exposure to other drugs acting on the autonomic nervous system		X43	1	1	<b>2</b>
Accidental poisoning by other unspecified drugs, medicaments, and biological substances		X44	222	136	<b>358</b>
<b>Total Deaths Related to Illicit &amp; Licit Drug Use</b>			<b>626</b>	<b>305</b>	<b>931</b>

### Mortality Statistics 2002 – Canada (Cont.)

DIAGNOSIS	ICD10	Males	Females	Total
<b>2002- Alcohol related deaths</b>				
Mental and behavioural disorders due to use of alcohol, withdrawal state with delirium	F10.4	16	3	<b>19</b>
Mental and behavioural disorders due to use of alcohol, amnesic syndrome	F10.6	20	8	<b>28</b>
Mental and behavioural disorders due to use of alcohol, residual and late onset psychotic disorder	F10.7	43	16	<b>59</b>
Mental and behavioural disorders due to use of alcohol, psychotic disorder	F10.5	0	0	<b>0</b>
Mental and behavioural disorders due to use of alcohol, acute intoxication	F10.0	51	11	<b>62</b>
Mental and behavioural disorders due to use of alcohol, other mental and behavioural disorders	F10.8	0	0	<b>0</b>
Mental and behavioural disorders due to use of alcohol, unspecified mental and behavioural disorder	F10.9	19	8	<b>27</b>
Mental and behavioural disorders due to use of alcohol, dependence syndrome	F10.2	300	76	<b>376</b>
Mental and behavioural disorders due to use of alcohol, harmful use	F10.1	107	40	<b>147</b>
Alcoholic polyneuropathy	G62.1	3	1	<b>4</b>
Alcoholic cardiomyopathy	I42.6	50	12	<b>62</b>
Alcoholic gastritis	K29.2	3	1	<b>4</b>
Alcoholic fatty liver	K70.0	8	2	<b>10</b>
Alcoholic hepatitis	K70.1	55	33	<b>88</b>

**CCENDU-Canadian Community Epidemiology Network on Drug Use**  
**Indicator: Mortality - Canada**

Alcoholic cirrhosis of liver		K70.3	593	217	<b>810</b>
Alcoholic liver disease unspecified		K70.9	62	26	<b>88</b>
Hepatic fibrosis		K74.0	1	1	<b>2</b>
Other and unspecified cirrhosis of the liver		K74.6	684	394	<b>1078</b>
Liver disease unspecified		K76.9	35	27	<b>62</b>
Acute pancreatitis		K85	158	144	<b>302</b>
Alcohol-induced chronic pancreatitis		K86.0	18	3	<b>21</b>
Other chronic pancreatitis		K86.1	15	6	<b>21</b>
Finding of alcohol in blood		R78.0	0	0	<b>0</b>
Accidental poisoning by and exposure to alcohol		X45	50	18	<b>68</b>
Mental disorders and diseases of the nervous system, complicating pregnancy, childbirth, and puerperium	This category includes licit/illicit drug related incidences. In other words, it is no longer exclusive to the alcohol classification.	O99.3 (mom)	0	1	<b>1</b>
<b>Total Deaths Related to Alcohol</b>			<b>2291</b>	<b>1048</b>	<b>3339</b>