

# 2005 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 7

## Canada Western Red Winter

Percentage of seeded acres / ( ) = last year

Variety	Prairie Average		Manitoba		Saskatchewan		Alberta		British Columbia	
AC Bellatrix	<b>25.6</b>	(9.4)	<b>0.6</b>	(0.0)	<b>19.3</b>	(4.8)	<b>41.7</b>	(28.6)	<b>0.0</b>	(0.0)
CDC Falcon	<b>20.2</b>	(41.2)	<b>70.4</b>	(74.7)	<b>10.0</b>	(11.1)	<b>2.0</b>	(4.3)	<b>0.0</b>	(0.0)
CDC Clair	<b>19.2</b>	(17.0)	<b>4.1</b>	(4.6)	<b>33.2</b>	(40.0)	<b>18.1</b>	(22.4)	<b>0.0</b>	(0.0)
CDC Osprey	<b>9.3</b>	(8.2)	<b>0.0</b>	(0.4)	<b>13.9</b>	(14.1)	<b>11.0</b>	(17.6)	<b>0.0</b>	(0.0)
CDC Raptor	<b>7.5</b>	(6.8)	<b>10.4</b>	(8.0)	<b>17.4</b>	(13.9)	<b>0.0</b>	(0.0)	<b>0.0</b>	(0.0)
AC Readymade	<b>6.6</b>	(4.4)	<b>0.0</b>	(0.0)	<b>0.0</b>	(0.0)	<b>13.9</b>	(14.9)	<b>0.0</b>	(0.0)
AC Tempest	<b>1.9</b>	(2.2)	<b>0.0</b>	(0.0)	<b>0.0</b>	(0.0)	<b>3.9</b>	(7.4)	<b>0.0</b>	(0.0)
Radiant	<b>1.8</b>	(0.0)	<b>0.0</b>	(0.0)	<b>1.3</b>	(0.0)	<b>3.0</b>	(0.0)	<b>0.0</b>	(0.0)
Other	<b>1.6</b>	(0.0)	<b>0.0</b>	(0.0)	<b>0.0</b>	(0.0)	<b>3.3</b>	(0.0)	<b>0.0</b>	(0.0)
CDC Buteo	<b>1.5</b>	(0.4)	<b>3.4</b>	(0.1)	<b>2.5</b>	(1.5)	<b>0.0</b>	(0.0)	<b>0.0</b>	(0.0)
CDC Kestrel	<b>1.5</b>	(2.4)	<b>2.3</b>	(1.4)	<b>1.0</b>	(5.0)	<b>1.4</b>	(2.3)	<b>0.0</b>	(0.0)
Norstar	<b>1.2</b>	(0.5)	<b>1.2</b>	(0.0)	<b>0.3</b>	(0.9)	<b>1.7</b>	(1.0)	<b>0.0</b>	(0.0)
McClintock	<b>1.1</b>	(2.2)	<b>4.3</b>	(2.5)	<b>0.3</b>	(3.3)	<b>0.0</b>	(1.2)	<b>0.0</b>	(0.0)
CDC Harrier	<b>1.0</b>	(5.4)	<b>3.4</b>	(8.3)	<b>0.7</b>	(5.5)	<b>0.0</b>	(0.3)	<b>0.0</b>	(0.0)

\* Trace: less than 0.05 percent