

# 2004 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 1

## Canada Western Red Spring

Percentage of seeded acres / ( ) = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
AC Barrie	<b>25.8</b> (31.8)	<b>43.1</b> (45.8)	<b>28.8</b> (34.8)	<b>11.8</b> (17.0)	<b>0.0</b> (0.0)
Superb	<b>14.4</b> (3.9)	<b>19.0</b> (7.5)	<b>11.4</b> (2.3)	<b>16.2</b> (3.6)	<b>1.0</b> (0.0)
McKenzie	<b>7.3</b> (6.4)	<b>2.0</b> (2.9)	<b>10.3</b> (8.7)	<b>6.2</b> (5.7)	<b>1.1</b> (0.0)
Prodigy	<b>7.0</b> (7.4)	<b>0.8</b> (1.2)	<b>8.9</b> (8.6)	<b>7.8</b> (10.3)	<b>3.0</b> (8.0)
AC Intrepid	<b>6.1</b> (6.6)	<b>3.4</b> (4.8)	<b>5.5</b> (6.0)	<b>8.2</b> (8.9)	<b>32.5</b> (12.3)
CDC Teal	<b>6.0</b> (7.2)	<b>2.3</b> (3.5)	<b>4.3</b> (5.0)	<b>10.5</b> (13.5)	<b>26.3</b> (11.3)
AC Splendor	<b>4.8</b> (5.4)	<b>1.9</b> (1.5)	<b>3.8</b> (4.4)	<b>7.9</b> (9.6)	<b>22.5</b> (28.2)
AC Cadillac	<b>4.1</b> (4.6)	<b>1.4</b> (2.1)	<b>6.7</b> (7.8)	<b>2.0</b> (1.8)	<b>0.0</b> (0.0)
AC Domain	<b>3.5</b> (4.7)	<b>17.1</b> (19.2)	<b>0.5</b> (0.5)	<b>0.1</b> (0.1)	<b>0.0</b> (0.0)
AC Eatonia	<b>3.1</b> (2.4)	<b>T</b> (T)	<b>1.3</b> (1.0)	<b>7.6</b> (6.4)	<b>0.0</b> (0.0)
AC Elsa	<b>2.8</b> (3.3)	<b>0.5</b> (0.5)	<b>4.2</b> (4.7)	<b>2.3</b> (3.2)	<b>0.0</b> (0.0)
AC Abbey	<b>2.1</b> (1.6)	<b>T</b> (0.0)	<b>1.7</b> (1.2)	<b>4.0</b> (3.2)	<b>0.0</b> (0.0)
Columbus	<b>2.1</b> (2.1)	<b>0.1</b> (0.2)	<b>3.0</b> (3.7)	<b>1.8</b> (1.3)	<b>0.0</b> (0.0)
Katepwa	<b>1.7</b> (2.0)	<b>0.2</b> (0.6)	<b>2.5</b> (2.6)	<b>1.6</b> (2.0)	<b>1.5</b> (0.0)
Roblin	<b>1.5</b> (1.6)	<b>0.1</b> (0.3)	<b>0.4</b> (0.2)	<b>4.0</b> (4.3)	<b>8.5</b> (35.1)
CDC Bounty	<b>1.5</b> (0.0)	<b>3.0</b> (0.0)	<b>1.1</b> (0.0)	<b>1.2</b> (0.0)	<b>0.0</b> (0.0)
Other	<b>1.0</b> (2.3)	<b>0.9</b> (4.8)	<b>0.9</b> (1.6)	<b>1.3</b> (1.6)	<b>0.0</b> (0.0)
Laura	<b>1.0</b> (1.1)	<b>0.0</b> (0.1)	<b>1.4</b> (1.3)	<b>0.8</b> (1.7)	<b>0.0</b> (0.0)
CDC Makwa	<b>0.6</b> (0.6)	<b>0.0</b> (0.0)	<b>0.9</b> (1.0)	<b>0.4</b> (0.6)	<b>0.0</b> (0.0)
AC Majestic	<b>0.6</b> (0.8)	<b>1.5</b> (1.6)	<b>0.5</b> (0.5)	<b>0.2</b> (0.8)	<b>0.0</b> (0.0)
Neepawa	<b>0.5</b> (0.5)	<b>T</b> (T)	<b>0.5</b> (0.7)	<b>0.8</b> (0.5)	<b>2.7</b> (0.0)
AC Cora	<b>0.3</b> (0.5)	<b>1.6</b> (2.0)	<b>0.1</b> (T)	<b>T</b> (0.0)	<b>0.0</b> (0.0)
5600 HR	<b>0.3</b> (1.3)	<b>0.3</b> (0.4)	<b>0.4</b> (2.3)	<b>0.3</b> (0.5)	<b>0.0</b> (0.0)
Pasqua	<b>0.3</b> (0.3)	<b>0.0</b> (0.0)	<b>0.5</b> (0.6)	<b>0.2</b> (0.2)	<b>0.0</b> (0.0)
AC Michael	<b>0.3</b> (0.2)	<b>T</b> (0.0)	<b>0.1</b> (T)	<b>0.7</b> (0.6)	<b>0.0</b> (0.0)
Leader	<b>0.2</b> (0.3)	<b>0.0</b> (0.0)	<b>0.2</b> (0.1)	<b>0.5</b> (0.9)	<b>0.0</b> (0.0)
5500 HR	<b>0.2</b> (0.0)	<b>0.2</b> (0.0)	<b>0.1</b> (0.0)	<b>0.3</b> (0.0)	<b>0.0</b> (0.0)
Park	<b>0.2</b> (0.2)	<b>0.0</b> (0.0)	<b>T</b> (0.0)	<b>0.6</b> (0.7)	<b>0.0</b> (0.0)
Conway	<b>0.2</b> (0.2)	<b>0.0</b> (T)	<b>0.1</b> (T)	<b>0.5</b> (0.7)	<b>1.0</b> (5.1)
Alsen	<b>0.1</b> (0.2)	<b>0.4</b> (1.0)	<b>0.1</b> (T)	<b>0.1</b> (T)	<b>0.0</b> (0.0)
Invader	<b>0.1</b> (0.2)	<b>0.0</b> (T)	<b>0.1</b> (0.2)	<b>0.1</b> (0.2)	<b>0.0</b> (0.0)
AC Minto	<b>T</b> (0.1)	<b>0.0</b> (0.0)	<b>0.1</b> (0.2)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)
Alikat	<b>T</b> (T)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.1</b> (T)	<b>0.0</b> (0.0)

\* Trace: less than 0.05 percent

# 2004 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 2

## Canada Western Amber Durum

Percentage of seeded acres / ( ) = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
Kyle	<b>41.2</b> (49.2)	<b>58.4</b> (61.6)	<b>44.0</b> (51.3)	<b>31.2</b> (43.1)	<b>0.0</b> (0.0)
AC Avonlea	<b>39.7</b> (31.0)	<b>22.7</b> (27.1)	<b>38.2</b> (30.4)	<b>45.2</b> (32.9)	<b>0.0</b> (0.0)
AC Navigator	<b>7.4</b> (4.8)	<b>3.0</b> (1.9)	<b>7.6</b> (5.4)	<b>7.0</b> (3.3)	<b>0.0</b> (0.0)
AC Morse	<b>6.2</b> (9.1)	<b>7.5</b> (2.5)	<b>4.7</b> (7.2)	<b>11.1</b> (14.4)	<b>0.0</b> (0.0)
Napoleon	<b>1.6</b> (0.6)	<b>4.2</b> (3.4)	<b>1.9</b> (0.8)	<b>0.4</b> (0.2)	<b>0.0</b> (0.0)
Other	<b>1.4</b> (1.4)	<b>0.0</b> (1.8)	<b>1.3</b> (1.7)	<b>1.9</b> (0.7)	<b>0.0</b> (0.0)
Plenty	<b>0.9</b> (1.0)	<b>0.0</b> (0.0)	<b>0.6</b> (0.8)	<b>1.9</b> (1.4)	<b>0.0</b> (0.0)
Wakooma	<b>0.7</b> (1.3)	<b>0.0</b> (1.9)	<b>0.8</b> (1.2)	<b>0.3</b> (1.7)	<b>0.0</b> (0.0)
Sceptre	<b>0.6</b> (0.9)	<b>2.8</b> (0.0)	<b>0.6</b> (0.8)	<b>0.7</b> (1.3)	<b>0.0</b> (0.0)
AC Melita	<b>0.3</b> (0.4)	<b>0.0</b> (0.0)	<b>0.3</b> (0.1)	<b>0.3</b> (1.0)	<b>0.0</b> (0.0)
Medora	<b>0.1</b> (0.2)	<b>1.4</b> (0.0)	<b>0.1</b> (0.3)	T (0.0)	<b>0.0</b> (0.0)

\* Trace: less than 0.05 percent

# 2004 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 3

## Canada Western Hard White

Percentage of seeded acres / ( ) = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
Snowbird	98.2 (99.0)	98.6 (98.8)	97.2 (98.8)	100.0 (100.0)	0.0 (0.0)
Kanata	1.8 (1.0)	1.4 (1.2)	2.8 (1.2)	0.0 (0.0)	0.0 (0.0)
* Trace: less than 0.05 percent					

# 2004 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 4

## Canada Western Extra Strong

Percentage of seeded acres / ( ) = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
Bluesky	<b>41.5</b> (36.9)	<b>0.0</b> (21.7)	<b>25.5</b> (43.0)	<b>55.0</b> (38.9)	<b>0.0</b> (0.0)
Glenlea	<b>26.9</b> (28.9)	<b>0.0</b> (22.8)	<b>51.3</b> (34.0)	<b>18.5</b> (27.9)	<b>0.0</b> (0.0)
Laser	<b>22.0</b> (21.3)	<b>69.1</b> (18.1)	<b>0.0</b> (5.9)	<b>26.4</b> (33.3)	<b>0.0</b> (0.0)
Amazon	<b>7.0</b> (2.5)	<b>0.0</b> (7.3)	<b>23.2</b> (3.5)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)
Wildcat	<b>2.6</b> (8.8)	<b>30.9</b> (22.6)	<b>0.0</b> (13.7)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)
AC Corinne	<b>0.0</b> (1.4)	<b>0.0</b> (7.5)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)
CDC Rama	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)
Glenavon	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)

\* Trace: less than 0.05 percent

# 2004 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 5

## Canada Prairie Spring White

Percentage of seeded acres / ( ) = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
AC Vista	<b>48.0</b> (54.1)	<b>100.0</b> (72.9)	<b>46.3</b> (52.7)	<b>54.7</b> (59.8)	<b>0.0</b> (0.0)
Genesis	<b>30.4</b> (23.9)	<b>0.0</b> (0.0)	<b>32.1</b> (25.7)	<b>14.1</b> (15.9)	<b>0.0</b> (0.0)
AC Karma	<b>20.0</b> (17.9)	<b>0.0</b> (0.0)	<b>21.6</b> (18.6)	<b>0.0</b> (24.3)	<b>0.0</b> (0.0)
AC 2000	<b>1.5</b> (4.1)	<b>0.0</b> (27.1)	<b>0.0</b> (3.0)	<b>31.3</b> (0.0)	<b>0.0</b> (0.0)

\* Trace: less than 0.05 percent

# 2004 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 6

## Canada Prairie Spring Red

Percentage of seeded acres / ( ) = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
AC Crystal	<b>51.8</b> (64.1)	<b>26.1</b> (24.3)	<b>67.4</b> (85.2)	<b>46.4</b> (59.7)	<b>69.7</b> (27.0)
5700PR	<b>26.0</b> (5.3)	<b>23.5</b> (4.9)	<b>24.4</b> (4.3)	<b>26.6</b> (4.5)	<b>30.3</b> (73.0)
AC Taber	<b>10.2</b> (13.4)	<b>40.1</b> (61.4)	<b>4.3</b> (7.6)	<b>11.8</b> (14.0)	<b>0.0</b> (0.0)
AC Foremost	<b>9.7</b> (15.2)	<b>0.0</b> (0.0)	<b>3.8</b> (2.9)	<b>12.1</b> (19.4)	<b>0.0</b> (0.0)
Oslo	<b>2.2</b> (1.7)	<b>10.3</b> (9.4)	<b>0.0</b> (0.0)	<b>2.9</b> (2.0)	<b>0.0</b> (0.0)
Cutler	<b>0.2</b> (0.3)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.2</b> (0.4)	<b>0.0</b> (0.0)

\* Trace: less than 0.05 percent

# 2004 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 7

## Canada Western Red Winter

Percentage of seeded acres / ( ) = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
CDC Falcon	<b>41.2</b> (32.8)	<b>75.2</b> (64.5)	<b>11.1</b> (9.7)	<b>4.3</b> (6.4)	<b>0.0</b> (0.0)
CDC Clair	<b>16.8</b> (26.6)	<b>4.0</b> (12.1)	<b>40.1</b> (55.4)	<b>22.5</b> (24.1)	<b>0.0</b> (0.0)
AC Bellatrix	<b>9.5</b> (4.8)	<b>0.0</b> (0.0)	<b>4.8</b> (1.8)	<b>28.7</b> (14.0)	<b>0.0</b> (0.0)
CDC Osprey	<b>8.2</b> (10.3)	<b>0.4</b> (0.3)	<b>14.1</b> (9.2)	<b>17.4</b> (25.3)	<b>0.0</b> (0.0)
CDC Raptor	<b>6.8</b> (4.8)	<b>8.1</b> (5.6)	<b>13.7</b> (8.8)	<b>0.0</b> (0.5)	<b>0.0</b> (0.0)
CDC Harrier	<b>5.3</b> (7.9)	<b>8.2</b> (14.6)	<b>5.5</b> (5.7)	<b>0.3</b> (0.0)	<b>0.0</b> (0.0)
AC Readymade	<b>4.4</b> (3.5)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>14.9</b> (11.2)	<b>0.0</b> (0.0)
CDC Kestrel	<b>2.4</b> (6.8)	<b>1.4</b> (2.3)	<b>5.0</b> (8.4)	<b>2.3</b> (12.0)	<b>0.0</b> (0.0)
McClintock	<b>2.3</b> (0.3)	<b>2.5</b> (0.5)	<b>3.3</b> (0.3)	<b>1.2</b> (0.0)	<b>0.0</b> (0.0)
AC Tempest	<b>2.2</b> (1.1)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>7.4</b> (3.6)	<b>0.0</b> (0.0)
Norstar	<b>0.5</b> (1.2)	<b>0.0</b> (0.3)	<b>0.9</b> (0.7)	<b>1.0</b> (2.9)	<b>0.0</b> (0.0)
CDC Buteo	<b>0.4</b> (T)	<b>0.1</b> (T)	<b>1.5</b> (T)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)

\* Trace: less than 0.05 percent

# 2004 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 8

## Canada Western Soft White Spring

Percentage of seeded acres / ( ) = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
AC Andrew	<b>85.6</b> (39.5)	<b>62.8</b> (0.0)	<b>97.9</b> (100.0)	<b>83.1</b> (39.3)	<b>0.0</b> (0.0)
AC Reed	<b>5.2</b> (36.2)	<b>14.8</b> (20.1)	<b>0.0</b> (0.0)	<b>6.2</b> (37.6)	<b>0.0</b> (0.0)
AC Meena	<b>4.4</b> (1.6)	<b>0.0</b> (0.0)	<b>2.1</b> (0.0)	<b>5.2</b> (1.7)	<b>0.0</b> (0.0)
AC Nanda	<b>3.7</b> (15.1)	<b>22.3</b> (79.9)	<b>0.0</b> (0.0)	<b>4.0</b> (13.3)	<b>0.0</b> (0.0)
AC Phil	<b>1.2</b> (7.7)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>1.5</b> (8.2)	<b>0.0</b> (0.0)
* Trace: less than 0.05 percent					

# 2004 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 9

## Malting -- Two-Row

Percentage of seeded acres / ( ) = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
AC Metcalfe	<b>47.8</b> (39.6)	<b>61.5</b> (49.3)	<b>44.9</b> (39.0)	<b>48.6</b> (38.2)	<b>80.1</b> (89.0)
CDC Kendall	<b>14.5</b> (13.2)	<b>3.2</b> (2.3)	<b>19.6</b> (17.9)	<b>9.9</b> (9.4)	<b>0.0</b> (0.0)
Harrington	<b>14.4</b> (22.1)	<b>1.6</b> (3.7)	<b>11.3</b> (19.1)	<b>20.9</b> (29.8)	<b>19.9</b> (11.0)
CDC Copeland	<b>8.4</b> (1.4)	<b>3.3</b> (0.5)	<b>10.9</b> (1.7)	<b>6.1</b> (1.3)	<b>0.0</b> (0.0)
Merit	<b>4.7</b> (5.7)	<b>1.8</b> (0.9)	<b>4.9</b> (7.1)	<b>5.0</b> (5.0)	<b>0.0</b> (0.0)
CDC Stratus	<b>4.4</b> (9.1)	<b>16.7</b> (29.8)	<b>4.3</b> (8.7)	<b>2.3</b> (5.3)	<b>0.0</b> (0.0)
Stein	<b>2.7</b> (5.0)	<b>0.0</b> (0.0)	<b>2.6</b> (4.1)	<b>3.3</b> (7.2)	<b>0.0</b> (0.0)
Other	<b>1.6</b> (3.1)	<b>4.5</b> (12.0)	<b>0.7</b> (2.0)	<b>2.4</b> (2.8)	<b>0.0</b> (0.0)
Newdale	<b>0.7</b> (0.0)	<b>6.8</b> (0.0)	<b>0.4</b> (0.0)	<b>0.1</b> (0.0)	<b>0.0</b> (0.0)
B1202	<b>0.4</b> (0.6)	<b>0.0</b> (1.4)	<b>0.1</b> (0.4)	<b>0.9</b> (0.8)	<b>0.0</b> (0.0)
CDC Select	<b>0.2</b> (0.0)	<b>0.6</b> (0.0)	<b>0.2</b> (0.0)	<b>0.2</b> (0.0)	<b>0.0</b> (0.0)
AC Bountiful	<b>0.2</b> (0.1)	<b>0.0</b> (0.2)	<b>0.1</b> (0.1)	<b>0.2</b> (0.1)	<b>0.0</b> (0.0)
Calder	<b>T</b> (0.0)	<b>0.0</b> (0.0)	<b>0.1</b> (0.0)	<b>T</b> (0.0)	<b>0.0</b> (0.0)

\* Trace: less than 0.05 percent

# 2004 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 10

## Malting -- Six-Row

Percentage of seeded acres / ( ) = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
Robust	<b>35.1</b> (37.0)	<b>59.8</b> (68.0)	<b>16.8</b> (18.8)	<b>7.1</b> (1.9)	<b>0.0</b> (0.0)
Excel	<b>30.9</b> (31.8)	<b>21.5</b> (20.1)	<b>42.3</b> (40.6)	<b>14.5</b> (35.8)	<b>0.0</b> (0.0)
Legacy	<b>15.4</b> (6.9)	<b>9.3</b> (1.7)	<b>22.5</b> (11.6)	<b>5.6</b> (2.8)	<b>0.0</b> (0.0)
B1602	<b>6.6</b> (13.8)	<b>1.5</b> (4.5)	<b>6.2</b> (16.4)	<b>35.4</b> (38.1)	<b>72.1</b> (100.0)
Lacey	<b>4.7</b> (0.0)	<b>5.6</b> (0.0)	<b>4.6</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)
Other	<b>3.1</b> (4.5)	<b>2.0</b> (4.4)	<b>1.3</b> (3.5)	<b>20.8</b> (11.1)	<b>27.9</b> (0.0)
CDC Battleford	<b>2.4</b> (0.7)	<b>0.1</b> (0.0)	<b>4.1</b> (0.5)	<b>6.2</b> (5.6)	<b>0.0</b> (0.0)
CDC Sisler	<b>1.1</b> (4.9)	<b>0.2</b> (1.2)	<b>1.8</b> (8.1)	<b>2.9</b> (3.7)	<b>0.0</b> (0.0)
CDC Yorkton	<b>0.5</b> (0.3)	<b>0.0</b> (0.0)	<b>0.1</b> (0.4)	<b>7.0</b> (1.0)	<b>0.0</b> (0.0)
CDC Tisdale	<b>0.1</b> (0.0)	<b>0.0</b> (0.0)	<b>0.2</b> (0.0)	<b>0.4</b> (0.0)	<b>0.0</b> (0.0)
Tradition	<b>0.0</b> (0.0)				
CDC Springside	<b>0.0</b> (0.0)				

\* Trace: less than 0.05 percent

# 2004 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 11

## Two-Row and Six-Row Malting Barley

Percentage of seeded acres / ( ) = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
Malting -- Two-Ro	87.3 (81.6)	52.7 (47.3)	88.3 (81.7)	97.8 (95.6)	78.9 (72.8)
Malting -- Six-Ro	12.7 (18.4)	47.3 (52.7)	11.7 (18.3)	2.2 (4.4)	21.1 (27.2)
* Trace: less than 0.05 percent					

# 2004 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 12

## Hulless -- Two-Row

Percentage of seeded acres / ( ) = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
Other	<b>61.6</b> (20.9)	<b>13.3</b> (14.1)	<b>79.2</b> (12.9)	<b>58.2</b> (36.6)	<b>0.0</b> (0.0)
CDC McGwire	<b>27.1</b> (43.6)	<b>82.7</b> (68.0)	<b>9.0</b> (59.0)	<b>17.3</b> (4.1)	<b>0.0</b> (0.0)
CDC Dawn	<b>3.4</b> (7.7)	<b>0.0</b> (0.0)	<b>5.1</b> (10.5)	<b>0.0</b> (10.4)	<b>0.0</b> (0.0)
Phoenix	<b>2.5</b> (3.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>24.5</b> (9.3)	<b>0.0</b> (0.0)
CDC Richard	<b>1.3</b> (12.2)	<b>0.0</b> (0.0)	<b>2.0</b> (2.4)	<b>0.0</b> (34.7)	<b>0.0</b> (0.0)
CDC Gainer	<b>1.2</b> (8.7)	<b>0.0</b> (9.2)	<b>1.8</b> (15.2)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)
Condor	<b>1.1</b> (0.0)	<b>0.0</b> (0.0)	<b>1.6</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)
CDC Freedom	<b>0.9</b> (3.9)	<b>4.0</b> (8.7)	<b>0.0</b> (0.0)	<b>0.0</b> (4.9)	<b>0.0</b> (0.0)
CDC Alamo	<b>0.9</b> (0.0)	<b>0.0</b> (0.0)	<b>1.3</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)
HB803	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)
HB805	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)
Merlin	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)
Tercel	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)
CDC Fibar	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)
CDC Candle	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)
CDC Rattan	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)
CDC Speedy	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)

\* Trace: less than 0.05 percent

# 2004 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 13

## Hulless -- Six-Row

Percentage of seeded acres / ( ) = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
Other	<b>80.5</b> (32.8)	<b>0.0</b> (56.8)	<b>87.6</b> (40.0)	<b>77.1</b> (26.4)	<b>0.0</b> (0.0)
AC Bacon	<b>11.5</b> (34.2)	<b>0.0</b> (0.0)	<b>12.4</b> (0.0)	<b>11.2</b> (52.0)	<b>0.0</b> (0.0)
Falcon	<b>8.0</b> (11.7)	<b>100.0</b> (43.2)	<b>0.0</b> (0.0)	<b>11.6</b> (11.1)	<b>0.0</b> (0.0)
CDC Silky	<b>0.0</b> (14.4)	<b>0.0</b> (0.0)	<b>0.0</b> (60.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)
Peregrine	<b>0.0</b> (6.9)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (10.4)	<b>0.0</b> (0.0)
Jaeger	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)
CDC Buck	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)
AC Hawkeye	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)

\* Trace: less than 0.05 percent

# 2004 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 14

## Feed Barley -- Two-Row

Percentage of seeded acres / ( ) = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
Xena	<b>37.2</b> (20.7)	<b>3.7</b> (2.8)	<b>45.3</b> (22.7)	<b>39.9</b> (21.9)	<b>66.1</b> (0.0)
CDC Dolly	<b>30.7</b> (44.9)	<b>5.4</b> (26.7)	<b>43.1</b> (64.6)	<b>31.0</b> (39.8)	<b>21.0</b> (70.0)
Other	<b>11.7</b> (20.5)	<b>3.0</b> (55.2)	<b>7.7</b> (9.1)	<b>14.2</b> (20.9)	<b>12.9</b> (23.1)
CONLON	<b>9.3</b> (0.0)	<b>84.5</b> (0.0)	<b>0.6</b> (0.0)	<b>T</b> (0.0)	<b>0.0</b> (0.0)
CDC Bold	<b>5.7</b> (8.0)	<b>2.7</b> (11.8)	<b>2.5</b> (2.7)	<b>7.1</b> (9.5)	<b>0.0</b> (6.9)
Seebe	<b>5.4</b> (5.1)	<b>0.7</b> (1.4)	<b>0.7</b> (0.0)	<b>7.6</b> (7.2)	<b>0.0</b> (0.0)
CDC Helgason	<b>0.1</b> (0.4)	<b>0.0</b> (2.1)	<b>0.1</b> (0.6)	<b>0.1</b> (0.2)	<b>0.0</b> (0.0)
Abee	<b>T</b> (0.3)	<b>0.0</b> (0.0)	<b>0.0</b> (0.1)	<b>T</b> (0.4)	<b>0.0</b> (0.0)
CDC Fleet	<b>T</b> (0.2)	<b>0.0</b> (0.0)	<b>T</b> (0.3)	<b>0.0</b> (0.1)	<b>0.0</b> (0.0)

\* Trace: less than 0.05 percent

# 2004 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 15

## Feed Barley -- Six-Row

Percentage of seeded acres / ( ) = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
Other	<b>32.2</b> (22.3)	<b>37.3</b> (28.0)	<b>25.9</b> (20.7)	<b>37.1</b> (22.1)	<b>5.1</b> (5.2)
AC Lacombe	<b>19.8</b> (28.4)	<b>6.5</b> (18.7)	<b>10.3</b> (14.0)	<b>30.3</b> (38.1)	<b>20.0</b> (15.0)
Brier	<b>11.1</b> (9.5)	<b>0.0</b> (0.0)	<b>27.0</b> (23.0)	<b>4.3</b> (5.5)	<b>2.6</b> (0.0)
AC Rosser	<b>10.6</b> (11.6)	<b>15.0</b> (14.6)	<b>24.2</b> (27.0)	<b>0.8</b> (3.6)	<b>0.0</b> (0.0)
AC Harper	<b>7.4</b> (6.1)	<b>8.1</b> (0.7)	<b>0.5</b> (0.1)	<b>12.4</b> (10.5)	<b>0.0</b> (0.0)
Bedford	<b>4.0</b> (3.8)	<b>26.5</b> (26.8)	<b>0.2</b> (0.1)	<b>0.3</b> (T)	<b>0.0</b> (0.0)
Virden	<b>3.6</b> (3.0)	<b>3.3</b> (1.5)	<b>7.5</b> (9.0)	<b>1.2</b> (0.4)	<b>0.0</b> (0.0)
Vivar	<b>2.8</b> (1.6)	<b>2.5</b> (4.6)	<b>0.0</b> (0.0)	<b>5.0</b> (1.8)	<b>0.0</b> (0.0)
AC Albright	<b>2.2</b> (2.5)	<b>0.0</b> (1.4)	<b>0.0</b> (0.1)	<b>1.5</b> (2.9)	<b>37.9</b> (48.0)
Kasota	<b>1.7</b> (2.2)	<b>T</b> (1.1)	<b>0.5</b> (0.0)	<b>1.2</b> (2.9)	<b>24.7</b> (31.8)
Trochu	<b>1.3</b> (1.9)	<b>0.0</b> (0.0)	<b>1.1</b> (1.0)	<b>2.0</b> (2.8)	<b>0.0</b> (0.0)
Bronco	<b>1.2</b> (1.1)	<b>0.8</b> (1.3)	<b>1.9</b> (2.7)	<b>0.9</b> (0.3)	<b>0.0</b> (0.0)
Leduc	<b>1.0</b> (2.2)	<b>0.0</b> (0.0)	<b>1.0</b> (2.4)	<b>1.3</b> (2.7)	<b>0.0</b> (0.0)
Mahigan	<b>0.6</b> (1.3)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.6</b> (2.4)	<b>9.7</b> (0.0)
CDC Earl	<b>0.4</b> (1.8)	<b>0.0</b> (1.1)	<b>0.0</b> (0.0)	<b>0.8</b> (2.9)	<b>0.0</b> (0.0)
Stetson	<b>0.2</b> (0.7)	<b>0.0</b> (0.3)	<b>0.0</b> (0.0)	<b>0.4</b> (1.2)	<b>0.0</b> (0.0)

\* Trace: less than 0.05 percent

# 2004 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 16

## All Barley (Malting, Feed and Hulless)

Percentage of seeded acres / ( ) = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
Malting -- Two-Ro	<b>56.3</b> (54.5)	<b>34.2</b> (35.8)	<b>70.8</b> (65.7)	<b>48.7</b> (49.1)	<b>42.9</b> (38.1)
Feed -- (2 & 6 Ro	<b>35.0</b> (32.7)	<b>34.3</b> (23.5)	<b>19.0</b> (19.2)	<b>50.0</b> (48.2)	<b>45.6</b> (47.7)
Malting -- Six-Ro	<b>8.2</b> (12.3)	<b>30.8</b> (39.9)	<b>9.4</b> (14.7)	<b>1.1</b> (2.3)	<b>11.5</b> (14.2)
Hulless -- (2 & 6 R	<b>0.5</b> (0.4)	<b>0.8</b> (0.7)	<b>0.8</b> (0.4)	<b>0.2</b> (0.4)	<b>0.0</b> (0.0)

\* Trace: less than 0.05 percent