

2005 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 1

Canada Western Red Spring

Percentage of seeded acres / () = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
AC Barrie	21.4 (26.0)	41.0 (43.2)	24.0 (28.9)	8.5 (12.1)	0.0 (0.0)
Superb	18.9 (14.3)	16.7 (18.7)	14.7 (11.3)	25.2 (16.1)	10.1 (1.0)
McKenzie	8.3 (7.3)	2.2 (2.0)	13.8 (10.2)	4.6 (6.2)	6.1 (1.1)
Prodigy	6.6 (7.0)	0.1 (0.8)	8.3 (8.9)	7.8 (7.9)	3.0 (3.0)
AC Intrepid	6.2 (6.2)	3.5 (3.5)	5.8 (5.7)	7.9 (8.4)	26.9 (32.5)
CDC Teal	4.8 (6.0)	2.7 (2.4)	3.6 (4.2)	7.3 (10.4)	16.5 (26.3)
AC Eatonia	4.5 (3.1)	0.0 (T)	1.8 (1.3)	10.3 (7.5)	0.0 (0.0)
AC Domain	4.5 (3.6)	21.5 (17.3)	1.1 (0.5)	0.2 (0.1)	0.0 (0.0)
AC Splendor	4.5 (4.8)	1.3 (1.9)	4.0 (3.8)	6.6 (7.9)	27.6 (22.5)
AC Cadillac	3.2 (4.2)	1.6 (1.4)	5.4 (6.8)	1.5 (2.0)	0.0 (0.0)
AC Elsa	2.6 (2.8)	0.8 (0.5)	3.8 (4.2)	2.0 (2.2)	0.0 (0.0)
Other	2.3 (1.0)	1.2 (0.9)	2.2 (0.9)	3.1 (1.3)	2.4 (0.0)
AC Abbey	2.0 (2.1)	0.0 (T)	2.1 (1.7)	2.8 (3.9)	0.0 (0.0)
Columbus	1.7 (2.0)	0.1 (0.1)	2.8 (2.9)	1.2 (1.8)	0.0 (0.0)
Katepwa	1.5 (1.7)	0.1 (0.2)	2.0 (2.4)	1.4 (1.5)	0.0 (1.5)
Harvest	1.4 (0.0)	2.3 (0.0)	0.2 (0.0)	2.5 (0.0)	0.0 (0.0)
CDC Imagine	1.3 (0.0)	0.3 (0.0)	1.6 (0.0)	1.4 (0.0)	0.0 (0.0)
Roblin	1.3 (1.5)	T (0.1)	0.2 (0.4)	3.1 (3.9)	7.3 (8.5)
CDC Bounty	1.2 (1.5)	1.6 (2.9)	1.2 (1.1)	1.0 (1.2)	0.0 (0.0)
Laura	0.6 (0.9)	0.0 (0.0)	1.0 (1.4)	0.5 (0.8)	0.0 (0.0)
5600 HR	0.2 (0.3)	0.2 (0.3)	0.1 (0.4)	0.4 (0.3)	0.0 (0.0)
AC Cora	0.2 (0.3)	0.9 (1.6)	0.1 (0.1)	0.0 (T)	0.0 (0.0)
5601 HR	0.2 (0.0)	1.0 (0.0)	T (0.0)	T (0.0)	0.0 (0.0)
5500 HR	0.1 (0.2)	0.2 (0.2)	T (0.1)	0.2 (0.3)	0.0 (0.0)
Journey	0.1 (0.0)	0.1 (0.0)	0.1 (0.0)	0.2 (0.0)	0.0 (0.0)
Alsen	0.1 (0.1)	0.6 (0.4)	T (0.1)	T (0.1)	0.0 (0.0)
Lillian	0.1 (0.0)	0.0 (0.0)	0.1 (0.0)	0.2 (0.0)	0.0 (0.0)
Lovitt	0.1 (0.0)	T (0.0)	0.1 (0.0)	T (0.0)	0.0 (0.0)
CDC Go	T (0.0)	T (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)
CDC Osler	T (0.0)	0.0 (0.0)	T (0.0)	0.0 (0.0)	0.0 (0.0)
CDC Makwa	0.0 (0.6)	0.0 (0.0)	0.0 (1.0)	0.0 (0.4)	0.0 (0.0)
AC Majestic	0.0 (0.6)	0.0 (1.4)	0.0 (0.4)	0.0 (0.2)	0.0 (0.0)
Neepawa	0.0 (0.5)	0.0 (T)	0.0 (0.5)	0.0 (0.8)	0.0 (2.7)
Pasqua	0.0 (0.3)	0.0 (0.0)	0.0 (0.5)	0.0 (0.2)	0.0 (0.0)
AC Michael	0.0 (0.3)	0.0 (T)	0.0 (0.1)	0.0 (0.7)	0.0 (0.0)
Leader	0.0 (0.2)	0.0 (0.0)	0.0 (0.2)	0.0 (0.5)	0.0 (0.0)
Park	0.0 (0.2)	0.0 (0.0)	0.0 (T)	0.0 (0.6)	0.0 (0.0)
Conway	0.0 (0.2)	0.0 (0.0)	0.0 (0.1)	0.0 (0.5)	0.0 (1.0)
Invader	0.0 (0.1)	0.0 (0.0)	0.0 (0.1)	0.0 (0.1)	0.0 (0.0)

2005 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 1

Canada Western Red Spring

Percentage of seeded acres / () = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
AC Minto	0.0 (T)	0.0 (0.0)	0.0 (0.1)	0.0 (0.0)	0.0 (0.0)
Alikat	0.0 (T)	0.0 (0.0)	0.0 (0.0)	0.0 (0.1)	0.0 (0.0)
Peace	0.0 (0.0)				
* Trace: less than 0.05 percent					

2005 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 2

Canada Western Amber Durum

Percentage of seeded acres / () = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
AC Avonlea	45.9 (39.6)	11.5 (22.7)	45.0 (38.0)	49.2 (45.1)	0.0 (0.0)
Kyle	34.2 (41.2)	52.5 (58.4)	37.3 (44.2)	23.4 (30.8)	0.0 (0.0)
AC Navigator	9.2 (7.5)	0.0 (3.0)	9.6 (7.6)	8.0 (7.0)	0.0 (0.0)
AC Morse	5.7 (6.2)	12.4 (7.5)	3.2 (4.7)	14.0 (11.4)	0.0 (0.0)
Other	1.7 (1.4)	0.0 (0.0)	1.6 (1.2)	1.8 (1.9)	0.0 (0.0)
Napoleon	1.5 (1.6)	23.6 (4.2)	1.7 (1.9)	0.6 (0.4)	0.0 (0.0)
Plenty	0.7 (0.9)	0.0 (0.0)	0.4 (0.6)	1.8 (2.0)	0.0 (0.0)
Wakooma	0.5 (0.7)	0.0 (0.0)	0.5 (0.8)	0.3 (0.3)	0.0 (0.0)
Sceptre	0.2 (0.6)	0.0 (2.8)	0.2 (0.6)	0.4 (0.7)	0.0 (0.0)
AC Melita	0.2 (0.2)	0.0 (0.0)	0.1 (0.2)	0.4 (0.3)	0.0 (0.0)
Medora	0.2 (0.1)	0.0 (1.4)	0.2 (0.1)	0.0 (T)	0.0 (0.0)

* Trace: less than 0.05 percent

2005 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 3

Canada Western Hard White

Percentage of seeded acres / () = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
Snowbird	97.5 (98.3)	97.5 (98.4)	96.3 (97.5)	99.5 (100.0)	100.0 (0.0)
Kanata	2.5 (1.7)	2.5 (1.6)	3.7 (2.5)	0.5 (0.0)	0.0 (0.0)
* Trace: less than 0.05 percent					

2005 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 4

Canada Western Extra Strong

Percentage of seeded acres / () = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
Glenlea	38.4 (26.9)	0.0 (0.0)	51.3 (51.3)	13.4 (18.5)	0.0 (0.0)
Laser	20.4 (22.0)	0.0 (69.1)	0.0 (0.0)	59.7 (26.4)	0.0 (0.0)
CDC Rama	14.2 (0.0)	0.0 (0.0)	21.6 (0.0)	0.0 (0.0)	0.0 (0.0)
AC Corinne	9.9 (0.0)	0.0 (0.0)	15.0 (0.0)	0.0 (0.0)	0.0 (0.0)
Amazon	8.0 (7.0)	0.0 (0.0)	12.2 (23.2)	0.0 (0.0)	0.0 (0.0)
Bluesky	6.3 (41.5)	0.0 (0.0)	0.0 (25.5)	18.5 (55.0)	0.0 (0.0)
Wildcat	2.9 (2.6)	0.0 (30.9)	0.0 (0.0)	8.4 (0.0)	0.0 (0.0)
Walrus	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)
Burnside	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)
Glenavon	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)

* Trace: less than 0.05 percent

2005 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 5

Canada Prairie Spring White

Percentage of seeded acres / () = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
AC Vista	57.8 (49.1)	100.0 (100.0)	51.1 (47.4)	100.0 (54.7)	0.0 (0.0)
AC Karma	42.2 (19.5)	0.0 (0.0)	48.9 (21.1)	0.0 (0.0)	0.0 (0.0)
Genesis	0.0 (29.9)	0.0 (0.0)	0.0 (31.5)	0.0 (14.1)	0.0 (0.0)
AC 2000	0.0 (1.5)	0.0 (0.0)	0.0 (0.0)	0.0 (31.3)	0.0 (0.0)
Other	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)
* Trace: less than 0.05 percent					

2005 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 6

Canada Prairie Spring Red

Percentage of seeded acres / () = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
AC Crystal	34.4 (51.7)	0.0 (25.7)	43.9 (68.1)	33.1 (46.0)	0.0 (69.7)
5700PR	34.3 (26.3)	52.3 (24.8)	39.7 (24.0)	32.4 (27.2)	0.0 (30.3)
AC Foremost	12.6 (9.7)	0.0 (0.0)	3.8 (3.8)	15.2 (12.1)	0.0 (0.0)
5701PR	11.7 (0.0)	42.0 (0.0)	8.7 (0.0)	11.5 (0.0)	0.0 (0.0)
AC Taber	6.2 (10.0)	1.3 (39.4)	4.0 (4.2)	6.9 (11.6)	0.0 (0.0)
Oslo	0.8 (2.2)	4.4 (10.2)	0.0 (0.0)	0.9 (2.8)	0.0 (0.0)
Cutler	0.0 (0.2)	0.0 (0.0)	0.0 (0.0)	0.0 (0.2)	0.0 (0.0)
Other	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)

* Trace: less than 0.05 percent

2005 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 7

Canada Western Red Winter

Percentage of seeded acres / () = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
AC Bellatrix	25.6 (9.4)	0.6 (0.0)	19.3 (4.8)	41.7 (28.6)	0.0 (0.0)
CDC Falcon	20.2 (41.2)	70.4 (74.7)	10.0 (11.1)	2.0 (4.3)	0.0 (0.0)
CDC Clair	19.2 (17.0)	4.1 (4.6)	33.2 (40.0)	18.1 (22.4)	0.0 (0.0)
CDC Osprey	9.3 (8.2)	0.0 (0.4)	13.9 (14.1)	11.0 (17.6)	0.0 (0.0)
CDC Raptor	7.5 (6.8)	10.4 (8.0)	17.4 (13.9)	0.0 (0.0)	0.0 (0.0)
AC Readymade	6.6 (4.4)	0.0 (0.0)	0.0 (0.0)	13.9 (14.9)	0.0 (0.0)
AC Tempest	1.9 (2.2)	0.0 (0.0)	0.0 (0.0)	3.9 (7.4)	0.0 (0.0)
Radiant	1.8 (0.0)	0.0 (0.0)	1.3 (0.0)	3.0 (0.0)	0.0 (0.0)
Other	1.6 (0.0)	0.0 (0.0)	0.0 (0.0)	3.3 (0.0)	0.0 (0.0)
CDC Buteo	1.5 (0.4)	3.4 (0.1)	2.5 (1.5)	0.0 (0.0)	0.0 (0.0)
CDC Kestrel	1.5 (2.4)	2.3 (1.4)	1.0 (5.0)	1.4 (2.3)	0.0 (0.0)
Norstar	1.2 (0.5)	1.2 (0.0)	0.3 (0.9)	1.7 (1.0)	0.0 (0.0)
McClintock	1.1 (2.2)	4.3 (2.5)	0.3 (3.3)	0.0 (1.2)	0.0 (0.0)
CDC Harrier	1.0 (5.4)	3.4 (8.3)	0.7 (5.5)	0.0 (0.3)	0.0 (0.0)

* Trace: less than 0.05 percent

2005 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 8

Canada Western Soft White Spring

Percentage of seeded acres / () = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
AC Andrew	69.1 (85.4)	0.0 (62.8)	100.0 (98.3)	63.8 (82.2)	0.0 (0.0)
AC Meena	14.6 (5.5)	0.0 (0.0)	0.0 (1.7)	17.2 (6.8)	0.0 (0.0)
Bhishaj	9.7 (0.0)	0.0 (0.0)	0.0 (0.0)	11.4 (0.0)	0.0 (0.0)
AC Nanda	3.8 (3.3)	0.0 (22.3)	0.0 (0.0)	4.5 (3.7)	0.0 (0.0)
AC Reed	2.7 (4.7)	0.0 (14.8)	0.0 (0.0)	3.2 (5.8)	0.0 (0.0)
AC Phil	0.0 (1.1)	0.0 (0.0)	0.0 (0.0)	0.0 (1.4)	0.0 (0.0)

* Trace: less than 0.05 percent

2005 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 9

Malting -- Two-Row

Percentage of seeded acres / () = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
AC Metcalfe	55.1 (47.6)	58.5 (61.6)	54.2 (45.1)	55.2 (48.0)	98.0 (79.2)
CDC Kendall	15.2 (14.4)	3.3 (3.2)	20.3 (19.6)	9.1 (9.7)	0.0 (0.0)
CDC Copeland	12.4 (8.5)	14.8 (3.4)	14.2 (10.8)	9.0 (6.4)	0.0 (0.0)
Harrington	8.5 (14.3)	0.0 (1.5)	5.3 (11.3)	15.8 (20.8)	2.0 (20.8)
Merit	1.9 (4.7)	0.0 (1.7)	1.9 (4.8)	2.4 (5.2)	0.0 (0.0)
Stein	1.8 (2.6)	0.0 (0.0)	1.5 (2.6)	2.6 (3.2)	0.0 (0.0)
Other	1.6 (1.6)	0.7 (4.6)	0.8 (0.7)	3.2 (2.3)	0.0 (0.0)
CDC Stratus	1.6 (4.7)	9.4 (16.9)	1.0 (4.3)	0.9 (3.1)	0.0 (0.0)
Newdale	1.4 (0.7)	13.3 (6.5)	0.6 (0.4)	0.4 (0.1)	0.0 (0.0)
B1202	0.4 (0.4)	0.0 (0.0)	0.1 (0.1)	1.0 (0.9)	0.0 (0.0)
AC Bountiful	0.2 (0.2)	0.0 (0.0)	0.1 (0.1)	0.5 (0.2)	0.0 (0.0)
CDC Select	T (0.2)	0.0 (0.5)	T (0.2)	0.0 (0.2)	0.0 (0.0)
Calder	0.0 (T)	0.0 (0.0)	0.0 (0.1)	0.0 (T)	0.0 (0.0)

* Trace: less than 0.05 percent

2005 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 10

Malting -- Six-Row

Percentage of seeded acres / () = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
Excel	31.0 (31.5)	24.1 (21.1)	36.8 (43.2)	25.0 (14.5)	0.0 (0.0)
Legacy	30.5 (15.9)	16.7 (9.5)	42.1 (23.1)	16.4 (5.6)	0.0 (0.0)
Robust	19.1 (34.3)	41.9 (60.2)	4.7 (15.6)	1.4 (7.1)	0.0 (0.0)
Lacey	8.4 (4.8)	11.1 (5.5)	7.4 (4.9)	0.0 (0.0)	0.0 (0.0)
Tradition	2.6 (0.0)	2.4 (0.0)	3.1 (0.0)	0.0 (0.0)	0.0 (0.0)
CDC Battleford	2.0 (2.3)	0.1 (0.1)	0.8 (3.8)	23.2 (6.2)	0.0 (0.0)
B1602	2.0 (6.2)	0.0 (1.5)	1.6 (5.7)	16.4 (35.4)	0.0 (72.1)
Other	1.8 (3.0)	2.9 (2.0)	1.1 (1.4)	0.7 (20.8)	0.0 (27.9)
CDC Yorkton	1.3 (0.5)	0.4 (0.0)	0.7 (0.1)	12.4 (7.0)	0.0 (0.0)
CDC Sisler	1.2 (1.3)	0.3 (0.2)	1.5 (2.0)	3.9 (2.9)	0.0 (0.0)
CDC Springside	0.2 (0.0)	0.0 (0.0)	0.3 (0.0)	0.6 (0.0)	0.0 (0.0)
CDC Tisdale	0.0 (0.1)	0.0 (0.0)	0.0 (0.2)	0.0 (0.4)	0.0 (0.0)

* Trace: less than 0.05 percent

2005 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 11

Two-Row and Six-Row Malting Barley

Percentage of seeded acres / () = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
Malting -- Two-Row	89.0 (87.0)	58.2 (53.1)	89.7 (87.7)	97.7 (97.8)	100.0 (79.1)
Malting -- Six-Row	11.0 (13.0)	41.8 (46.9)	10.3 (12.3)	2.3 (2.2)	0.0 (20.9)

* Trace: less than 0.05 percent

2005 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 12

Hulless -- Two-Row

Percentage of seeded acres / () = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
CDC McGwire	39.8 (28.4)	34.8 (82.7)	49.3 (11.5)	0.0 (17.3)	0.0 (0.0)
Other	28.1 (60.4)	65.2 (13.3)	6.0 (76.8)	0.0 (58.2)	0.0 (0.0)
CDC Fibar	20.5 (0.0)	0.0 (0.0)	23.3 (0.0)	100.0 (0.0)	0.0 (0.0)
CDC Alamo	8.5 (0.9)	0.0 (0.0)	15.8 (1.3)	0.0 (0.0)	0.0 (0.0)
Condor	1.6 (1.1)	0.0 (0.0)	3.0 (1.6)	0.0 (0.0)	0.0 (0.0)
CDC Gainer	1.5 (1.2)	0.0 (0.0)	2.7 (1.7)	0.0 (0.0)	0.0 (0.0)
CDC Dawn	0.0 (3.5)	0.0 (0.0)	0.0 (5.3)	0.0 (0.0)	0.0 (0.0)
Phoenix	0.0 (2.4)	0.0 (0.0)	0.0 (0.0)	0.0 (24.5)	0.0 (0.0)
CDC Richard	0.0 (1.3)	0.0 (0.0)	0.0 (1.9)	0.0 (0.0)	0.0 (0.0)
CDC Freedom	0.0 (0.9)	0.0 (4.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)
HB803	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)
HB805	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)
Merlin	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)
Tercel	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)
CDC Candle	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)
CDC Rattan	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)
CDC Speedy	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)

* Trace: less than 0.05 percent

2005 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 13

Hulless -- Six-Row

Percentage of seeded acres / () = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
Other	58.5 (80.5)	100.0 (0.0)	85.7 (87.6)	42.3 (77.1)	0.0 (0.0)
AC Bacon	24.7 (11.5)	0.0 (0.0)	14.3 (12.4)	31.3 (11.2)	0.0 (0.0)
Falcon	16.8 (8.0)	0.0 (100.0)	0.0 (0.0)	26.4 (11.6)	0.0 (0.0)
Jaeger	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)
CDC Buck	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)
CDC Silky	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)
Peregrine	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)
AC Hawkeye	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)

* Trace: less than 0.05 percent

2005 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 14

Feed Barley -- Two-Row

Percentage of seeded acres / () = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
Xena	48.5 (36.9)	3.1 (3.7)	58.3 (44.8)	51.5 (39.7)	90.3 (66.1)
CDC Dolly	21.4 (30.8)	0.9 (5.8)	33.7 (43.2)	20.9 (31.0)	9.7 (21.0)
Other	9.0 (11.8)	2.7 (2.9)	5.7 (7.9)	10.8 (14.4)	0.0 (12.9)
CONLON	8.9 (9.2)	92.7 (84.2)	1.4 (0.6)	0.1 (T)	0.0 (0.0)
Seebe	5.8 (5.5)	0.0 (0.7)	0.1 (0.8)	8.1 (7.7)	0.0 (0.0)
CDC Bold	5.5 (5.7)	0.5 (2.7)	0.7 (2.6)	7.5 (7.2)	0.0 (0.0)
PONOKA	0.8 (0.0)	0.0 (0.0)	0.0 (0.0)	1.2 (0.0)	0.0 (0.0)
CDC Helgason	0.0 (0.1)	0.0 (0.0)	0.0 (0.1)	0.0 (0.1)	0.0 (0.0)
Abee	0.0 (T)	0.0 (0.0)	0.0 (0.0)	0.0 (T)	0.0 (0.0)
CDC Fleet	0.0 (T)	0.0 (0.0)	0.0 (T)	0.0 (0.0)	0.0 (0.0)

* Trace: less than 0.05 percent

2005 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 15

Feed Barley -- Six-Row

Percentage of seeded acres / () = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
Other	38.2 (32.2)	71.1 (34.6)	32.9 (26.7)	35.1 (37.4)	41.7 (5.1)
AC Lacombe	13.9 (19.7)	2.4 (6.6)	12.1 (10.3)	16.7 (30.2)	27.8 (20.0)
AC Rosser	10.3 (10.6)	11.7 (16.2)	18.9 (23.6)	5.1 (0.8)	0.0 (0.0)
Brier	8.5 (11.1)	0.0 (0.0)	21.1 (26.8)	2.6 (4.3)	0.0 (2.6)
Vivar	8.4 (2.7)	0.0 (2.5)	0.0 (0.0)	15.4 (4.9)	0.0 (0.0)
AC Harper	6.9 (7.3)	0.0 (8.2)	3.1 (0.5)	10.6 (12.3)	0.0 (0.0)
Virden	4.9 (3.6)	0.4 (3.4)	11.8 (7.5)	1.7 (1.2)	0.0 (0.0)
Trochu	4.0 (1.3)	0.0 (0.0)	0.0 (1.1)	7.4 (1.9)	0.0 (0.0)
AC Albright	2.1 (2.2)	0.0 (0.0)	0.0 (0.0)	3.7 (1.5)	9.2 (37.9)
Bedford	1.5 (4.2)	14.5 (27.7)	0.1 (0.2)	0.0 (0.3)	0.0 (0.0)
Kasota	0.9 (1.7)	0.0 (T)	0.0 (0.5)	1.1 (1.2)	21.3 (24.7)
PONOKA	0.4 (0.0)	0.0 (0.0)	0.0 (0.0)	0.7 (0.0)	0.0 (0.0)
Bronco	0.0 (1.2)	0.0 (0.8)	0.0 (1.9)	0.0 (0.9)	0.0 (0.0)
Leduc	0.0 (1.0)	0.0 (0.0)	0.0 (0.9)	0.0 (1.3)	0.0 (0.0)
Mahigan	0.0 (0.6)	0.0 (0.0)	0.0 (0.0)	0.0 (0.6)	0.0 (9.7)
CDC Earl	0.0 (0.4)	0.0 (0.0)	0.0 (0.0)	0.0 (0.8)	0.0 (0.0)
Stetson	0.0 (0.2)	0.0 (0.0)	0.0 (0.0)	0.0 (0.4)	0.0 (0.0)

* Trace: less than 0.05 percent

2005 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 16

All Barley (Malting, Feed and Hulless)

Percentage of seeded acres / () = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
Malting -- Two-Row	61.2 (56.4)	40.7 (34.8)	76.7 (70.4)	49.1 (49.1)	62.9 (43.2)
Feed -- (2 & 6 Row)	30.9 (34.7)	29.0 (33.6)	14.1 (18.9)	49.5 (49.6)	37.1 (45.4)
Malting -- Six-Row	7.5 (8.4)	29.3 (30.8)	8.8 (9.9)	1.2 (1.1)	0.0 (11.4)
Hulless -- (2 & 6 Row)	0.4 (0.5)	1.0 (0.8)	0.4 (0.8)	0.2 (0.2)	0.0 (0.0)

* Trace: less than 0.05 percent