

## 2005 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 15

### Feed Barley -- Six-Row

Percentage of seeded acres / ( ) = last year

Variety	Prairie Average		Manitoba		Saskatchewan		Alberta		British Columbia	
Other	<b>38.2</b>	(32.2)	<b>71.1</b>	(34.6)	<b>32.9</b>	(26.7)	<b>35.1</b>	(37.4)	<b>41.7</b>	(5.1)
AC Lacombe	<b>13.9</b>	(19.7)	<b>2.4</b>	(6.6)	<b>12.1</b>	(10.3)	<b>16.7</b>	(30.2)	<b>27.8</b>	(20.0)
AC Rosser	<b>10.3</b>	(10.6)	<b>11.7</b>	(16.2)	<b>18.9</b>	(23.6)	<b>5.1</b>	(0.8)	<b>0.0</b>	(0.0)
Brier	<b>8.5</b>	(11.1)	<b>0.0</b>	(0.0)	<b>21.1</b>	(26.8)	<b>2.6</b>	(4.3)	<b>0.0</b>	(2.6)
Vivar	<b>8.4</b>	(2.7)	<b>0.0</b>	(2.5)	<b>0.0</b>	(0.0)	<b>15.4</b>	(4.9)	<b>0.0</b>	(0.0)
AC Harper	<b>6.9</b>	(7.3)	<b>0.0</b>	(8.2)	<b>3.1</b>	(0.5)	<b>10.6</b>	(12.3)	<b>0.0</b>	(0.0)
Virden	<b>4.9</b>	(3.6)	<b>0.4</b>	(3.4)	<b>11.8</b>	(7.5)	<b>1.7</b>	(1.2)	<b>0.0</b>	(0.0)
Trochu	<b>4.0</b>	(1.3)	<b>0.0</b>	(0.0)	<b>0.0</b>	(1.1)	<b>7.4</b>	(1.9)	<b>0.0</b>	(0.0)
AC Albright	<b>2.1</b>	(2.2)	<b>0.0</b>	(0.0)	<b>0.0</b>	(0.0)	<b>3.7</b>	(1.5)	<b>9.2</b>	(37.9)
Bedford	<b>1.5</b>	(4.2)	<b>14.5</b>	(27.7)	<b>0.1</b>	(0.2)	<b>0.0</b>	(0.3)	<b>0.0</b>	(0.0)
Kasota	<b>0.9</b>	(1.7)	<b>0.0</b>	(T)	<b>0.0</b>	(0.5)	<b>1.1</b>	(1.2)	<b>21.3</b>	(24.7)
PONOKA	<b>0.4</b>	(0.0)	<b>0.0</b>	(0.0)	<b>0.0</b>	(0.0)	<b>0.7</b>	(0.0)	<b>0.0</b>	(0.0)
Bronco	<b>0.0</b>	(1.2)	<b>0.0</b>	(0.8)	<b>0.0</b>	(1.9)	<b>0.0</b>	(0.9)	<b>0.0</b>	(0.0)
Leduc	<b>0.0</b>	(1.0)	<b>0.0</b>	(0.0)	<b>0.0</b>	(0.9)	<b>0.0</b>	(1.3)	<b>0.0</b>	(0.0)
Mahigan	<b>0.0</b>	(0.6)	<b>0.0</b>	(0.0)	<b>0.0</b>	(0.0)	<b>0.0</b>	(0.6)	<b>0.0</b>	(9.7)
CDC Earl	<b>0.0</b>	(0.4)	<b>0.0</b>	(0.0)	<b>0.0</b>	(0.0)	<b>0.0</b>	(0.8)	<b>0.0</b>	(0.0)
Stetson	<b>0.0</b>	(0.2)	<b>0.0</b>	(0.0)	<b>0.0</b>	(0.0)	<b>0.0</b>	(0.4)	<b>0.0</b>	(0.0)

\* Trace: less than 0.05 percent