

## 2006 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 14

### Feed Barley -- Two-Row

Percentage of seeded acres / ( ) = last year

Variety	Prairie Average		Manitoba		Saskatchewan		Alberta		British Columbia	
Xena	<b>62.1</b>	(48.4)	<b>3.2</b>	(3.4)	<b>74.2</b>	(58.0)	<b>68.6</b>	(51.4)	<b>0.0</b>	(90.3)
Conlon	<b>12.2</b>	(8.9)	<b>86.1</b>	(92.2)	<b>5.0</b>	(1.3)	<b>2.6</b>	(0.1)	<b>0.0</b>	(0.0)
CDC Dolly	<b>9.0</b>	(21.6)	<b>0.8</b>	(0.5)	<b>12.7</b>	(33.8)	<b>9.5</b>	(21.1)	<b>0.0</b>	(9.7)
Seebe	<b>4.9</b>	(5.9)	<b>0.0</b>	(0.0)	<b>0.9</b>	(0.1)	<b>6.4</b>	(8.3)	<b>0.0</b>	(0.0)
Other	<b>3.7</b>	(9.1)	<b>0.1</b>	(3.3)	<b>2.1</b>	(6.1)	<b>4.5</b>	(10.7)	<b>0.0</b>	(0.0)
CDC Bold	<b>2.7</b>	(5.3)	<b>0.0</b>	(0.5)	<b>0.1</b>	(0.6)	<b>3.7</b>	(7.2)	<b>0.0</b>	(0.0)
Ponoka	<b>2.4</b>	(0.8)	<b>0.0</b>	(0.0)	<b>0.0</b>	(0.0)	<b>3.2</b>	(1.2)	<b>100.0</b>	(0.0)
TREY	<b>1.5</b>	(0.0)	<b>7.7</b>	(0.0)	<b>1.6</b>	(0.0)	<b>0.5</b>	(0.0)	<b>0.0</b>	(0.0)
CDC Helgalson	<b>1.5</b>	(0.0)	<b>2.1</b>	(0.0)	<b>3.5</b>	(0.0)	<b>1.0</b>	(0.0)	<b>0.0</b>	(0.0)
RIVERS	<b>0.0</b>	(0.0)	<b>0.0</b>	(0.0)	<b>0.0</b>	(0.0)	<b>0.0</b>	(0.0)	<b>0.0</b>	(0.0)

\* Trace: less than 0.05 percent