

# 2006 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 1

## Canada Western Red Spring

Percentage of seeded acres / ( ) = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
AC Barrie	18.8 (21.4)	43.3 (41.0)	17.1 (23.9)	6.1 (8.5)	0.0 (0.0)
Superb	18.1 (18.7)	14.2 (16.2)	16.1 (14.8)	22.8 (24.9)	0.0 (9.4)
McKenzie	9.4 (8.2)	4.4 (2.1)	16.4 (13.6)	4.2 (4.6)	14.3 (5.7)
AC Intrepid	5.5 (6.3)	2.3 (3.8)	5.2 (5.8)	7.4 (8.0)	48.9 (28.9)
Harvest	5.4 (1.4)	3.2 (2.2)	1.6 (0.3)	11.3 (2.5)	0.0 (2.9)
AC Eatonia	4.7 (4.5)	0.0 (0.0)	2.6 (1.7)	10.2 (10.2)	0.0 (0.0)
Prodigy	4.6 (6.7)	0.1 (T)	7.0 (8.5)	4.4 (8.0)	2.9 (2.8)
Lillian	3.9 (0.1)	T (0.0)	4.1 (0.1)	6.1 (0.2)	0.0 (0.0)
Other	3.7 (2.3)	2.2 (1.2)	4.5 (2.1)	3.9 (3.0)	0.8 (2.3)
AC Domain	3.6 (4.6)	15.9 (21.9)	0.5 (1.1)	T (0.2)	0.0 (0.0)
CDC Teal	3.5 (5.0)	1.8 (2.6)	2.5 (3.7)	5.7 (7.7)	6.2 (15.4)
AC Splendor	3.4 (4.6)	1.2 (1.3)	2.8 (4.0)	5.5 (6.7)	27.0 (25.8)
CDC Imagine	3.1 (1.3)	1.5 (0.3)	4.1 (1.7)	3.0 (1.4)	0.0 (0.0)
AC Cadillac	2.8 (3.2)	1.0 (1.5)	5.2 (5.4)	1.1 (1.5)	0.0 (0.0)
AC Elsa	2.1 (2.6)	0.6 (0.8)	3.5 (3.7)	1.2 (2.2)	0.0 (0.0)
AC Abbey	1.7 (1.9)	0.0 (0.0)	2.0 (2.0)	2.3 (2.7)	0.0 (0.0)
Columbus	1.2 (1.7)	T (0.1)	2.3 (2.8)	0.7 (1.1)	0.0 (0.0)
CDC Bounty	1.1 (1.2)	1.5 (1.6)	1.1 (1.2)	0.9 (1.0)	0.0 (0.0)
5602 HR	0.8 (0.0)	2.0 (0.0)	0.2 (0.0)	0.9 (0.0)	0.0 (0.0)
5601 HR	0.8 (0.2)	3.3 (1.3)	0.1 (T)	T (T)	0.0 (0.0)
Journey	0.4 (0.1)	0.3 (0.1)	0.1 (0.1)	0.9 (0.2)	0.0 (0.0)
Lovitt	0.3 (0.1)	0.8 (T)	0.2 (0.1)	0.2 (T)	0.0 (0.0)
5500 HR	0.2 (0.1)	T (0.2)	0.1 (T)	0.5 (0.2)	0.0 (0.0)
5600 HR	0.2 (0.2)	0.1 (0.2)	0.2 (0.1)	0.3 (0.4)	0.0 (0.0)
Infinity	0.2 (0.0)	0.1 (0.0)	0.3 (0.0)	0.1 (0.0)	0.0 (0.0)
Alsen	0.1 (0.1)	0.2 (0.5)	0.1 (T)	0.0 (T)	0.0 (0.0)
CDC Go	0.1 (T)	0.0 (0.0)	T (0.0)	0.2 (T)	0.0 (0.0)
CDC Osler	0.1 (T)	0.0 (0.0)	0.1 (T)	T (0.0)	0.0 (0.0)
Somerset	T (0.0)	0.1 (0.0)	T (0.0)	0.0 (0.0)	0.0 (0.0)
Peace	T (0.0)	0.0 (0.0)	0.0 (0.0)	T (0.0)	0.0 (0.0)
CDC Alsask	T (0.0)	0.0 (0.0)	T (0.0)	0.0 (0.0)	0.0 (0.0)
Katepwa	0.0 (1.5)	0.0 (0.1)	0.0 (2.0)	0.0 (1.4)	0.0 (0.0)
Roblin	0.0 (1.2)	0.0 (T)	0.0 (0.2)	0.0 (3.1)	0.0 (6.8)
Laura	0.0 (0.6)	0.0 (0.0)	0.0 (1.0)	0.0 (0.6)	0.0 (0.0)
AC Cora	0.0 (0.2)	0.0 (0.9)	0.0 (0.1)	0.0 (0.0)	0.0 (0.0)

\* Trace: less than 0.05 percent

## 2006 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 2

### Canada Western Amber Durum

Percentage of seeded acres / ( ) = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
AC Avonlea	38.4 (45.3)	0.0 (7.0)	37.0 (44.4)	43.6 (49.1)	0.0 (0.0)
Kyle	23.0 (34.8)	0.0 (61.7)	24.9 (37.9)	16.2 (23.6)	0.0 (0.0)
Strongfield	19.2 (0.0)	0.0 (0.0)	17.5 (0.0)	25.6 (0.0)	0.0 (0.0)
AC Navigator	11.2 (9.2)	0.0 (0.0)	12.4 (9.6)	6.9 (8.0)	0.0 (0.0)
AC Morse	2.9 (5.7)	0.0 (5.9)	1.9 (3.4)	6.6 (14.0)	0.0 (0.0)
Napoleon	2.1 (1.5)	100.0 (25.5)	2.6 (1.6)	0.0 (0.6)	0.0 (0.0)
Commander	1.8 (0.0)	0.0 (0.0)	2.3 (0.0)	0.1 (0.0)	0.0 (0.0)
Other	1.3 (1.6)	0.0 (0.0)	1.4 (1.6)	0.8 (1.8)	0.0 (0.0)
Plenty	0.0 (0.8)	0.0 (0.0)	0.0 (0.4)	0.0 (1.9)	0.0 (0.0)
Wakooma	0.0 (0.5)	0.0 (0.0)	0.0 (0.6)	0.0 (0.3)	0.0 (0.0)
Sceptre	0.0 (0.2)	0.0 (0.0)	0.0 (0.2)	0.0 (0.4)	0.0 (0.0)
AC Melita	0.0 (0.2)	0.0 (0.0)	0.0 (0.1)	0.0 (0.4)	0.0 (0.0)
Medora	0.0 (0.2)	0.0 (0.0)	0.0 (0.2)	0.0 (0.0)	0.0 (0.0)

\* Trace: less than 0.05 percent

# 2006 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 3

## Canada Western Hard White

Percentage of seeded acres / ( ) = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
Snowbird	96.8 (97.5)	98.4 (97.6)	97.2 (96.4)	93.8 (99.5)	0.0 (100.0)
Kanata	3.2 (2.5)	1.6 (2.4)	2.8 (3.6)	6.2 (0.5)	0.0 (0.0)

\* Trace: less than 0.05 percent

# 2006 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 4

## Canada Western Extra Strong

Percentage of seeded acres / ( ) = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
Glenlea	<b>53.8</b> (34.1)	<b>100.0</b> (0.0)	<b>79.0</b> (45.2)	<b>0.0</b> (13.4)	<b>0.0</b> (0.0)
Laser	<b>18.0</b> (18.1)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>55.9</b> (59.7)	<b>0.0</b> (0.0)
Bluesky	<b>14.2</b> (5.6)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>44.1</b> (18.5)	<b>0.0</b> (0.0)
CDC Rama	<b>14.1</b> (12.6)	<b>0.0</b> (0.0)	<b>21.0</b> (19.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)
Amazon	<b>0.0</b> (14.9)	<b>0.0</b> (0.0)	<b>0.0</b> (22.5)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)
AC Corinne	<b>0.0</b> (8.8)	<b>0.0</b> (0.0)	<b>0.0</b> (13.2)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)
Burnside	<b>0.0</b> (3.3)	<b>0.0</b> (100.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)
Wildcat	<b>0.0</b> (2.5)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (8.4)	<b>0.0</b> (0.0)
Walrus	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)
Glenavon	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)

\* Trace: less than 0.05 percent

# 2006 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 5

## Canada Prairie Spring White

Percentage of seeded acres / ( ) = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
AC Vista	<b>93.6</b> (58.0)	<b>100.0</b> (100.0)	<b>92.9</b> (52.6)	<b>0.0</b> (70.7)	<b>0.0</b> (0.0)
AC Karma	<b>6.4</b> (42.0)	<b>0.0</b> (0.0)	<b>7.1</b> (47.4)	<b>0.0</b> (29.3)	<b>0.0</b> (0.0)
Other	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)

\* Trace: less than 0.05 percent

# 2006 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 6

## Canada Prairie Spring Red

Percentage of seeded acres / ( ) = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
5700PR	37.3 (34.3)	52.6 (54.6)	31.9 (39.1)	39.2 (32.4)	100.0 (0.0)
AC Crystal	32.0 (35.0)	0.0 (0.0)	45.0 (45.1)	26.5 (33.6)	0.0 (0.0)
AC Foremost	16.4 (12.3)	0.0 (0.0)	1.3 (3.7)	25.2 (14.9)	0.0 (0.0)
5701PR	12.3 (11.4)	11.3 (40.0)	20.7 (8.3)	8.0 (11.3)	0.0 (0.0)
AC Taber	2.0 (6.2)	36.1 (1.2)	1.1 (3.8)	1.1 (7.0)	0.0 (0.0)
Oslo	0.0 (0.8)	0.0 (4.2)	0.0 (0.0)	0.0 (0.8)	0.0 (0.0)
Other	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)

\* Trace: less than 0.05 percent

# 2006 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 7

## Canada Western Red Winter

Percentage of seeded acres / ( ) = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
CDC Falcon	37.7 (20.0)	73.9 (68.5)	8.3 (10.3)	1.1 (2.0)	0.0 (0.0)
AC Bellatrix	14.9 (25.0)	0.0 (0.5)	3.8 (18.9)	61.2 (40.8)	0.0 (0.0)
CDC Buteo	11.5 (1.8)	7.5 (4.2)	26.6 (2.9)	0.0 (0.0)	0.0 (0.0)
CDC Clair	8.7 (19.1)	1.1 (3.9)	21.3 (33.7)	7.8 (17.7)	0.0 (0.0)
CDC Raptor	6.6 (7.6)	3.7 (11.3)	16.0 (17.0)	0.0 (0.0)	0.0 (0.0)
Radiant	5.0 (1.8)	0.0 (0.0)	3.6 (1.3)	17.4 (3.0)	0.0 (0.0)
McClintock	4.5 (1.1)	7.8 (4.2)	2.6 (0.3)	0.0 (0.0)	0.0 (0.0)
CDC Osprey	3.8 (9.1)	0.0 (0.0)	10.3 (13.7)	3.1 (10.7)	0.0 (0.0)
CDC Harrier	2.9 (1.1)	4.1 (4.0)	3.4 (0.6)	0.0 (0.0)	0.0 (0.0)
AC Tempest	1.6 (1.8)	0.0 (0.0)	1.4 (0.0)	5.3 (3.8)	0.0 (0.0)
Norstar	1.3 (1.1)	1.6 (1.2)	1.8 (0.3)	0.0 (1.6)	0.0 (0.0)
AC Readymade	0.9 (7.4)	0.0 (0.0)	0.0 (0.0)	3.8 (15.6)	0.0 (0.0)
CDC Kestrel	0.5 (1.4)	0.4 (2.2)	1.0 (0.9)	0.3 (1.4)	0.0 (0.0)
Other	0.0 (1.6)	0.0 (0.0)	0.0 (0.0)	0.0 (3.4)	0.0 (0.0)

\* Trace: less than 0.05 percent

# 2006 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 8

## Canada Western Soft White Spring

Percentage of seeded acres / ( ) = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
AC Andrew	<b>83.5</b> (70.0)	<b>0.0</b> (0.0)	<b>100.0</b> (100.0)	<b>52.2</b> (65.0)	<b>0.0</b> (0.0)
AC Meena	<b>11.6</b> (14.2)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>33.7</b> (16.6)	<b>0.0</b> (0.0)
AC Nanda	<b>3.5</b> (3.7)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>10.3</b> (4.3)	<b>0.0</b> (0.0)
AC Reed	<b>1.3</b> (2.6)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>3.9</b> (3.1)	<b>0.0</b> (0.0)
Bhishaj	<b>0.0</b> (9.5)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (11.0)	<b>0.0</b> (0.0)
AC Phil	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)

\* Trace: less than 0.05 percent



## 2006 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 9

### Malting -- Two-Row

Percentage of seeded acres / ( ) = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
AC Metcalfe	59.3 (55.1)	61.5 (59.9)	57.8 (54.3)	60.2 (55.0)	85.8 (98.0)
CDC Copeland	14.6 (12.3)	12.2 (14.3)	15.8 (14.0)	13.8 (9.1)	0.0 (0.0)
CDC Kendall	13.4 (15.2)	0.6 (3.1)	19.2 (20.3)	8.8 (9.2)	0.0 (0.0)
Harrington	4.7 (8.6)	0.0 (0.0)	3.5 (5.4)	7.1 (15.9)	14.2 (2.0)
Newdale	2.2 (1.3)	16.4 (12.7)	0.7 (0.6)	0.9 (0.4)	0.0 (0.0)
Other	1.8 (1.6)	1.7 (0.7)	0.8 (0.8)	3.1 (3.1)	0.0 (0.0)
Merit	1.8 (1.9)	0.0 (0.0)	0.9 (1.8)	3.3 (2.4)	0.0 (0.0)
CDC Stratus	1.5 (1.7)	7.6 (9.4)	1.2 (1.2)	0.6 (0.9)	0.0 (0.0)
Stein	0.8 (1.8)	0.0 (0.0)	0.0 (1.5)	2.0 (2.6)	0.0 (0.0)
CDC Select	0.1 (T)	0.0 (0.0)	0.1 (T)	0.2 (0.0)	0.0 (0.0)
B1202	0.0 (0.4)	0.0 (0.0)	0.0 (0.1)	0.0 (0.9)	0.0 (0.0)
AC Bountiful	0.0 (0.2)	0.0 (0.0)	0.0 (0.1)	0.0 (0.5)	0.0 (0.0)
Calder	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)

\* Trace: less than 0.05 percent

## 2006 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 10

### Malting -- Six-Row

Percentage of seeded acres / ( ) = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
Legacy	43.0 (30.4)	35.9 (16.9)	50.9 (42.0)	27.8 (16.5)	0.0 (0.0)
Robust	20.8 (19.3)	31.2 (41.2)	7.6 (5.1)	52.9 (4.5)	0.0 (0.0)
Excel	12.6 (30.5)	15.5 (23.1)	11.2 (36.5)	9.8 (25.9)	0.0 (0.0)
Tradition	12.3 (2.7)	7.5 (2.3)	18.0 (3.3)	0.0 (0.0)	0.0 (0.0)
Lacey	8.1 (8.6)	8.2 (11.8)	8.8 (7.3)	3.7 (0.0)	0.0 (0.0)
CDC Sisler	1.2 (1.1)	0.0 (0.3)	2.3 (1.4)	0.0 (3.6)	0.0 (0.0)
B1602	0.9 (2.0)	0.6 (0.0)	0.5 (1.8)	2.0 (15.3)	100.0 (0.0)
CDC Battleford	0.7 (1.9)	0.9 (0.1)	0.1 (0.8)	2.8 (21.5)	0.0 (0.0)
Other	0.5 (1.7)	0.2 (2.7)	0.7 (1.0)	1.1 (0.6)	0.0 (0.0)
CDC Yorkton	0.0 (1.3)	0.0 (0.4)	0.0 (0.6)	0.0 (11.5)	0.0 (0.0)
CDC Tisdale	0.0 (0.4)	0.0 (1.1)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)
CDC Springside	0.0 (0.2)	0.0 (0.0)	0.0 (0.2)	0.0 (0.6)	0.0 (0.0)
CDC Clyde	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)
CDC Laurence	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)

\* Trace: less than 0.05 percent

## 2006 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 11

### Two-Row and Six-Row Malting Barley

Percentage of seeded acres / ( ) = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
Malting -- Two-Row	84.5 (89.0)	56.2 (58.0)	84.1 (89.8)	95.7 (97.6)	94.6 (100.0)
Malting -- Six-Row	15.5 (11.0)	43.8 (42.0)	15.9 (10.2)	4.3 (2.4)	5.4 (0.0)

\* Trace: less than 0.05 percent

# 2006 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 12

## Hulless -- Two-Row

Percentage of seeded acres / ( ) = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
Other	100.0 (2.6)	100.0 (0.0)	100.0 (3.6)	100.0 (0.0)	0.0 (0.0)
CDC McGwire	0.0 (54.0)	0.0 (100.0)	0.0 (50.6)	0.0 (0.0)	0.0 (0.0)
CDC Fibar	0.0 (27.8)	0.0 (0.0)	0.0 (23.8)	0.0 (100.0)	0.0 (0.0)
CDC Alamo	0.0 (11.5)	0.0 (0.0)	0.0 (16.2)	0.0 (0.0)	0.0 (0.0)
Condor	0.0 (2.2)	0.0 (0.0)	0.0 (3.0)	0.0 (0.0)	0.0 (0.0)
CDC Gainer	0.0 (2.0)	0.0 (0.0)	0.0 (2.8)	0.0 (0.0)	0.0 (0.0)

\* Trace: less than 0.05 percent

# 2006 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 13

## Hulless -- Six-Row

Percentage of seeded acres / ( ) = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
Other	100.0 (58.5)	0.0 (100.0)	100.0 (85.7)	100.0 (42.3)	0.0 (0.0)
AC Bacon	0.0 (24.7)	0.0 (0.0)	0.0 (14.3)	0.0 (31.3)	0.0 (0.0)
Falcon	0.0 (16.8)	0.0 (0.0)	0.0 (0.0)	0.0 (26.4)	0.0 (0.0)

\* Trace: less than 0.05 percent

## 2006 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 14

### Feed Barley -- Two-Row

Percentage of seeded acres / ( ) = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
Xena	<b>63.0</b> (48.4)	<b>3.6</b> (3.4)	<b>73.7</b> (58.0)	<b>70.1</b> (51.4)	<b>0.0</b> (90.3)
Conlon	<b>12.3</b> (0.0)	<b>85.8</b> (0.0)	<b>5.8</b> (0.0)	<b>2.6</b> (0.0)	<b>0.0</b> (0.0)
CDC Dolly	<b>7.7</b> (21.6)	<b>0.0</b> (0.5)	<b>12.5</b> (33.8)	<b>8.1</b> (21.1)	<b>0.0</b> (9.7)
Other	<b>4.8</b> (9.1)	<b>1.3</b> (3.3)	<b>3.1</b> (6.1)	<b>5.4</b> (10.7)	<b>85.7</b> (0.0)
Seebe	<b>4.3</b> (5.9)	<b>0.0</b> (0.0)	<b>0.9</b> (0.1)	<b>5.5</b> (8.3)	<b>0.0</b> (0.0)
CDC Bold	<b>3.2</b> (5.3)	<b>0.0</b> (0.5)	<b>0.0</b> (0.6)	<b>4.2</b> (7.2)	<b>0.0</b> (0.0)
Ponoka	<b>2.1</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>2.7</b> (0.0)	<b>14.3</b> (0.0)
TREY	<b>1.4</b> (0.0)	<b>7.0</b> (0.0)	<b>1.0</b> (0.0)	<b>0.6</b> (0.0)	<b>0.0</b> (0.0)
CDC Helgalson	<b>1.3</b> (0.0)	<b>2.3</b> (0.0)	<b>3.1</b> (0.0)	<b>0.8</b> (0.0)	<b>0.0</b> (0.0)
CONLON	<b>0.0</b> (8.9)	<b>0.0</b> (92.2)	<b>0.0</b> (1.3)	<b>0.0</b> (0.1)	<b>0.0</b> (0.0)
PONOKA	<b>0.0</b> (0.8)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (1.2)	<b>0.0</b> (0.0)
RIVERS	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)	<b>0.0</b> (0.0)

\* Trace: less than 0.05 percent

## 2006 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 15

### Feed Barley -- Six-Row

Percentage of seeded acres / ( ) = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
AC RANGER	23.4 (0.0)	59.8 (0.0)	13.9 (0.0)	17.2 (0.0)	0.0 (0.0)
Stander	13.4 (0.0)	8.9 (0.0)	13.6 (0.0)	15.1 (0.0)	0.0 (0.0)
Brier	10.8 (7.9)	0.0 (0.0)	30.4 (20.2)	4.5 (2.6)	0.0 (0.0)
Other	10.3 (42.2)	12.3 (70.4)	8.2 (34.3)	10.6 (41.7)	18.3 (51.9)
AC Lacombe	8.8 (12.9)	2.0 (2.3)	11.7 (11.6)	9.2 (15.1)	22.9 (23.0)
Vivar	8.2 (7.6)	0.0 (0.0)	0.0 (0.0)	15.3 (13.3)	0.0 (0.0)
Trochu	6.3 (3.7)	0.0 (0.0)	0.3 (0.0)	11.5 (6.4)	0.0 (0.0)
AC Rosser	5.5 (9.8)	5.4 (11.6)	11.8 (18.1)	2.5 (5.2)	0.0 (0.0)
AC Harper	3.8 (6.2)	0.0 (0.0)	0.0 (3.0)	7.1 (9.2)	0.0 (0.0)
Virden	2.8 (4.9)	0.0 (0.4)	10.2 (12.7)	0.0 (1.5)	0.0 (0.0)
AC Albright	2.4 (2.0)	0.0 (0.0)	0.0 (0.2)	3.0 (3.2)	58.8 (7.6)
Bedford	2.0 (1.5)	11.6 (15.3)	0.0 (0.1)	0.0 (0.0)	0.0 (0.0)
Kasota	1.3 (1.0)	0.0 (0.0)	0.0 (0.0)	2.4 (1.2)	0.0 (17.6)
CDC Earl	0.8 (0.0)	0.0 (0.0)	0.0 (0.0)	1.5 (0.0)	0.0 (0.0)
PONOKA	0.0 (0.3)	0.0 (0.0)	0.0 (0.0)	0.0 (0.6)	0.0 (0.0)
MANNY	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)

\* Trace: less than 0.05 percent

## 2006 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 16

### All Barley (Malting, Feed and Hulless)

Percentage of seeded acres / ( ) = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
Malting -- Two-Row	49.9 (61.0)	34.5 (41.0)	69.7 (76.6)	39.1 (48.6)	68.2 (61.2)
Feed -- (2 & 6 Row)	40.1 (31.2)	38.1 (29.0)	15.4 (14.3)	59.0 (50.0)	27.9 (38.8)
Malting -- Six-Row	9.2 (7.6)	26.9 (29.6)	13.1 (8.7)	1.7 (1.2)	3.9 (0.0)
Hulless -- (2 & 6 Row)	0.8 (0.3)	0.4 (0.4)	1.7 (0.4)	0.2 (0.2)	0.0 (0.0)

\* Trace: less than 0.05 percent



## 2006 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 17

### All Wheat

Percentage of seeded acres / ( ) = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
Canada Western Red Spring	<b>69.2</b> (69.2)	<b>84.9</b> (84.9)	<b>60.4</b> (60.4)	<b>74.8</b> (74.8)	<b>95.4</b> (95.4)
Canada Western Amber Durum	<b>21.4</b> (21.4)	<b>0.2</b> (0.2)	<b>33.2</b> (33.2)	<b>14.0</b> (14.0)	<b>0.0</b> (0.0)
Canada Western Hard White	<b>3.9</b> (3.9)	<b>8.8</b> (8.8)	<b>3.0</b> (3.0)	<b>2.7</b> (2.7)	<b>2.3</b> (2.3)
Canada Western Red Winter	<b>2.5</b> (2.5)	<b>5.6</b> (5.6)	<b>1.5</b> (1.5)	<b>2.6</b> (2.6)	<b>0.0</b> (0.0)
Canada Prairie Spring Red	<b>2.5</b> (2.5)	<b>0.4</b> (0.4)	<b>1.2</b> (1.2)	<b>5.3</b> (5.3)	<b>2.3</b> (2.3)
Canada Western Soft White Spring	<b>0.2</b> (0.2)	<b>0.0</b> (0.0)	<b>0.2</b> (0.2)	<b>0.4</b> (0.4)	<b>0.0</b> (0.0)
Canada Prairie Spring White	<b>0.2</b> (0.2)	<b>0.1</b> (0.1)	<b>0.3</b> (0.3)	<b>T</b> (T)	<b>0.0</b> (0.0)
Canada Western Extra Strong	<b>0.1</b> (0.1)	<b>T</b> (T)	<b>0.1</b> (0.1)	<b>0.1</b> (0.1)	<b>0.0</b> (0.0)

\* Trace: less than 0.05 percent