

SCHEDULE I
Sections 5 to 7, 11 and 12, 18, 23 to 27, 30 and 40
TABLE I

Applicable to: (a) Wheat, common - *Triticum aestivum* L. emend. Fiori et Paol
(b) Wheat, durum - *Triticum durum* Desf.

I	II	III	IV	V	VI	VII	VIII	IX	X
	Maximum Number of Seeds per kg								
	Noxious Weeds							Minimum Percent Germination	
GRADE NAME	PRIMARY	PRIMARY PLUS SECONDARY	TOTAL WEEDS	TOTAL BARLEY, RYE AND TRITICALE	TOTAL OTHER CROPS INCLUDING BARLEY, RYE & TRITICALE	ADDITIONAL COMMON WHEAT IN DURUM WHEAT AND DURUM WHEAT IN COMMON WHEAT	MAXIMUM NUMBER OF ERGOT BODIES PER KILOGRAM	COMMON WHEAT	DURUM WHEAT
1. Canada Foundation No. 1	0	0.0	2	0.2	0.5	0	1	85	80
2. Canada Foundation No. 2	0	0.1	4	1.0	2.0	0	8	75	70
3. Canada Registered No. 1	0	0.0	3	0.5	1.0	0	1	85	80
4. Canada Registered No. 2	0	0.1	6	1.0	2.0	0	8	75	70
5. Canada Certified No. 1	0	0.0	3	1.0	2.0	5	1	85	80
6. Canada Certified No. 2	0	0.5	6	2.0	5.0	10	8	75	70
7. Common No. 1	0	2.0	10	10.0	10.0	12	1	85	80
8. Common No. 2	2	4.0	20	20.0	20.0	24	8	70	70

1. Seed to which Table 1 to Schedule I applies shall be free from Tartarian buckwheat in Manitoba, Saskatchewan, Alberta and British Columbia.
2. Seed to which Table I to Schedule I applies shall be free from wild oats in Quebec, Nova Scotia, New Brunswick and Prince Edward Island.

TABLE II

Applicable to:

- | | | | |
|--|--|--|--|
| (a) Barley
-- six-row
-- two-row | - <i>Hordeum vulgare</i> L. convar. <i>hexastichon</i> Alef.
- <i>Hordeum vulgare</i> L. convar. <i>distichon</i> Alef. | (l) Rye
(j) Sainfoin
(k) Spelt
(l) Triticale
(m) Vetch, hairy
(n) Vetch, Hungarian
(o) Vetch, spring or common | - <i>Secale cereale</i> L.
- <i>Onobrychis viciifolia</i> Scop.
- <i>Triticum spelta</i> L.
- <i>X Triticosecale</i> Wittmack
- <i>Vicia villosa</i> Roth. ssp. <i>villosa</i>
- <i>Vicia pannonica</i> Crantz
- <i>Vicia sativa</i> L. ssp. <i>sativa</i> |
| (b) Bean, mung | - <i>Vigna radiata</i> (L.) Wilczek var. <i>radiata</i> | | |
| (c) Buckwheat, common | - <i>Fagopyrum esculentum</i> Moench | | |
| (d) Buckwheat, tartarian | - <i>Fagopyrum tataricum</i> (L.) Gaertner | | |
| (e) Emmer | - <i>Triticum dicoccum</i> Schrank | | |
| (f) Lentil | - <i>Lens culinaris</i> Medikus | | |
| (g) Lupine, field (non-ornamental) | - <i>Lupinus</i> spp. | | |
| (h) Oat | - <i>Avena sativa</i> L., <i>A. nuda</i> L. | | |

I	II		III		IV	V	VI	VII	VIII	IX	X	XI
	Maximum Number of Seeds Per kg											
GRADE NAME	PRIMARY	PRIMARY PLUS SECONDARY		TOTAL WEEDS	TOTAL OTHER CROPS	Maximum Number of Ergot Bodies per kg		MAXIMUM PER CENT TRUE LOOSE SMUT IN BARLEY	Minimum Per cent Germination			
		IN OATS	IN OTHER KINDS			IN BARLEY & OATS	IN TRITICALE & RYE		RYE, TRITICALE & SAINFOIN	OTHER KINDS		
1. Canada Foundation No. 1	0	0.0	0.0	2	1	1	2	2	75	85		
2. Canada Foundation No. 2	0	0.1	0.2	4	2	8	10	4	65	75		
3. Canada Registered No. 1	0	0.0	0.0	3	2	1	2	2	75	85		
4. Canada Registered No. 2	0	0.1	0.2	6	4	8	10	4	65	75		
5. Canada Certified No. 1	0	0.5	0.2	3	4	2	4	2	75	85		
6. Canada Certified No. 2	0	1.0	1.0	6	10	8	15	4	65	75		
7. Common No. 1	0	2.0	2.0	10	25	2	4	4	75	85		
8. Common No. 2	2	4.0	4.0	20	50	8	15	6	65	75		

- Seed to which Table II applies, shall be free from Tartarian buckwheat in Manitoba, Saskatchewan, Alberta and British Columbia.
- Seed to which Table II applies shall be free from Wild oats in Quebec, Nova Scotia, New Brunswick and Prince Edward Island.
- In the grading of seeds of lentils referred to in Table II to Schedule I, a seed head of mallow (*Maha* spp.) shall be considered as a secondary noxious weed.
- Barley seed that has been treated with a product registered as a control product under the *Pest Control Products Act*, for the control of true loose smut (*Ustilago nuda*) is exempt from the standards for true loose smut set out in Column IX of Table II to Schedule I.

Note: In Grade Table II, false wild oats are considered as oats in the grading of oats. However, if found in any crop other than oats, it should be considered as other crop seeds.

TABLE III

Applicable to:

Cereal mixtures composed of seeds of two or more kinds listed in Tables I and II, including field beans, horse beans, tick beans and peas.

I	II	III	IV	V	VI	VII	VIII	IX
	Maximum Number of Seeds Per Kilogram							
	Noxious Weeds						Minimum Percent Germination for each Component	
GRADE NAME	PRIMARY	PRIMARY PLUS SECONDARY	TOTAL WEEDS	TOTAL OTHER CROPS	MAXIMUM PERCENT TRUE LOOSE SMUT IN BARLEY	MAXIMUM NUMBER OF ERGOT BODIES PER KILOGRAM	DURUM WHEAT	OTHER SPECIES
1. Canada Certified No. 1 Cereal Mixture	0	0.2	3	5	2	3	80	85
2. Canada Certified No. 2 Cereal Mixture	0	1.0	6	20	4	12	70	75
3. Common No. 1 Cereal Mixture	0	2.0	10	30	4	3	80	85
4. Common No. 2 Cereal Mixture	2	4.0	20	60	6	12	70	75

1. Seed to which Table III to applies shall be free from Tartarian buckwheat in Manitoba, Saskatchewan, Alberta and British Columbia.
2. Seed to which Table III to Schedule I applies shall be free from wild oats in Quebec, Nova Scotia, New Brunswick and Prince Edward Island.

Note: In Grade Table III, false wild oats are considered as oats in the grading of oats. However, if found in any crop other than oats, it should be considered as other crop seeds.

TABLE IV

Applicable to:

- (a) Canarygrass - *Pbalaris canariensis* L.
- (b) Flax, oil and fibre - *Linum usitatissimum* L.
- (c) Sorghum - *Sorghum bicolor* (L.) Moench (= *S. vulgare* Pers.), *S. alnum* Parodi
- (d) Sorghum Sudangrass hybrids
- (e) Sudan grass - *Sorghum sudanense* (Piper) Stapf

I	II	III	IV	V	VI	VII
	Maximum Number of Seeds per 25 g					
	Noxious Weeds			Other Crops		
GRADE NAME	PRIMARY	PRIMARY PLUS SECONDARY	TOTAL WEEDS	IN SORGHUM, SUDANGRASS and CANARYGRASS	IN OTHER KINDS	Minimum Percent Germination
1. Canada Foundation No. 1	0	0.0	2	0.5	0.5	85
2. Canada Foundation No. 2	0	0.5	3	1.0	1.0	70
3. Canada Registered No. 1	0	0.0	2	0.5	1.0	85
4. Canada Registered No. 2	0	0.5	3	1.0	2.0	70
5. Canada Certified No. 1	0	0.5	3	1.0	1.0	85
6. Canada Certified No. 2	0	2.0	5	2.0	2.0	70
7. Common No.1	0	2.0	6	25.0	2.0	85
8. Common No. 2	2	6.0	10	50.0	8.0	70

TABLE V

Applicable to:

- (a) Bean, field and garden - *Phaseolus vulgaris* L.
- (b) Bean, broad, horse, tick and faba - *Vicia faba* L.
- (c) Bean, lima - *Phaseolus lunatus* L.
- (d) Bean, scarlet runner - *Phaseolus coccineus* L.
- (e) Chickpea - *Cicer arietinum* L.
- (f) Corn (open-pollinated, synthetic) - *Zea mays* L.
- (g) Cowpea - *Vigna unguiculata* (L.) Walpers ssp. *unguiculata*
- (h) Pea, canning, field and garden - *Pisum sativum* L.
- (I) Safflower - *Carthamus tinctorius* L.
- (j) Soybean - *Glycine max* (L.) Merrill
- (k) Sunflower (open-pollinated) - *Helianthus annuus* L.

I	II	III	IV	V	VI	VII
	Maximum Number of Seeds per kg			Minimum Percent Germination		
GRADE NAME	TOTAL WEEDS	OTHER CROP SEEDS	MAXIMUM NUMBER OF SCLEROTIA BODIES IN SUNFLOWER PER KG	PEAS	BEANS & SOYBEANS	OTHER KINDS
1. Canada Foundation No. 1	0	0.0	1	80	85	90
2. Canada Foundation No. 2	0	0.5	8	70	75	80
3. Canada Registered No. 1	0	0.0	1	80	85	90
4. Canada Registered No. 2	0	1.0	8	70	75	80
5. Canada Certified No. 1	0	1.0	1	80	85	90
6. Canada Certified No. 2	0	3.0	8	70	75	80
7. Common No. 1	1	4.0	4	75	80	90
8. Common No. 2	3	10.0	8	65	65	75

TABLE VI

Applicable to:

- (a) Corn, hybrid - *Zea mays L.*
- (b) Sunflower, hybrid - *Helianthus annuus L.*

I	II	III	IV	V
	Maximum Number of Seeds per kg			
GRADE NAME	WEEDS	OTHER CROPS	MAXIMUM NUMBER OF SCLEROTIA BODIES PER KG IN SUNFLOWER	Minimum Percent Germination
1. Canada Foundation	0	N/A	8	N/A
2. Canada Certified No. 1	0	1	1	90
3. Canada Certified No. 2	0	5	8	80
4. Common	4	10	8	75

1. Hybrid seed of corn and hybrid seed of sunflower, sold or offered for sale as a Canada Foundation grade shall be labelled with the total number of seeds of other crops, per kilogram and the percentage germination.

TABLE VII

Applicable to:

- (a) Mustard, black - *Brassica nigra* (L.) Koch
- (b) Mustard, oriental or Indian - *Brassica juncea* (L.) Czern. et Coss.
- (c) Mustard, white (=yellow) - *Sinapis alba* L. (= *B. hirta* Moench)
- (d) Rapeseed, Oilseed Rape including Canola
 - polish type - *Brassica rapa* L.
 - argentine type - *Brassica napus* L.

I	II	III	IV	V	VI	VII	VIII	XI
	Maximum Number of Seeds per 25 g							
	Noxious Weeds							
GRADE NAME	PRIMARY	PRIMARY PLUS SECONDARY	TOTAL WEEDS	<i>B. rapa</i> IN <i>B. napus</i> OR <i>B. napus</i> IN <i>B. rapa</i>	OTHER BRASSICA CROPS INCLUDING <i>S. alba</i>	NON-BRASSICA CROPS	MAXIMUM NO. OF SCLEROTIA BODIES PER 50 G	MINIMUM PERCENT GERMINATION
1. Canada Foundation No. 1	0	0	5	0	0	1	1	90
2. Canada Foundation No. 2	0	0	10	2	1	2	2	75
3. Canada Registered No. 1	0	0	5	0	0	1	1	90
4. Canada Registered No. 2	0	0	10	2	1	2	2	75
5. Canada Certified No. 1	0	1	5	5	1	1	1	90
6. Canada Certified No. 2	0	2	10	10	5	2	2	80
7. Common	1	3	15	20	15	2	2	90

1. Seed to which Table VII applies shall be free from seeds of cleavers (*Galium aparine* L. and *Galium spurium* L.)

TABLE VIII

Applicable to:

- (a) Alfalfa - *Medicago sativa* L. (incl. *M. sativa* L. ssp. *falcata* (L.) Arcangeli)
- (b) Clover, crimson - *Trifolium incarnatum* L.
- (c) Clover, red - *Trifolium pratense* L.
- (d) Clover, subterranean - *Trifolium subterraneum* L.
- (e) Clover, sweet
 - white blossom - *Melilotus alba* Medikus
 - yellow blossom - *Melilotus officinalis* (L.) Pallas
- (f) Lespedeza, common or Kobe - *Lespedeza striata* (Thunb. ex Murray) Hook. & Arn.
- (g) Lespedeza, Korean - *Lespedeza stipulacea* Maxim.
- (h) Lespedeza, sericea or Chinese - *Lespedeza cuneata* (Dum.-Cours.) G. Don
- (i) Millet, foxtail or Italian - *Setaria italica* (L.) Beauv.
- (j) Millet, Japanese - *Echinochloa crusgalli* (L.) Beauv. var. *frumentacea* (Link) W.F. Wight
- (k) Millet, pearl - *Pennisetum americanum* (L.) Schumann (= *P. typhoides* (Burm.) Stapf & C.E. Hubb.)
- (l) Millet, proso - *Panicum miliaceum* L.
- (m) Vetch, crown - *Cornonilla varia* L.
- (n) Vetch, kidney - *Anthyllis vulneraria* L.
- (o) Vetch, milk, cicer - *Astragalus cicer* L.

I	II	III	IV	V	VI	VII	VIII
	Maximum Number of Seeds per 25 g Except Where Otherwise Stated						
	Noxious Weeds			Other Crops			
GRADE NAME	PRIMARY	PRIMARY PLUS SECONDARY	TOTAL WEEDS	SWEET-CLOVER	BRASSICA CROPS INCLUDING <i>S. alba</i>	OTHER NON-BRASSICA CROPS	MINIMUM PERCENT GERMINATION
1. Canada Foundation No. 1	0	0	5	1	0	5	85
2. Canada Foundation No. 2	0	2	30	2	2	20	70
3. Canada Registered No. 1	0	0	5	1	0	5	85
4. Canada Registered No. 2	0	2	30	2	2	20	70
5. Canada Certified No. 1	0	5	50	20	4	1% by weight	80
6. Canada Certified No. 2	0	10	75	30	6	2% by weight	70
7. Varietal Blend No. 1	0	5	50	20	4	1% by weight	80
8. Varietal Blend No. 2	0	10	75	30	6	2% by weight	70
9. Common No. 1	0	5	75	25	8	2% by weight	80
10. Common No. 2	5	10	100	50	10	3% by weight	70

1. Column V of Table VIII does not apply to sweet-clover seed.

TABLE IX

Applicable to:

- | | |
|-----------------------------------|---|
| (a) Clover, alsike | - <i>Trifolium hybridum</i> L. |
| (b) Clover, hop or yellow | - <i>Trifolium aureum</i> Poll. (= <i>T. agrarium</i> L.) |
| (c) Clover, hop, large | - <i>Trifolium campestre</i> Schreber (= <i>T. procumbens</i> L.) |
| (d) Clover, hop, small (suckling) | - <i>Trifolium dubium</i> Sibth. |
| (e) Clover, Persian | - <i>Trifolium resupinatum</i> L. |
| (f) Clover, strawberry | - <i>Trifolium fragiferum</i> L. |
| (g) Clover, white | - <i>Trifolium repens</i> L. |
| (h) Medick, black | - <i>Medicago lupulina</i> L. |
| (I) Timothy | |
| - common | - <i>Pbleum pratense</i> L. |
| - dwarf | - <i>Pbleum bertolonii</i> DC. (= <i>P. nodosum</i> auct.) |

I	II	III	IV	V	VI	VII	VIII	IX
	Maximum Number of Seeds per 25 g Except Where Otherwise Stated							
	Noxious Weeds			Other Crops				
GRADE NAME	PRIMARY	PRIMARY PLUS SECONDARY	TOTAL WEEDS	SWEET CLOVER	BRASSICA CROPS INCLUDING <i>S. alba</i>	OTHER NON-BRASSICA CROPS	MAXIMUM PERCENT ERGOT OR SCLEROTIA BODIES	MINIMUM PERCENT GERMINATION
1. Canada Foundation No. 1	0	0	8	0	0	5	1	80
2. Canada Foundation No. 2	0	2	40	1	2	30	2	70
3. Canada Registered No. 1	0	0	8	0	0	5	1	80
4. Canada Registered No. 2	0	2	40	1	2	30	2	70
5. Canada Certified No. 1	0	5	50	10	4	1% by weight	1	80
6. Canada Certified No. 2	0	10	75	25	6	2% by weight	2	70
7. Varietal Blend No. 1	0	5	50	10	4	1% by weight	1	80
8. Varietal Blend No. 2	0	10	75	25	6	2% by weight	2	70
9. Common No. 1	0	10	75	25	8	2% by weight	1	80
10. Common No. 2	5	20	125	50	10	3% by weight	2	70

TABLE X

Applicable to:

Trefoil, bird's-foot

- *Lotus corniculatus* L.

I	II	III	IV	V	VI	VII	VIII
	Maximum Number of Seeds per 25 g Except Where Otherwise Stated						
	Noxious Weeds			Other Crops			
GRADE NAME	PRIMARY	PRIMARY PLUS SECONDARY	TOTAL WEEDS	SWEET CLOVER	BRASSICA CROPS INCLUDING <i>S. ALBA</i>	OTHER NON-BRASSICA CROPS	MINIMUM PERCENT GERMINATION
1. Canada Foundation No. 1	0	1	10	1	0	50	75
2. Canada Foundation No. 2	0	2	75	1	2	300	65
3. Canada Registered No. 1	0	1	10	1	0	50	75
4. Canada Registered No. 2	0	2	75	1	2	300	65
5. Canada Certified No. 1	0	10	100	25	4	2% by weight	75
6. Canada Certified No. 2	0	20	200	50	6	5% by weight	65
7. Varietal Blend No. 1	0	10	100	25	4	2% by weight	75
8. Varietal Blend No. 2	0	20	200	50	6	5% by weight	65
9. Common No. 1	0	10	100	25	8	3% by weight	75
10. Common No. 2	5	20	200	100	10	5% by weight	65

TABLE XI

Applicable to:

(a) Bromegrass, meadow	- <i>Bromus riparius</i> Rehm.
(b) Bromegrass, smooth	- <i>Bromus inermis</i> Leysser
(c) Bromegrass, sweet	- <i>Bromus carinatus</i> Hook. & Arn.
(d) Canarygrass, reed	- <i>Phalaris arundinacea</i> L.
(e) Fescue, Cheving's	- <i>Festuca rubra</i> L. var. <i>commutata</i> Gaudin
(f) Fescue, fine-leaved	- <i>Festuca tenuifolia</i> Sibth. (= <i>F. ovina</i> L. var. <i>tenuifolia</i> (Sibth.) Dumort.) (= <i>F. capillata</i> Lam.)
(g) Fescue, hard	- <i>Festuca longifolia</i> Thuill. (= <i>F. ovina</i> L. var. <i>duriuscula</i> auct. amer.)
(h) Fescue, meadow	- <i>Festuca pratensis</i> Hudson
(i) Fescue, red and creeping red	- <i>Festuca rubra</i> L. var. <i>rubra</i>
(j) Fescue, sheep	- <i>Festuca ovina</i> L.
(k) Fescue, tall	- <i>Festuca arundinacea</i> Schreber
(l) Fescue, various-leaved	- <i>Festuca heterophylla</i> Lam.
(m) Foxtail, creeping	- <i>Alopecurus arundinaceus</i> Poirlet
(n) Foxtail, meadow	- <i>Alopecurus pratensis</i> L.
(o) Oatgrass, tall	- <i>Arrhenatherum elatius</i> (L.) Beauv. ex J. & K. Presl.
(p) Orchardgrass	- <i>Dactylis glomerata</i> L.
(q) Ryegrass, annual	- <i>Lolium multiflorum</i> Lam.
(r) Ryegrass, intermediate	- <i>Lolium × hybridum</i> Hausskn.
(s) Ryegrass, perennial	- <i>Lolium perenne</i> L.
(t) Wheatgrass, beardless	- <i>Agropyron spicatum</i> (Pursh) Scribn. & J.G. Smith f. <i>inermis</i> (Scribn. & J.G. Smith) Beetle
(u) Wheatgrass, crested	- <i>Agropyron cristatum</i> (L.) Gaertner or <i>Agropyron desertorum</i> (Fischer ex Link) Schult.
(v) Wheatgrass, intermediate	- <i>Agropyron intermedium</i> (Host) Beauv.
(w) Wheatgrass, northern	- <i>Agropyron dasystachyum</i> (Hook.) Scribn.
(x) Wheatgrass, pubescent	- <i>Agropyron trichophorum</i> (Link) Richter
(y) Wheatgrass, Siberian	- <i>Agropyron sibiricum</i> (Willd.) Beauv.
(z) Wheatgrass, slender	- <i>Agropyron trachycaulum</i> (Link) Malte ex H.F. Lewis
(aa) Wheatgrass, streambank	- <i>Agropyron riparium</i> Scribn. & Smith
(bb) Wheatgrass, tall	- <i>Agropyron elongatum</i> (Host) Beauv.
(cc) Wheatgrass, western	- <i>Agropyron smithii</i> Rydb.
(dd) Wildrye, Altai	- <i>Elymus angustus</i> Trin.
(ee) Wildrye, Dahurian	- <i>Elymus daburicus</i> Turcz ex Griseb.
(ff) Wildrye, Russian	- <i>Elymus junceus</i> Fischer

TABLE XI (cont'd)

I	II	III	IV	V	VI	VII	VIII	IX					XI	
	Maximum Number of Seeds per 25 g Except Where Otherwise Stated													
	Noxious Weeds				Other Crops									
GRADE NAME	PRIMARY	PRIMARY PLUS SECONDARY	TOTAL WEEDS	BRASSICA CROPS INCLUDING <i>S. Alba</i>	SWEET CLOVER	IN MEADOW FOXTAIL, RYEGRASS AND ORCHARDGRASS	IN OTHER KINDS	MAXIMUM PERCENT ERGOT BODIES	MINIMUM PERCENT PURE LIVING SEED				MINIMUM PERCENT PURE SEED BY WEIGHT	
									1	2	3	4	1	2
1. Canada Foundation No. 1	0	1	5	0	0	2	2	1	80	75	70	55	95	90
2. Canada Foundation No. 2	0	2	25	2	2	20	20	2	70	65	60	50	90	80
3. Canada Registered No. 1	0	1	5	0	0	2	2	1	80	75	70	55	95	90
4. Canada Registered No. 2	0	2	25	2	2	20	20	2	70	65	60	50	90	80
5. Canada Certified No. 1	0	4	50	4	5	2% by weight	1% by weight	1	80	75	70	55	95	90
6. Canada Certified No. 2	0	10	75	6	20	3% by weight	2% by weight	2	70	65	60	50	90	80
7. Varietal Blend No. 1	0	4	50	4	5	2% by weight	1% by weight	1	80	75	70	55	95	90
8. Varietal Blend No. 2	0	10	75	6	20	3% by weight	2% by weight	2	70	65	60	50	90	80
9. Common No. 1	0	5	75	8	25	3% by weight	2% by weight	1	80	75	70	55	95	90
10. Common No. 2	4	15	100	10	50	5% by weight	3% by weight	2	70	65	60	50	90	80

1. Notwithstanding the kinds and species of seeds set out in Table XI,

- a) subcolumn 1 of column X of Table XI applies only to Chewing's fescue, meadow fescue, red fescue, creeping red fescue, tall fescue, annual ryegrass, intermediate ryegrass and perennial ryegrass;
- b) subcolumn 2 of column X of Table XI applies only to smooth bromegrass, meadow bromegrass, sweet bromegrass, fine-leaved fescue, hard fescue, sheep fescue, various-leaved fescue, creeping foxtail, meadow foxtail, orchardgrass beardless wheatgrass, northern wheatgrass, pubescent wheatgrass, slender wheatgrass, streambank wheatgrass, Altai wildrye, Dahurian wildrye and Russian wildrye;
- c) subcolumn 3 of column X of Table XI applies only to crested wheatgrass, intermediate wheatgrass, Siberian wheatgrass, tall wheatgrass, western wheatgrass and tall oatgrass;
- d) subcolumn 4 of column X of Table XI applies only to reed canarygrass;
- e) subcolumn 1 of column XI of Table XI applies only to Chewing's fescue, meadow fescue, red fescue, creeping red fescue, tall fescue, annual ryegrass, intermediate ryegrass, perennial ryegrass and reed canarygrass; and
- f) subcolumn 2 of column XI of Table XI applies only to smooth bromegrass, meadow bromegrass, sweet bromegrass, fine-leaved fescue, hard fescue, sheep fescue, various-leaved fescue, creeping foxtail, meadow foxtail, tall oatgrass, orchardgrass, beardless wheatgrass, crested standard wheatgrass, crested fairway wheatgrass, intermediate wheatgrass, northern wheatgrass, pubescent wheatgrass, Siberian wheatgrass, slender wheatgrass, streambank wheatgrass, tall wheatgrass, western wheatgrass, Altai wildrye, Dahurian wildrye and Russian wildrye.

TABLE XII

Applicable to:

- (a) Alkaligrass, weeping - *Puccinellia distans* (L.) Parl.
- (b) Bentgrass, colonial(browntop) - *Agrostis capillaris* L. (= *A. tenuis* Sibth.)
- (c) Bentgrass, creeping - *Agrostis stolonifera* L. (= *A. palustris* Hudson)
- (d) Bentgrass, velvet - *Agrostis canina* L.
- (e) Bluegrass, annual - *Poa annua* L.
- (f) Bluegrass, Canada - *Poa compressa* L.
- (g) Bluegrass, fowl - *Poa palustris* L.
- (h) Bluegrass, Kentucky - *Poa pratensis* L.
- (I) Bluegrass, rough - *Poa trivialis* L.
- (j) Bluegrass, wood - *Poa nemoralis* L.
- (k) Crested dogtail - *Cynosurus cristatus* L.
- (l) Redtop - *Agrostis gigantea* Roth (= *A. alba* auct.)

I	II	III	IV	V	VI	VII	VIII	IX
	Maximum Number of Noxious Weed Seeds per 25 g		Maximum Percent by Weight				Minimum Percent Pure Living Seed	
GRADE NAME	PRIMARY	PRIMARY PLUS SECONDARY	TOTAL WEED SEEDS	OTHE R CROPS	MAXIMUM PERCENT ERGOT BODIES	MINIMUM PERCENT PURE SEED BY WEIGHT	KENTUCKY BLUEGRASS AND CANADA BLUEGRASS	OTHER KINDS
1. Canada Foundation No. 1	0	5	0.2	1	2	85	70	75
2. Canada Foundtion No. 2	1	20	0.5	2	3	75	60	65
3. Canada Registered No. 1	0	5	0.2	1	2	85	70	75
4. Canada Registered No. 2	1	2	0.5	2	3	75	60	65
5. Canada Certified No. 1	2	20	0.5	2	2	85	65	75
6. Canada Certified No. 2	4	50	0.8	4	3	75	55	65
7. Varietal Blend No. 1	2	20	0.5	2	2	85	65	75
8. Varietal Blend No. 2	4	50	0.8	4	3	75	55	65
9. Common No. 1	4	20	0.5	3	2	85	65	75
10. Common No. 2	8	100	1.0	5	3	75	55	65

TABLE XIII

Applicable to: Mixtures of forage seeds composed of seeds of two or more kinds listed in Tables VIII to XII, except mixtures of grass seeds designated by the seller as Lawn or Turf Grass Mixtures or Ground Cover Mixtures

I	II	III	IV	V	VI	VII	VIII	IX	X	XI	
	Maximum Number of Seeds per 25g except where otherwise stated										
	Noxious Weeds							Minimum Percent Germination			
GRADE NAME	PRIMARY	PRIMARY PLUS SECONDARY	TOTAL WEEDS	SWEET CLOVER	OTHER CROPS	BRASSICA CROPS INCLUDIN G <i>S. alba</i>	MAXIMUM PERCENT ERGOT BODIES	REED CANARYGRASS	KENTUCKY BLUEGRASS & CANADA BLUEGRASS	EACH OTHER COMPONENT	
1. Canada Certified No. 1 Forage Mixture	0	10	100	25	2% by weight	4	1	55	70	75	
2. Canada Certified No. 2 Forage Mixture	0	20	200	50	3% by weight	6	2	50	60	65	
3. Varietal Blend No. 1 Forage Mixture	0	10	100	25	2% by weight	4	1	55	70	75	
4. Varietal Blend No. 2 Forage Mixture	0	20	200	50	3% by weight	6	2	50	60	65	
5. Common No. 1 Forage Mixture	0	10	100	50	3% by weight	8	1	55	70	75	
6. Common No. 2 Forage Mixture	5	20	200	100	5% by weight	10	2	50	60	65	

1. Notwithstanding the kinds and species of seeds set out in Table XIII, column V of that Table does not apply to mixtures containing one percent or more of sweet-clover seed.
2. Notwithstanding Table XIII, Common No. 1 Forage Mixture may contain one primary noxious weed per 25 g where there is present, singly or combined, 10 percent or more of common timothy, Canada bluegrass, Kentucky bluegrass or redtop seed.

Note: A Varietal Blend Forage Mixture is a mixture of two or more kinds of seeds containing two or more varieties of a kind.

TABLE XIV - PART I

Applicable to: Lawn or Turf Grass Mixtures of Two or more of the kinds of seeds listed in Part II of this Table.

I	II	III	IV	V	VI	VII	VIII	IX	X
	Maximum Number of Noxious Weed Seeds per 25 g	Maximum Percent by Weight							
GRADE NAME	NOXIOUS WEEDS EXCLUDING COUCHGRASS	COUCHGRASS	TOTAL WEEDS	WHITE CLOVER	BROMEGRASS, ORCHARDGRASS & TALL FESCUE	OTHER KINDS	MAXIMUM PERCENT ERGOT BODIES	MINIMUM PERCENT PURE SEEDS BY WEIGHT	MINIMUM PERCENT GERMINATION OF EACH COMPONENT
1. Canada No. 1 Lawn Grass Mixture	20	10	0.5	5	1	2	2	85	70
2. Canada No. 2 Lawn Grass Mixture	75	20	1.0	5	2	3	3	75	60
4. Varietal Blend No. 1 Lawn Grass Mixture	20	10	0.5	5	1	2	2	85	70
5. Varietal Blend No. 2 Lawn Grass Mixture	75	20	1.0	5	2	3	3	75	60

Note: A Varietal Blend Lawn Grass Mixture is a mixture of two or more kinds of seeds containing two or more varieties of a kind.

TABLE XIV - PART II

I	II
1. Group A	Kentucky bluegrass, creeping red fescue, Chewing's fescue, hard fescue, colonial bentgrass, creeping bentgrass, velvet bentgrass, rough bluegrass, Canada bluegrass, sheep fescue, turf-type tall fescue varieties and turf-type perennial ryegrass varieties
2. Group B	Redtop, crested wheatgrass, timothy (dwarf-type) and annual ryegrass provided if none of those species constitutes more than 30 percent by weight singly and no more than 40 percent by weight combined of the total mixture
3. Group C	All other kinds of grass if those kinds are indicated on the label

Notwithstanding Table XIV, all mixtures described on a label to be suitable for shady places shall contain not less than 40 percent by weight, singly or combined, of Chewing's fescue, creeping red fescue or rough bluegrass seed.

TABLE XV

Applicable to: Ground Cover Mixtures composed of seed of two or more kinds other than Forage Mixtures and Lawn or Turf Grass Mixtures.

I	II	III	IV	V	VI
		Maximum Percent by Weight			
GRADE NAME	MAXIMUM NO. OF NOXIOUS WEED SEEDS PER 25G	TOTAL WEED SEEDS	OTHER KINDS	MINIMUM PERCENT PURE SEED BY WEIGHT	MINIMUM PERCENT GERMINATION OF EACH COMPONENT
1. Canada No. 1 Ground Cover Mixture	25	0.5	3	85	70
2. Canada No. 2 Ground Cover Mixture	75	1.0	5	75	60

TABLE XVI

Applicable to:

- (a) Beet - *Beta vulgaris* L.
- (b) Beet, sugar - *Beta vulgaris* L. *vulgaris* L. var. *saacbarifera* Lange
- (c) Mangel - *Beta vulgaris* L.
- (d) Swiss chard - *Beta vulgaris* L. var. *cicla* L.

I	II	III	IV	V	VI	VII
	Maximum Number of Seeds per 500 g					
	Noxious Weeds					
GRADE NAME	PRIMARY	PRIMARY PLUS SECONDARY	TOTAL WEEDS	OTHER CROPS	MAXIMUM NUMBER OF HEADS OF MAYWEED IN BEET AND SUGAR BEET PER 500g	MINIMUM PERCENT GERMINATION
1. Canada Foundation No. 1	0	2	10	5	1	75
2. Canada Foundation No. 2	2	10	30	20	1	65
3. Canada Registered No. 1	0	2	10	5	1	75
4. Canada Registered No. 2	2	10	30	20	1	65
5. Canada Certified No. 1	0	5	20	10	1	75
6. Canada Certified no. 2	2	20	50	20	1	65
7. Canada No. 1	1	10	20	10	1	75
8. Canada No. 2	2	30	100	30	1	65

TABLE XVII

Applicable to:

- (a) Cantaloupe - *Cucumis melo* L.
- (b) Citron - *Citrullus lanatus* (Thunb.) Matsum. & Nakai var. *citroides* L. (Bailey) Mansf.
- (c) Cucumber - *Cucumis sativus* L.
- (d) Gherkin - *Cucumis anguria* L.
- (e) Muskmelon - *Cucumis melo* L.
- (f) Pumpkin - *Cucurbita pepo* L. convar. *citrullinina* Greb.
- (g) Squash, crookneck - *Cucurbita moschata* Duchesne ex. Poirer
- (h) Squash, summer - *Cucurbita pepo* L. var. *meloepo* (L.) Alef.
- (i) Squash, winter or autumn - *Cucurbita maxima* Duch.
- (j) Vegetable marrow - *Cucurbita pepo* L. var. *medullosa* Alef.
- (k) Watermelon - *Citrullus lanatus* (Thunb.) Matsum. & Nakai var. *lanatus*

I	II	III	IV	V	VI
	Maximum Number of Seeds per 25g				
	Noxious Weeds				
GRADE NAME	PRIMARY	PRIMARY PLUS SECONDARY	TOTAL WEEDS	OTHER CROP	MINIMUM PERCENT GERMINATION
1. Canada Foundation No. 1	0	0	0	0	80
2. Canada Foundation No. 2	0	0	0	0	60
3. Canda Registered No. 1	0	0	0	0	80
4. Canada Registered No. 2	0	0	0	0	60
5. Canada Certified No. 1	0	1	1	1	80
6. Canada Certified No. 2	0	1	2	2	75
7. Canada No. 1	0	0	1	1	80
8. Canada No. 2	0	2	5	5	70

TABLE XVIII

Applicable to seed of the following species when designated by the seller as vegetable types for home gardens:

- (a) Bean, broad - *Vicia faba* L.
- (b) Bean, lima - *Phaseolus lunatus* L.
- (c) Bean, runner - *Phaseolus coccineus* L.
- (d) Chickpea - *Cicer arietinum* L.
- (e) Corn, pop - *Zea mays* L. var. *praecox* Bonaf. (= *Z. mays* L. var. *everta* (Sturter) Bailey)
- (f) Corn, sweet - *Zea mays* L. var. *rugosa* Bonaf. (= *Z. mays* L. var. *saccharata* (Struter) Bailey)
- (g) Pea - *Pisum sativum* L.
- (h) Soybean - *Glycine max* (L.) Merrill

I	II	III	IV	V	VI
	Maximum Number of Seeds per kg		Minimum Percent Germination		
GRADE NAME	WEEDS	OTHER CROPS	PEA	BEAN AND SOYBEAN	SWEET CORN
1. Canada Foundation No. 1	0	0.0	80	85	90
2. Canada Foundation No. 2	0	0.5	70	75	80
3. Canada Registered No. 1	0	0.0	80	85	90
4. Canada Registered No. 2	0	1.0	70	75	80
5. Canada Certified No. 1	0	1.0	80	85	90
6. Canada Certified No. 2	0	3.0	70	75	80
7. Canada No. 1	1	4.0	75	80	90
8. Canada No. 2	3	10.0	65	65	75

TABLE XIX

Applicable to:

- (a) Broccoli - *Brassica oleracea* L. convar. *botrytis* (L.) Alef. var. *italica* Plenck
- (b) Brussels sprouts - *Brassica oleracea* L. var. *bullata* DC. subvar. *gemmifera* DC.
- (c) Cabbage - *Brassica oleracea* L. var. *capitata* L.
- (d) Cauliflower - *Brassica oleracea* L. Alef. var. *botrytis* L.
- (e) Collard - *Brassica oleracea* L. var. *acephala* DC.
- (f) Kale, vegetable - *Brassica oleracea* L. var. *acephala* DC.
- (g) Kohlrabi - *Brassica oleracea* L. var. *gongylodes* L.
- (h) Radish - *Rapbanus sativus* L.
- (l) Rape, forage - *Brassica napus* L., *B. campestris* L.
- (j) Rutabaga (swede) - *Brassica napus* L. var. *napobrassica* (L.) Reichb.
- (k) Spinach mustard - *Brassica perviridis* Bailey, and
- (l) Turnip - *Brassica rapa* L. var. *rapa* (L.) Thell.

I	II	III	IV	V	VI	VII
	Maximum Number of Seeds per 25g					
	Noxious Weeds					
	Primary					
GRADE NAME	IN BROCCOLI, BRUSSEL SPROUTS, CABBAGE AND CAULIFLOWER	IN OTHER KINDS	PRIMARY PLUS SECONDARY	TOTAL WEEDS	OTHER CROPS	MINIMUM PERCENT GERMINATION
1. Canada Foundation No. 1	0	0	2	5	5	80
2. Canada Foundation No. 2	2	2	10	25	20	60
3. Canada Registered No. 1	0	0	2	5	5	80
4. Canada Registered No. 2	2	2	10	25	20	60
5. Canada Certified No. 1	0	0	5	20	10	80
6. Canada Certified No. 2	2	2	20	50	20	70
7. Canada No. 1	3	1	10	30	10	80
8. Canada No. 2	5	2	50	100	30	70

TABLE XX

Applicable to:

Minimum Percent Germination

(a)	Artichoke	- <i>Cynara scolymus</i> L.	75
(b)	Asparagus	- <i>Asparagus officinalis</i> L. ssp. <i>officinalis</i>	75
(c)	Cardoon	- <i>Cynara cardunculus</i> L.	55
(d)	Carrot (cultivated)	- <i>Daucus carota</i> L. ssp. <i>sativa</i> (Hoffm.) Arcangeli	60
(e)	Celeriac	- <i>Apium graveolens</i> L. var. <i>rapaceum</i> (Miller) Gaudich.	60
(f)	Celery	- <i>Apium graveolens</i> L. var. <i>dulce</i> (Miller) Pers.	55
(g)	Celtus	- <i>Lactuca sativa</i> L. var. <i>asperangia</i> Bailey	75
(h)	Chervil	- <i>Chaerophyllum bulbosum</i> L.	75
(I)	Chicory (cultivated)	- <i>Cichorium intybus</i> L.	65
(j)	Chive	- <i>Allium schoenoprasum</i> L.	65
(k)	Cornsalad	- <i>Valerianella olitoria</i> (L.) Poll. (= <i>V. locusta</i> Betsche)	75
(l)	Cress, garden	- <i>Lepidium sativum</i> L.	75
(m)	Cress, water	- <i>Nasturtium officinale</i> R. Brown	35
(n)	Dandelion (cultivated)	- <i>Taraxacum officinale</i> Wiggers	55
(o)	Dill	- <i>Anethum graveolens</i> L.	50
(p)	Eggplant	- <i>Solanum melongena</i> L.	65
(q)	Endive	- <i>Cichorium endivia</i> L.	65
(r)	Leek	- <i>Allium porrum</i> L.	65
(s)	Lettuce:		
	- cos (romaine)	- <i>Lactuca sativa</i> L. var. <i>longifolia</i> Lam.	70
	- head	- <i>Lactuca sativa</i> L. var. <i>capitata</i> L.	70
	- leaf	- <i>Lactuca sativa</i> L. var. <i>crispa</i> L.	70
(t)	Okra	- <i>Hibiscus esculentus</i> L.	55
(u)	Onion	- <i>Allium cepa</i> L.	75
(v)	Parsley	- <i>Petroselinum crispum</i> (Miller) Nym. ex Hill	55
(w)	Parsnip	- <i>Pastinaca sativa</i> L.	60
(x)	Pepper	- <i>Capsicum annuum</i> L.	65
(y)	Rampion	- <i>Campanula rapunculus</i> L.	55
(z)	Rhubarb	- <i>Rheum rhabarbaricum</i> L.	65
(aa)	Sage	- <i>Salvia officinalis</i> L.	50
(bb)	Salsify	- <i>Tragopogon porrifolius</i> L.	75
(cc)	Savory	- <i>Satureja hortensis</i> L.	50
(dd)	Sorrel (cultivated)	- <i>Rumex acetosa</i> L.	65
(ee)	Spinach	- <i>Spinacia oleracea</i> L.	65
(ff)	Spinach, New Zealand	- <i>Tetragonia tetragonoides</i> (Pall.) Kuntze (= <i>T. expansa</i> Thunb. ex Murray)	50
(gg)	Thyme	- <i>Thymus vulgaris</i> L.	50
(hh)	Tobacco (fine-cured and burley types)	- <i>Nicotiana tabacum</i> L.	75
(ii)	Tomato	- <i>Lycopersicon lycopersicum</i> (L.) Karsten ex Farwell (= <i>L. esculentum</i> Miller)	75

TABLE XX CONT'D...

I	II	III	IV	V
	Maximum Number of Seeds per 25g			
	Noxious Weeds			
GRADE NAME	PRIMARY	PRIMARY PLUS SECONDARY	TOTAL WEEDS	OTHER CROPS
1. Canada Foundation No. 1	0	3	10	5
2. Canada Foundation no. 2	2	10	25	20
3. Canada Registered No. 1	0	3	10	5
4. Canada Registered No. 2	2	10	25	20
5. Canada Certified No. 1	0	5	20	10
6. Canada Certified No. 2	2	20	50	20
7. Canada No. 1	1	10	25	10
8. Canada No. 2	2	50	100	30

TABLE XXI

Applicable to:

Onion Sets and multiplier onions

I	II	III	IV
GRADE NAME	DIAMETER OF SETS (CM)	PERCENT ONE VARIETY AND COLOUR	GENERAL QUALITY
1. Canada No. 1	1 to 2.5	Not less than 98	<ol style="list-style-type: none"> 1. Mature, well cured, sound and dry. 2. Practically free from tops, dirt, leaves or other foreign matter, from disease, decay and moulds and from insect, mechanical, frost or other damage
2. Canada No. 1 (Large)	2.5 to 3.5		<ol style="list-style-type: none"> 3. Practically free from sprouted and soft bulbs when graded.
3. Canada No. 2	1 to 2.5	Not less than 90	<ol style="list-style-type: none"> 1. Mature, well cured, sound and reasonably dry. 2. Reasonably free from tops, dirt, leaves or other foreign matter, from disease, decay and moulds and from insect, mechanical, frost or other damage.
4. Canada No. 2 (Large)	2.5 to 3.5		<ol style="list-style-type: none"> 3. Reasonably free from soft bulbs and containing not more than 5% sprouts for yellow and 10% for white, red and globe.

TABLE XXI CONT'D...

DEFINITION OF TERMS USED IN COLUMN IV

“Mature, well cured” means that the sets have neck that are well dried out and have not more than 1% and 2% of thick necks in Canada No. 1 and Canada No. 2 sets, respectively.

“Sound” means that the set is firm and neat and will yield only slightly to moderate pressure.

“Practically free from tops, dirt, leaves or other foreign matter” means free for all intents and purposes. Tops, dirt, leaves or other foreign matter should not exceed 1 per cent by weight and not more than 25 per cent of the sets should have tops 6.5 cm long or longer.

“Practically free from disease, decay and moulds and from insect, mechanical, frost or other damage” means that not more than 3 per cent combined of the bulbs should be so affected.

“Practically free from sprouted bulbs” means that, for yellow sets, not over 3 per cent have visible sprouts and, for red, white and globe sets, that not over 6 per cent have visible sprouts

“Practically free from soft bulbs” means that such bulbs should not be in excess of 3 per cent.

“Reasonable free from tops, dirt, leaves or other foreign matter” means that not more than 35 per cent of the bulbs should have tops over 6.5 cm long and dirt, leaves or other foreign matter should not exceed 2 per cent by weight.

“Reasonably free from disease, decay and moulds and from insect, mechanical, frost or other damage” means that not more than 4 per cent combined of the bulbs should be so affected.

“Reasonably free from soft bulbs” means that not more than 5 per cent combined of the bulbs should be so affected.

“Sprouted bulbs” means those bulbs with sprouts exceeding 0.6 cm in length.

“Sound, reasonably dry and reasonably free from tops, dirt, leaves or other foreign matter, from decay, disease and moulds and from insect, mechanical, frost or other damage” means that such conditions and defects may be apparent on examination, but should not be present to an extent that will seriously affect the keeping quality under average storage conditions.

1. The diameter of sets set out in column II of Table XXI to Schedule I do not apply to multiplier onions.
2. For the purposes of Table XXI to Schedule I, a tolerance of three per cent by weight is allowed for the oversized or undersized sets for each grade of onion sets.

TABLE XXII

Applicable to all of the kinds of seeds referred to in Tables I to XX except Tables III, XIII, XIV and XV.

1. Canada Foundation Substandard seed is seed of Foundation status that meets the standards for purity for Canada Foundation No. 2 grade seed.
2. Canada Registered Substandard seed is seed of Registered status that meets the standards for purity for Canada Registered No. 2 grade seed.
3. Canada Certified Substandard seed is seed of Certified status that meets the standards for purity for Canada Certified No. 2 grade seed.
4. Canada Foundation Substandard (Purity) seed is seed of Foundation status that meets the standards for purity for common grade seed.