

Index

A

- Active labour
 - care required, 5.11
 - diagnosis, 5.11
 - initial assessment, 5.11–12
- Active management of labour, 5.26
 - components, 5.26
 - psychosocial support, 5.26
- Adoption, 8.9
- Allied health care
 - Level II
 - neonatal care, 2.24
 - postpartum, maternal, and newborn care, 2.23
- ALPHA Form, 4.46–48 (App.3)
 - preconception care, 3.11
 - prenatal care, 4.12, 4.13 (t4.3)
- Ambulatory prenatal care, 2.13
 - Level I, 2.14
 - Level II, 2.14
 - Level III, 2.14
- Amniotomy
 - early, 5.26
 - effects, 5.26
- Ancillary services, maternal, and newborn care, 2.28
- Antenatal Psychosocial Health Assessment, *see* ALPHA Form
- Augmentation of labour, 5.27, 5.47–48 (App.2)

B

- Baby-friendly hospital initiative, 7.36–40
- BFHI, *see* baby-friendly hospital initiative, 7.36
- Birth facility, design of, 5.6
- Birth plans, 5.7, 44
 - advantages, 5.7–8
 - example, 5.44–46 (App.1)
- BMI, *see* Prenatal care, Body Mass Index
- Breast assessment, 7.11–12
- Breast augmentation surgery, 7.12
- Breast feeding, after birth, 5.25

- Breast milk jaundice syndrome, 7.21
- Breast milk storage, 7.48–49 (App.8)
- Breast pumps, 7.47–48 (App.8)
- Breast reduction surgery, 7.12
- Breastfeeding
 - breast assessment, 7.11–12
 - contraindication, 7.12
 - early postnatal period, 7.13
 - eliminating obstacles, 7.23
 - facts, 7.7 (t7.1)
 - follow-up support, 7.29
 - hospital discharge and follow-up, 7.19
 - informed choice, 7.8
 - late postnatal period, 7.25
 - length of hospital stay, effect, 7.19
 - literature available, 7.6
 - loss of milk supply, 7.26
 - maternal medication, 7.24–25
 - medical reasons for supplementation, 7.22
 - mother's nutrition, 7.17
 - multiple babies, 7.23
 - myths, 7.9–10
 - nutrition myths, 7.18
 - pointers for health care providers, 7.13–14
 - policies, 7.8
 - positioning and latch, 7.14
 - potential difficulties, 7.21
 - premature infants, 7.23
 - reasons for stopping, 7.25
 - returning to paid employment, 7.27–28
 - role of health care providers, 7.9
 - signs of adequate nutrition for baby, 7.15
 - signs to seek help, 7.16–17
 - special situations, 7.22
 - supplementation, 7.22
 - vitamin and mineral supplementation, 7.26
 - weaning process, 7.28
 - workplace concerns, 7.26–29
- Breastfeeding clinics, 6.34

C

- Calcium intake, preconception care, 3.18
- Canada's Food Guide to Healthy Eating*, 4.20
- Canadian Perinatal Surveillance System, 2.31, 2.39–40 (App.1)
- Cardiopulmonary resuscitation, 6.10
- Cesarean birth, postpartum care, 6.27
- Cesarean births, 5.29, 10.10
 - facilitating family-centred care, 5.30–31 (t5.1)
 - rates, 5.29
 - reducing rate, 5.29
- Checklist for the care of women
 - experiencing early pregnancy loss, 8.30 (App.1)
- Child with a congenital anomaly, 8.9
- Childbirth care, 2.10
- Circumcision, 6.17
- Combined mother/baby care, 6.12–13
 - role of health care providers, 6.12–13
- Combined spinal epidural analgesia, 5.21–22
- Comfort measures during labour, 5.19–20
- Community participation, 1.16 (t1.5)
- Community programs, preconception care, 3.9
- Community support, perinatal loss, 8.23–24
- Community-based care, 2.9
- Continuous quality improvement, 5.49–50 (App.3)
 - effect on cesarean birth, 5.29
- CPR, *see* Cardiopulmonary resuscitation
- CPSS, *see* Canadian Perinatal Surveillance System
- CSE, *see* Combined spinal epidural analgesia

D

- Diversity, sensitivity to, 3.6
- Drug information, preconception care, 3.16
- Drug use, preconception care, 3.15
- Dyad care, *see* Combined mother/baby care

Dystocia

- definition, 5.25–26
- treatment, 5.27

E

- Early labour, diagnosis, 5.10–11
- Educating health care providers, perinatal loss, 8.21–22
- Education
 - adult learning principles, 2.32 (t2.5)
 - health professionals, 2.33–34
 - learning opportunities for health professionals, 2.34
 - principles for educational materials, 2.33 (t2.6)
 - women and families, 2.31–33
- Electronic fetal heart-rate monitoring, 5.17
 - Society of Obstetricians and Gynaecologists of Canada recommendations, 5.17
- Environmental hazards, preconception care, 3.16
- Epidural analgesia, 5.21
 - information on, 5.22
 - negative effects, 5.21
 - variations, 5.21
- Episiotomy
 - effects of, 5.22
 - recommendation to discontinue, 5.22–23
- Equipment for labour, birth, recovery, and postpartum room, 10.30 (App.2)
- Equipment for normal newborn nurseries, 10.31 (App.2)
- Equipment, neonatal transport, 9.21 (App.2)
- Erythromycin treatment, 6.11
- Evaluation
 - questions to consider, 2.36
 - treatments, technology and policies, 2.35–36
- Evaluation of care, 2.34–35
- Evidence-based practice, 1.6
 - limitations, 1.7
- Expressing and storing expressed breast milk, 7.46–49 (App.8)

F

- Facilities, design process, 10.28 (App.1)
- Facilities
- acoustics, 10.23
 - antenatal, birth, and postpartum, 10.8–10
 - architectural process, 10.28 (App.1)
 - communication, 10.23
 - electrical service, 10.23
 - environmental controls and engineering, 10.22–24
 - equipment requirements, 10.30–31 (App.2)
 - family-centred intensive care, evaluation, 10.17 (t10.1)
 - guidelines for environmental light, 10.18 (t10.2)
 - guidelines for environmental sound, 10.19 (t10.3)
 - guidelines for infant-sensitive developmental care, 10.19 (t10.4)
 - infant resuscitation area, 10.12–13
 - labour and birth unit, key components, 10.10–12
 - labour/birth/recovery, 10.21–22
 - labour/birth/recovery room, requirements, 10.10–12
 - labour/birth/recovery/postpartum, 10.21–22
 - Level II, 10.16
 - Level III, 10.16
 - requirements for space and equipment, 10.21
 - Level III nurseries, requirements for space and equipment, 10.20
 - neonatal intensive care unit, 10.16–20
 - nursery, 10.14–15
 - operative birth room, 10.12
 - planning principles, 10.5–7
 - planning process, 10.27–29 (App.1)
 - postpartum mother/baby rooms, 10.13–14
 - selection of materials and finishes, 10.24
 - single-room approach, 10.9
 - special care nursery, 10.17
 - support areas needed for antepartum, birth, and postpartum facilities, 10.32 (App.3)
 - terminology, 10.27–28 (App.1)
 - thermal environment, ventilation, and air conditioning, 10.22–23
 - use of existing, 10.7–8
- Family participation, 1.15
 - on committees, 1.16 (t1.4)
 - during labour and birth, 5.10
 - enabling, 1.15 (t1.3)
- Family support, perinatal loss, 8.13–14
- Family-centred cesarean birth, facilitating, 5.30–31 (t5.1)
- Family-centred maternity and newborn care
 - cesarean births, 5.30
 - definition, 1.8
 - guiding principles, 1.8–12, 2.5, 3.5, 4.5, 5.5, 7.6, 9.6, 10.5–6
 - implementing, 1.14 (t1.1)
 - informed choice, 1.10
 - mission statement, 2.11
 - policies and programs checklist, 2.12 (t2.2)
 - program administration, 2.11
 - research evidence, 1.9
 - role of health care providers, 1.11
 - use of technology, 1.11
- FAS, *see* Fetal alcohol syndrome
- FCMC, *see* Family-centred maternity and newborn care
- Female genital mutilation, and preconception counselling, 3.12
- Fetal alcohol syndrome, preconception care, 3.14
- Fetal health surveillance, 5.16–18
 - appropriate terminology, 5.17–18
- Fetal scalp blood sampling, 5.17
- FGM, *see* Female genital mutilation
- Fluid balance, newborn, 6.24
- Folic acid intake, preconception care, 3.19–20

G

- Genetic counselling, 3.24
- Grief support
 - the five Cs, 8.12
 - interventions, 8.12 (t8.5)
- Grieving tasks, 8.10

- Guidelines
 content, 1.7
 history and context, 1.5–6
 implementation, 1.12–16
 purpose, 1.6
 Guidelines for taking a feeding history, 7.44 (App.6)
 Guiding principles, family-centred maternity and newborn care, 1.8–12, 2.5, 3.5, 4.5, 5.5, 7.6, 9.6, 10.5–6
- H**
- Health care providers
 continuity of care, 5.9–10
 guidelines for transfer of care, 5.8
 role during labour and birth, 5.8–9
 training, 1.14
 Health professionals, education, 2.33
 Health records, 2.30–31
Healthy Beginnings: Guidelines for Care During Pregnancy and Childbirth, 4.9
 Home births, 2.10
 Hospital admission, 5.12
 procedures no longer recommended, 5.12
 Hospital discharge, guidelines, 7.19–20
 Hospital stay, guidelines, 6.36
 Hospital/community referral, 6.37
 Hospital/community referral form, 6.37 (t.6.3)
 Hospital/Maternity facility guidelines for the implementation of WHO/UNICEF baby-friendly hospital initiative, 7.36–40 (App.3)
 How can I tell if my baby is breast feeding well?, 7.45 (App.7)
 Hyperbilirubinemia, 6.24–25
 investigation, 6.25 (t.6.2)
 Hypoglycemia, newborn, 6.23–24
- I**
- ICD-9 and 10 coding interpretation and diagnosis of ALPHA, 2.41 (App.2)
 Immunization of newborn, 6.18
 Indications for seeking help, 6.45 (App.3)
 Induction and augmentation methods, 5.47–48 (App.2)
 Infant nutrition resources: assessment guide, 7.42–43 (App.5)
 Initial assessment, sources of information, 5.12
 Interdepartmental Perinatal Loss Flowsheet, 8.31–33 (App.2)
 International Code of Marketing of Breast Milk Substitutes, 7.35 (App.2)
 Intracutaneous sterile water, use of, 5.20
 Intravenous patient-controlled analgesia, 5.20
 Iron intake, preconception care, 3.19
 IV PCA, *see* Intravenous patient-controlled analgesia
- J**
- Jaundice, 6.24–25
 neonatal, 7.21
Joint Statement: Prevention of Fetal Alcohol Syndrome (FAS) and Fetal Alcohol Effects (FAE) in Canada, 3.14, 4.18
- L**
- La Leche League, 6.34
 Labour
 assessing progress, 5.15
 nutrition and hydration, 5.15
 position and ambulation, 5.15–16
 supportive care, 5.13
 Labour, birth, and immediate postpartum care, Level I personnel, 2.20–21
 Labour, birth, postpartum, and newborn care
 Level I, 2.14–15
 Level II, 2.16–17
 Level III, 2.17–18
 Labour, birth, postpartum, and newborn care, personnel requirements, 2.18–27
 Labour induction and cesarean birth, 5.28
 effects of, 5.27–29

- morbidity risk, 5.28
 - perinatal mortality, 5.28
 - Labour/birth/recovery room, 5.7, 10.9–10
 - Labour/birth/recovery/postpartum room, requirements, 10.10–12
 - Labour/birth/recovery/postpartum system, 10.9
 - Language, importance of, 1.12
 - Late postpartum hemorrhage, 6.22
 - LBR, *see* Labour/birth/recovery room
 - LBRP
 - see* Labour/birth/recovery/postpartum room
 - see* Labour/birth/recovery/postpartum system
 - Level I
 - postpartum, maternal, and newborn care, 2.22
 - minimal staffing requirements, 2.21
 - physician and midwifery care, 2.22
 - postpartum maternal/newborn care, 2.21–22
 - registered nurse responsibilities, 2.21–22
 - Level I hospital, 5.5
 - Level I personnel, 2.20–21
 - labour, birth, and immediate postpartum care, 2.20–21
 - physician and midwifery care, 2.20–21
 - registered nurse responsibilities, 2.20 (t8.3)
 - Level I personnel, nursing care, 2.20–21
 - Level II
 - labour, birth, and immediate postpartum care, 2.22
 - physician and midwifery care, 2.22–23
 - registered nurse responsibilities, 2.22
 - neonatal care, 2.23–24
 - allied health care, 2.24
 - nursing care, 2.24
 - physician care, 2.24
 - postpartum, maternal, and newborn care, minimal staffing requirements, 2.23
 - postpartum, maternal, and newborn care, 2.23
 - allied health care, 2.23
 - physician and midwifery care, 2.22–23
 - registered nurse responsibilities, 2.23
 - Level III
 - labour, birth, and immediate postpartum care, 2.24
 - nursing care, 2.24–25
 - physician and midwifery care, 2.25
 - registered nurse responsibilities, 2.24
 - neonatal care, 2.25–27
 - allied health care, 2.27
 - minimal staffing requirements, 2.25
 - nursing care, 2.25–26
 - physician care, 2.26–27
 - staff:infant ratios, 2.26
 - postpartum, maternal, and newborn care, 2.25
 - registered nurse responsibilities, 2.25
 - Loss and grief assessment, 8.11 (t8.4)
- M**
- Maternal and newborn care
 - ancillary services, 2.28
 - phases, 2.6
 - Maternal and newborn outreach programs, 2.10
 - Maternal and newborn support outcomes, 6.31
 - Maternal transport equipment, 9.21 (App.2)
 - Medication, and breastfeeding, 7.24–25
 - Minimal staffing requirements
 - Level I, postpartum, maternal, and newborn care, 2.21
 - Level II, neonatal care, 2.23
 - Level III, neonatal care, 2.25
 - Miscarriage, 8.8
 - Mother-baby contact, 7.14
 - benefits, 5.24–25
 - Mother-friendly workplace, 7.27

N

- Narcotics use, 5.20–22
 - drawbacks, 5.20
- NEC, *see* Necrotizing enterocolitis
- Necrotizing enterocolitis, 7.22
- Neonatal jaundice, 7.21
- Neonatal Resuscitation Program, 2.19
- Neural tube defects, 3.19–20
- Newborn, potential problems, 6.44 (App.2)
- Newborn cardiorespiratory distress, 6.23
- Newborn resuscitation, 2.19–20, 6.10–11
- NICU, *see* Facilities, neonatal intensive care unit
- NICU bereavement checklist, 8.35 (App.4)
- Nitrous oxide, 5.21
 - for labour analgesia, 5.21
- NTD, *see* Neural tube defects
- Nursing care
 - during labour and birth, 5.9
 - knowledge requirements, 5.14
 - Level I, 2.20
 - postpartum, maternal, and newborn care, 2.21–22
 - Level II
 - labour, birth, and immediate postpartum care, 2.22
 - neonatal care, 2.24
 - postpartum, maternal, and newborn care, 2.23
 - Level III
 - labour, birth, and immediate postpartum care, 2.24–25
 - neonatal care, 2.25–26
 - postpartum, maternal, and newborn care, 2.25
 - supportive role, 5.14
- Nutrition
 - during breastfeeding, 7.17
 - folic acid intake, 3.19
 - iron intake, 3.19
 - neural tube defects, 3.19–20
 - vitamin A, excess intake, 3.20
 - vitamin D intake, 3.18–19

- Nutrition counselling, preconception care, 3.17–18

Nutrition for a Healthy Pregnancy: National Guidelines for the Child-bearing Years, 3.21, 4.20

O

- Ophthalmia neonatorum, 6.11
- Oxytocic drugs use, 5.25
- Oxytocin, early administration, 5.26
- Oxytocin administration, 5.27

P

- Pain during labour, 5.18–22
 - comfort measures, 5.19–20
 - pharmacological methods for control, 5.20
 - relief of, 5.18
- Parent support groups, perinatal loss, 8.24–25
- Parent-infant contact
 - benefits, 6.6
 - postpartum period, 6.6
- Parenting classes, 6.34
- Patient-controlled epidural analgesia, 5.21–22
- PCEA, *see* patient-controlled epidural analgesia
- Perinatal loss
 - assessment of family needs, 8.9
 - bereavement guides, 8.18
 - cautions to health care providers, 8.18
 - communicating with other children, 8.14
 - community support, 8.23–24
 - cremation, burial, and funeral options, 8.15
 - educating health care providers, 8.21–22
 - elements of grief and loss assessment, 8.11 (t8.4)
 - facilitating communication, 8.14–15
 - facilitating and supporting grieving, 8.12
 - family self-help and mutual aid, 8.25–26

- family support, 8.13–14
- follow-up communication, 8.36 (App.5)
- grieving, religious, and cultural consideration, 8.19
- legal considerations, 8.19–21
 - burial and cremation, 8.20
 - postmortems, 8.19
 - returning to work, 8.21
- miscarriage, 8.16–17
 - D+C medical, 8.17
 - D+C surgical, 8.17
- NICU Bereavement Checklist, 8.35 (App.4)
- parent support groups, 8.24–25
- postmortem checklist, 8.34 (App.3)
- religious and cultural considerations, 8.19
- remembrance of baby, 8.16 (t8.6)
- resources for families, 8.28
- rights of the infant, 8.7 (t8.1)
- rights of the parent, 8.7 (t8.2)
- role of health care providers, 8.6
- self-esteem of parents, 8.9
- self-help and mutual aid
 - role of health care providers, 8.26
 - strategies, 8.27–28
- special considerations for babies with congenital anomalies, 8.17–18
- special considerations for premature babies, 8.17–18
- support groups, 8.27
- support for health care providers, 8.22
- transportation of baby, 8.20–21
- types of, 8.8
- Perinatal Loss Review Team, 8.6
- Perinatal loss, grieving process, 8.10
- Physical abuse, determination of, 4.17
- Physical abuse, 4.17 (t4.4)
- Physical activity
 - preconception care, 3.16–17
 - prenatal care, 4.19
- Physical Activity Guide to Healthy Active Living*, 3.21
- Physician care
 - Level II, neonatal care, 2.24
 - Level III, neonatal care, 2.26–27
- Physician and midwifery care
 - Level I, postpartum, maternal, and newborn care, 2.22
 - Level I personnel, 2.20–21
 - Level II, postpartum, maternal, and newborn care, 2.23
 - Level III, labour, birth, and immediate postpartum care, 2.25
- Place of birth, 5.6–7
- Policies and procedures, 2.29–30
 - written policies, 2.30 (t2.4)
- Population diversity, 1.6
- Portograms, use, 5.26
- Post-term pregnancy, 5.28
- Postpartum, planning for, 6.28–29
- Postpartum blues, 6.19–20
- Postpartum care
 - after cesarean birth, 6.27
 - assessment and care of mother, 6.7–8
 - assessment and care of newborn, 6.8–9, 16
 - assessment of mother, criteria, 6.8
 - assessment of newborn complications, 6.23–26
 - breastfeeding clinics, 6.34
 - circumcision, 6.17
 - clinics, 6.33
 - combined mother/baby, 6.12–13
 - community maps, 6.34
 - community programs, 6.36
 - community support, 6.28–29
 - community support strategies, 6.32–35
 - depression, 6.20–21
 - electronic information, 6.34
 - erythromycin treatment, 6.11
 - goals, 6.5
 - hospital/community referral, 6.37
 - immunization of newborn, 6.18
 - importance, 6.13
 - indicators of adjustment, 6.29–30
 - indicators of mother's adjustment, 6.29–30
 - indicators of newborn safety, 6.30–31
 - jaundice, 6.24–25
- La Leche League, 6.34

- late postpartum hemorrhage, 6.22
- learning needs of mother and baby, 6.14
- length of hospital stay, 6.35–36
- maternal and newborn support outcomes, 6.31
- needs and services, 6.5
- newborn cardiorespiratory distress, 6.23
- newborn dehydration, 6.26
- newborn fluid balance, 6.24
- newborn hearing loss, 6.18
- newborn hypoglycemia, 6.23–24
- newborn resuscitation, 6.10–11
- newborn risk assessment, 6.9–10
- newborn screening, 6.18–19
- newborn sepsis risk, 6.26
- newborns at risk, 6.19
- newborns with special needs, 6.38–39
- outcomes, 6.29–30
- parenting classes, 6.34
- phone follow-up, 6.33
- phone lines, 6.32–33
- physician follow-up, 6.33–34
- postpartum blues, 6.19–20
- principles for facilities, 6.15 (t6.1)
- professional home visits, 6.33
- puerperal infection/endometritis, 6.22
- psychosis, 6.21–22
- risk assessment, 6.10
- role of health care providers, 6.14–15
- skin and cord care of newborn, 6.16–17
- specific intervention, 6.11
- substance use, effect on newborn, 6.26–27
- temperature regulation, 6.9
- transition to home and community, 6.28–29
- treatment of depression, 6.20–21
- vitamin K administration, 6.11
- Postpartum period
 - phases, 6.14
 - three phases, 6.44 (App.2)
- Postpartum psychosis, 6.21–22
 - treatment, 6.21–22
- Preconception care
 - alcohol use, 3.14
 - ALPHA Form, 3.11
 - calcium intake, 3.18
 - Canada's Food Guide to Healthy Eating*, 3.18–19
 - components, 3.5
 - counseling, 3.21, 3.23 (t3.2)
 - dealing with abuse, 3.11–12
 - dealing with female genital mutilation, 3.12
 - dealing with stress, 3.10
 - drug information, 3.16
 - drug use, 3.15
 - environmental hazards, 3.16
 - fetal alcohol syndrome, 3.14
 - folic acid intake, 3.19–20
 - genetic counseling, 3.24
 - iron intake, 3.19
 - lifestyle counseling, 3.12–13
 - location, 3.7–8
 - nutrition counseling, 3.17–18
 - physical activity, 3.16–17
 - role of community programs, 3.9
 - role of health care providers, 3.6
 - role of primary care practitioners, 3.9
 - social support, 3.10
 - tobacco use, 3.13
- Preconception counselling, vitamin D intake, 3.18–19
- Preconception history taking, 3.22–24 (t3.2)
- Preconception programs, role of media, 3.8–9
- Pregnancy discomforts, 4.52–54 (App.5)
- Premature infants, breastfeeding, 7.23
- Prenatal care
 - alcohol use, 4.18
 - role of health care provider, 4.18
 - ALPHA Form, 4.12
 - ALPHA guide, 4.14
 - Body Mass Index, 4.21
 - complications of pregnancy, 4.23–24
 - continuing assessment, 4.23
 - continuity of care, 4.7
 - counseling, 4.12

- couple relationships, 4.15
 - determination of risk, 4.9
 - discomforts of pregnancy, 4.23, 4.52–54 (App.5)
 - drug use, 4.19
 - resources for health care providers, 4.19
 - elements of prenatal history, 4.10 (t4.2)
 - environmental exposure, 4.20
 - goals, 4.5
 - groups nutritionally at risk, 4.22
 - history taking, 4.9
 - initial care visits, 4.8
 - nutrition, 4.20
 - nutritional concerns, 4.21 (t4.6), 4.49–51 (App.4)
 - organization and provision, 4.5
 - personal support, 4.7
 - physical activity, 4.19
 - physical examination, 4.9
 - pregnancy, birth, and postpartum planning, 4.22
 - preterm labour, 4.24
 - psychosocial assessment, 4.12
 - psychosocial stress, 4.14
 - records, 4.8–9
 - records example, 4.38 (App.1)
 - role of health care providers, 4.6
 - alcohol use, 4.18
 - screening, 4.11
 - screening procedures, 4.40–46 (App.2)
 - sexuality, 4.15–16
 - signs of preterm labour, 4.24 (t4.7)
 - Society of Obstetricians and Gynaecologists of Canada, 4.11
 - Society of Obstetricians and Gynaecologists of Canada guidelines, 4.9
 - support for fathers, 4.14–15
 - tobacco use, 4.18
 - use of ALPHA Form, 4.13 (t4.3)
 - violence in pregnancy, 4.16–17
 - weight gain, 4.21–22
 - working conditions, 4.20
- Prenatal education, 4.25
- adolescents, 4.29
 - benefits, 4.25
 - content, 4.27–28
 - familiarization with birth facility, 4.28
 - group prenatal programs, 4.25
 - groups with special needs, 4.28–29
 - issues, 4.27–29
 - requirements of educator, 4.26–27
 - role of prenatal educator, 4.26
- Prenatal hospital “visit” programs, 6.32
- Prenatal visits, 4.7–8
- Preterm birth, 8.8
- Principal level, *see* Level I
- Program administration
- Level I, 2.29
 - Level II, 2.29
 - Level III, 2.29
- Puerperal infection/endometritis
- postpartum care, 6.22
 - treatment, 6.22
- ## Q
- Quality of care indicators, 1.11–12
- ## R
- Records, prenatal care, 4.8–9
- Regional maternal and newborn care, continuum of care, 2.9–11
- Regional Referral and Transport Program, 9.6–7
- Regionalization, *see* Regional maternal and newborn care
- Regionalization of maternal and newborn care, 2.6–11
- advisory groups, 2.8–9
 - definitions of regions, 2.7
 - description, 2.6
 - models, 2.7–11
- Registered nurse responsibilities, Level I, 2.20 (t2.3)
- Remembrance, 8.15
- Risk of newborn problems, factors, 6.43 (App.1)
- Role of health care providers
- nutrition counseling, 3.17–18
 - preconception care, 3.5
 - substance use, 6.26–27
 - violence in pregnancy, 4.16
- Role of media, preconception programs, 3.8–9

S

- Second stage of labour, 5.23–24
 - definition, 5.23
- Secondary level, *see* Level II
- Single-room maternity care, 5.7
- Social support, preconception care, 3.10
- Society of Obstetricians and Gynaecologists of Canada, guidelines, prenatal care, 4.9
- Society of Obstetricians Gynaecologists guidelines, feasibility of, 5.18
- SOGC, *see* Society of Obstetricians and Gynaecologists of Canada
- Staffing requirements, 2.21, 23, 25
- Stillbirth, 8.8
- Substance use
 - effect on newborn, 6.26–27
 - role of health care providers, 6.26–27
- Support areas for antepartum, labour, birth, and postpartum facilities, 10.32–33 (App.2)
- Support groups, perinatal loss, 8.27
- Supportive care
 - advantages, 5.13
 - elements, 5.13

T

- Ten steps to baby-friendly communities, 7.41 (App.4)
- Ten steps to creating a mother-friendly workplace, 7.50 (App.9)
- Ten steps to successful breastfeeding, 7.34 (App.1)
- Tertiary level, *see* Level III
- Therapeutic abortion, 8.8
- Third stage of labour, drug use, 5.25
- Tobacco use, prenatal care, 4.18
- Tobacco reduction resources, 3.29 (App.1)
- Training, health care providers, 1.14
- Transfer of care, guidelines, 5.8

Transport

- care on arrival, 9.12
- care during, 9.11
- care guidelines, 9.12
- contraindications for, 9.8
- equipment, 9.11, 21
- guiding principles, 9.6
- indications for, 9.7–8
- neonatal, 9.12–15
 - indications for, 9.13
 - planning, 9.14–15
 - return to origin, 9.15
- personnel, 9.7
- physician's and midwife's assessment, 9.9
- planning, 9.8–11
 - family communications, 9.8
- Regional Referral and Transport Program, components, 9.6–7
- role of health care providers, 9.10
- transfer record, 9.18–20 (App.1)
- Transport personnel requirements, 9.7

U

- UNICEF Baby Friendly Hospital Initiative and Programme Manual*, 7.22

V

- Violence in pregnancy, role of health care providers, 4.16
- Vitamin A, excessive intakes, 3.20
- Vitamin D intake, preconception counseling, 3.18–19
- Vitamin K, administration, 6.11
- Vitamin and mineral supplementation, during breastfeeding, 7.26

W

- Weaning process, 7.28
- Worden's Tasks of Grieving, 8.10 (t8.3)