


(O T R)

O N T A R I O T R A U M A R E G I S T R Y



2003 REPORT

INJURY DEATHS IN ONTARIO
(INCLUDES 2000–2001 DATA)



Canadian Institute
for Health Information

Institut canadien
d'information sur la santé

**Ontario Trauma Registry
2003 Report
Injury Deaths in Ontario
(includes 2000–2001 data)**

All rights reserved.

No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording, or by any information storage and retrieval system now known or to be invented, without the prior permission in writing from the owner of the copyright, except by a reviewer who wishes to quote brief passages in connection with a review written for inclusion in a magazine, newspaper or broadcast.

Requests for permission should be addressed to:

Canadian Institute for Health Information
377 Dalhousie Street
Suite 200
Ottawa, Ontario
K1N 9N8

Telephone: (613) 241-7860
Fax: (613) 241-8120
www.cihi.ca

ISBN 1-55392-388-X (PDF)

© 2004 Canadian Institute for Health Information

About the Canadian Institute for Health Information (CIHI)

The Canadian Institute for Health Information (CIHI) is an independent, pan-Canadian, not-for-profit organization working to improve the health of Canadians and the health care system by providing quality health information. Committed to safeguarding the privacy and confidentiality of personal health information, CIHI's mandate is to coordinate the development and maintenance of a common approach to health information for Canada. To this end, CIHI is responsible for providing accurate and timely information that is needed to establish sound health policies, manage the Canadian health system effectively and create public awareness of factors affecting good health.

The Institute's mandate is based upon collaborative planning with key stakeholder groups, including all provincial, territorial and federal governments, national health care agencies and service providers.

CIHI is governed by a Board of Directors whose 15 members strike a balance among the health stakeholders, sectors and regions of Canada.

The Institute's core functions are to:

- identify and promote national health indicators;
- coordinate and promote the development and maintenance of national health information standards;
- develop and manage health databases and registries;
- conduct analysis and special studies and participate in research;
- publish reports and disseminate health information; and
- coordinate and conduct education sessions and conferences.

Acknowledgements

This Ontario Trauma Registry (OTR) report is made possible by the contributions of the Office of the Chief Coroner of Ontario, the Trauma Registry Advisory Committee (TRAC), and the Ontario Ministry of Health and Long-Term Care.

The OTR extends sincere thanks to Dr. Barry McLellan, Acting Chief Coroner, and June Frank, Manager of the Coroner's Information System at the Office of the Chief Coroner, for their ongoing support. The OTR also thanks Steve Rach at the Ministry of the Solicitor General, Integrated Justice Information Technology Division for providing death data to the OTR in an electronic format on an annual basis.

The OTR is funded by the Emergency Health Services Branch of the Ontario Ministry of Health and Long-Term Care and managed by the Canadian Institute for Health Information.

This OTR report was developed at CIHI under the direction of Nizar Ladak, Director, Health Services Information, by:

- Cassandra Linton, Senior Analyst, Clinical Registries
- Nicole de Guia, Consultant, Clinical Registries
- Greg Webster, Director, Research and Indicator Development

CIHI staff are thanked for their contributions. CIHI Productions Systems, Health Services, are thanked for their technical support in the programming of the figures and tables. CIHI Publications assisted with the formatting and layout of this report.

All questions regarding this report should be directed to:

Nicole de Guia
Consultant, Clinical Registries
Ontario Trauma Registry
Canadian Institute for Health Information
90 Eglinton Avenue East, Suite 300
Toronto, Ontario
M4P 2Y3
Tel: (416) 481-2002 Ext. 3545
Fax: (416) 481-2950
e-mail: otr@cihi.ca

Executive Summary

The source of data for this report is the Ontario Trauma Registry (OTR) Death Data Set (DDS). This data set contains information on all injury deaths in Ontario, which is provided by the Office of the Chief Coroner of Ontario. The inclusion of an injury or trauma case is based on whether the case met the OTR definition of trauma, “injury resulting from the transfer of energy”, applied clinically.

Provincial Analyses

Five-year Trends

In 2000–2001, there were 3,115 injury deaths in Ontario. This represents a reduction of 10% since 1996–1997 and an average annual decrease of 3%. The age-standardized injury death rate in 2000–2001 was 2.5 deaths per 10,000 population compared to 3.0 per 10,000 in 1996–1997. This represents a five-year reduction of 17% and an average annual decrease of 4%.

Demographics

In 2000–2001, the mean age of injury deaths was 55 years. Males accounted for approximately two-thirds (64%, n = 1,999) of all injury deaths.

Persons 65 years of age and over constituted the largest percentage of injury deaths (41%, n = 1,266), followed by those between the ages of 35 and 64 years (32%, n = 986). Persons aged 20 to 34 years accounted for 17% (n = 544), and those under the age of 20 years accounted for 10% (n = 319).

Causes of Injury Death

Among injury deaths in 2000–2001, the three leading causes of injury were falls (34%, n = 1,063), motor vehicle collisions (25%, n = 769) and suicide and self-inflicted injury (excluding poisoning) (21%, n = 659). Other causes of injury death included drowning (4%, n = 138), homicide and injury purposely inflicted (excluding poisoning) (3%, n = 105) and fire and flames (3%, n = 95).

Injury Deaths Due to Falling

In 2000–2001, unintentional falls accounted for 34% (n = 1,063) of all injury deaths in the province. The majority (85%, n = 902) of these deaths occurred among persons aged 65 years of age and over. Females represented 52% (n = 552) of all fall-related deaths.

Injury Deaths Due to Motor Vehicle Collisions

In 2000–2001, 25% (n = 769) of all injury-related deaths in Ontario were attributed to motor vehicle collisions. Males comprised two-thirds (65%, n = 501) of these deaths.

Of the motor vehicle collision deaths, almost one-half (49%, n = 378) were drivers and one-quarter (27%, n = 207) were passengers. The remainder (24%, n = 184) included motorcyclists, pedestrians and snowmobile riders.

Seatbelts were known to be present in the vehicle for 334 motor vehicle occupant deaths. Among these cases, seatbelts were used in 60% (n = 202) of the cases.

Drugs and/or alcohol were involved in nearly one-quarter (23%, n = 177) of all injury deaths due to motor vehicle collision.

Injury Deaths Due to Suicide

Deaths due to suicide (excluding poisoning) accounted for 21% (n = 659) of all injury deaths in Ontario in 2000–2001. Persons between the ages of 35 and 64 years comprised one-half (50%, n = 329) of these deaths. Firearms were used in 19% (n = 128) of suicide and self-inflicted injury deaths (excluding poisoning) and 14% (n = 90) involved drugs and/or alcohol.

Firearm-Related Deaths

There were 168 firearm related deaths in 2000–2001, representing 5% of injury deaths in Ontario. Nearly all (92%, n = 154) of these deaths were among males. Of all firearm-related injury deaths, 76% (n = 128) were related to suicide and 20% (n = 34) were related to homicide. The remaining deaths were unintentional injuries or injuries where the intent was undetermined.

Work-Related Deaths

In 2000–2001, there were 85 work-related deaths in the province. The most commonly reported specific work environments were farming (25%, n = 21), and inside factory/warehouse work (14%, n = 12), and construction (13%, n = 11). Nearly all (94%, n = 80) of these cases were male, and the mean age was 46 years.

Regional Analyses

Regional analyses were based on where the injury occurred rather than the person's place of residence, and reflect the seven Ontario health planning regions as defined by the Ministry of Health and Long-Term Care. From 1996–1997 to 2000–2001, the North region of Ontario consistently had the highest age-standardized injury death rate compared to the other regions in Ontario (4.6 per 10,000 population in 2000–2001). In contrast, Toronto experienced the lowest age-standardized injury death rate during the five-year period (2.0 per 10,000 population in 2000–2001).

Ontario Trauma Registry 2003 Report

Injury Deaths in Ontario

(includes 2000–2001 data)

Table of Contents

About the Canadian Institute for Health Information (CIHI)	i
Acknowledgements	iii
Executive Summary	v
1. Introduction	1
A. Purpose of Report	1
B. About the Ontario Trauma Registry (OTR)	1
i) Goal	1
ii) History	1
iii) Structure	1
2. Methodological Notes	3
A. Data Source	3
B. Definition of Trauma for Injury Deaths	3
i) Selection Criteria	3
ii) The Classification System of the Office of the Chief Coroner	4
C. Reporting Guidelines.....	5
3. Provincial Analysis.....	6
A. Trend Analysis	6
i) 1991–1992 Through 2000–2001	6
ii) 1996–1997 Through 2000–2001	6
B. Demographics	7
C. Causes of Death	9
i) Overall Causes	9
ii) Causes by Age Group	10
iii) Falls.....	14
iv) Motor Vehicle Collisions	15
v) Drowning.....	17
D. Intentionality of Trauma Deaths	19
i) Suicide	19
ii) Homicide	22
E. Contextual Factors.....	23
i) Firearm-Related Deaths	23
ii) Work-Related Deaths.....	24
iii) Drug and Alcohol Involvement	25
4. Regional Analysis	26
A. Trend Analysis, 1996–1997 Through 2000–2001	26
B. 2000–2001	27
C. Demographics	28
D. Causes of Death	29

Ontario Trauma Registry 2003 Report
Injury Deaths in Ontario
(includes 2000–2001 data)

Table of Contents (cont'd)

List of Figures

Figure 1.	Injury Deaths in Ontario, 1991–1992 Through 2000–2001	6
Figure 2.	Injury Deaths by Age Group, 2000–2001	7
Figure 3.	Injury Deaths by Sex and Single Year of Age, 2000–2001	8
Figure 4.	Causes of Injury Death, 2000–2001	9
Figure 5.	Causes of Injury Deaths Among Persons Under 20 Years, 2000–2001	10
Figure 6.	Causes of Injury Deaths Among Persons Aged 20 to 34 Years, 2000–2001	11
Figure 7.	Causes of Injury Deaths Among Persons Aged 35 to 64 Years, 2000–2001	12
Figure 8.	Causes of Injury Deaths Among Persons Aged 65 Years and Over, 2000–2001	13
Figure 9.	Deaths Due to Unintentional Falls by Age Group, 2000–2001	14
Figure 10.	Motor Vehicle Collision Injury Deaths by Sex and Single Year of Age, 2000–2001	15
Figure 11.	Deaths Due to Drowning by Age Group, 2000–2001	17
Figure 12.	Deaths Due to Drowning by Death Factor, 2000–2001	18
Figure 13.	Deaths Due to Suicide (Excluding Poisoning) by Age Group, 2000–2001	19
Figure 14.	Deaths Due to Suicide (Including Poisoning) by Age Group, 2000–2001	20
Figure 15.	Deaths Due to Suicide (Including Poisoning) by Sex and Suicide Method, 2000–2001	21
Figure 16.	Deaths Due to Homicide (Excluding Poisonings) by Age Group, 2000–2001	22
Figure 17.	Firearm-Related Deaths by Age Group, 2000–2001	23
Figure 18.	Work-Related Deaths by Work Environment, 2000–2001	24
Figure 19.	Injury Deaths with Drug/Alcohol Involvement by Cause of Injury, 2000–2001	25

**Ontario Trauma Registry 2003 Report
Injury Deaths in Ontario
(includes 2000–2001 data)**

Table of Contents (cont'd)

List of Figures (cont'd)

Figure 20. Age-Adjusted Injury Death Rates by Ontario Health Planning Region, 1996–1997 Through 2000–2001	26
Figure 21. Age-Adjusted Injury Death Rates and Number of Injury Deaths by Ontario Health Planning Region, 2000–2001	27
Figure 22. Trauma Deaths by Age Group and Ontario Health Planning Region, 2000–2001	28

Appendices

Appendix A—Definitions of Terms	A-1
Appendix B—Trauma Definition: ICD-9 E Code Inclusions and Exclusions	B-1
Appendix C—Trauma Definition: Mapping Methodology	C-1
Appendix D—Trauma Definition: Death Factors	D-1
Appendix E—List of Tables	E-1

1. Introduction

A. Purpose of Report

The purpose of this report is to provide a descriptive analysis of injury-related deaths in Ontario. The data source for this report is the Ontario Trauma Registry Death Data Set (OTR DDS), which contains information from the Office of the Chief Coroner of Ontario as of September 2003.

B. About the Ontario Trauma Registry (OTR)

i) Goal

The goal of the Ontario Trauma Registry (OTR) is to facilitate the reduction of injury hospitalizations and deaths in the province of Ontario by identifying, describing and quantifying trauma in order to:

1. permit planning and evaluation of prevention programs, legislative changes and cost expenditures
2. aid in resource allocation decisions and contribute to cost reductions

ii) History

The OTR, funded by the Ontario Ministry of Health and Long-Term Care, was established in May 1992. A multidisciplinary advisory committee provides guidance to the OTR. The Trauma Registry Advisory Committee (TRAC) includes representatives from the Ontario Ministry of Health and Long-Term Care, Ministry of Labour, Ministry of Transportation, CIHI, epidemiologists, trauma care providers, the Office of the Chief Coroner of Ontario and the Trauma Association of Canada. The current structure and implementation of the OTR is based on the data elements, data collection procedures, report formats and management procedures determined by TRAC.

The primary users of the OTR include the eleven lead trauma hospitals in Ontario, the members of TRAC and Area Emergency Health Services (EHS) Committees. The Area EHS Committees are part of regional planning networks composed of committees at the provincial, regional and local levels involving health care planners, providers and consumers in emergency health initiatives.

iii) Structure

For injury prevention programs to be effective, data are needed to clearly define the nature and scope of injury in the province. The OTR consists of three major sources of data as listed below. Standard and ad hoc reports from these data sets detail demographic information, cause and nature of injury hospitalizations and deaths both provincially and regionally. This information is used by policy makers, planners, researchers and injury prevention specialists to develop and monitor injury prevention programs, and to improve trauma patient care.

The Ontario Trauma Registry is composed of 3 data sets:

1. The **Minimal Data Set (MDS)** contains demographic, diagnostic and procedural information on all hospitalizations due to injury in acute care hospitals in Ontario. These hospitalizations are selected from the Discharge Abstract Database (DAD) at CIHI and downloaded to the OTR data processing system. Selection criteria for inclusion in the OTR MDS are based on specific External Cause of Injury Codes (E Codes) within the International Classification of Disease (ICD), version 9, coding system.

Examples of E Codes that are included in the definition of trauma are motor vehicle collisions, including those involving pedestrians, motorcycles and bicycles, and falls, drownings and burns. E Codes that are excluded are poisonings, adverse effects and complications. Appendix B (Trauma Definition: ICD-9 E Code Inclusions and Exclusions) lists the E Codes that are included and excluded from the definition of trauma used in the OTR.

2. The **Comprehensive Data Set (CDS)** consists of detailed information on patients hospitalized with major trauma in 11 participating trauma facilities in the province. These lead/trauma hospitals have been funded by the Ministry of Health and Long-Term Care for hardware, software and dedicated trauma staff including a Medical Director, Trauma Coordinator, Data Analyst and Administrative Assistant. The definition of major trauma in the OTR CDS is based on the Injury Severity Score (ISS), an international scoring system created to calculate the severity of injury, and an E Code that falls within the OTR definition of trauma.

Specialized trauma software (COLLECTOR and TRI-CODE from Digital Innovation Inc. and Tri-Analytics, Inc.) is used to collect and analyze data on approximately 3,600 cases annually. This software has been customized for the province of Ontario with input from participating hospitals and TRAC. Detailed data are collected including demographics, pre-hospital and hospital care, and patient outcomes including a 6-month follow up interview. Data are electronically transmitted to the OTR on a monthly basis.

3. The **Death Data Set (DDS)**, the data source for this report, is described in detail in the next chapter.

2. Methodological Notes

A. Data Source

The data source for this report is the **Ontario Trauma Registry Death Data Set (OTR DDS)**. Data comprising the OTR DDS are provided by the Office of the Chief Coroner of Ontario. The OTR DDS contains information on all deaths in the province due to injury, including demographics, cause of death and factors contributing to death such as alcohol use. Reporting on all injury deaths rather than only in-hospital deaths provides a more complete picture of trauma in the province. Information contained in the database at the Office of the Chief Coroner is important to injury prevention programs because a large percentage of injured persons die before admission to hospital. Therefore, these persons are not captured in hospital-based statistics.

B. Definition of Trauma for Injury Deaths

As directed by the Ontario Trauma Patient Registry Task Force Report, the definition of trauma used by the OTR is injury resulting from the transfer of energy. The International Classification of Diseases, version 9 (ICD-9) External Cause of Injury Codes (E Codes) are used to define trauma hospitalizations in the Minimal and Comprehensive Data Sets of the OTR. E Code categories that are included and excluded from the definition of trauma are found in Appendix B (Trauma Definition: ICD-9 E Code Inclusions and Exclusions). However, the death data provided by the Office of the Chief Coroner does not include E Codes. The Office of the Chief Coroner categorizes deaths using a classification system including death types, death factors, environments and involvements. These components describe the cause of death and the events surrounding the death.

To take advantage of the information documented by the Coroner's Office and to allow comparability with other sources of injury information, the OTR has developed a system to map the death type, death factor and environment components of the classification system used by the Office of the Chief Coroner to ICD-9 E Code categories. This allows standardized reporting across the data sets of the OTR and comparisons to other sources of data. However, trauma deaths cannot always be mapped to specific E Codes because of the differences in the ICD and Coroner's coding systems. A summary of the mapping methodology used is found in Appendix C (Trauma Definition: Mapping Methodology).

i) Selection Criteria

The electronic file provided to the OTR by the Integrated Justice Information Technology Division includes:

- all deaths documented with a death type of unintentional, homicide, suicide and undetermined; and
- deaths documented with a death type of natural with documentation indicating a fall in an institution.

ii) The Classification System of the Office of the Chief Coroner

The following are the components of the classification system used by the Office of the Chief Coroner to categorize deaths.

Death Type

Death type is the classification of the intent of the action, force, instrument or disease that caused death. There are six death types defined by the Office of the Chief Coroner including unintentional, homicide, suicide, natural, undetermined and archaeological/skeletal/animal remains. It is important to note that suicide deaths resulting from poisoning are not included in the definition of trauma and therefore are generally not included in trauma reports provided by the OTR. However, Appendix E—Tables 10 and 11 report all suicide deaths (including poisoning) and have been included in this report to provide a more complete representation of suicide deaths in the province.

Death Factors

A death factor is an action, force, instrument or disease occurring in an environment, which leads directly toward death. For each death, the Coroner codes a primary death factor that identifies the most significant circumstances or events leading to death. There are over 80 death factors used by the Office of the Chief Coroner to describe the cause of death. TRAC and lead/trauma hospital staff reviewed these death factors in 1992. This group identified 40 death factors that parallel the definition of trauma used in the Minimal Data Set and that are used to define trauma in this report. A list of these death factors is found in Appendix D (Trauma Definition: Death Factors).

Motor vehicle crashes, shooting and drowning are examples of death factors that meet the definition of trauma. Examples of excluded death factors include alcohol poisoning and intestinal obstruction.

Environments

An environment is a combination of the location and the activity of the deceased when an action, force, instrument or disease was applied which led toward death. Up to four environments may be documented for each death. Over 90 environments are defined by the Office of the Chief Coroner. Environments are divided into the following categories: occupational, recreation and sporting, institutions (patients and inmates), general living and traveling. Examples of environments are hunting, snowmobiling, bicycles, drivers, passengers and pedestrians.

Involvements

An involvement is an activity or circumstance related to the deceased that did not directly lead to death but which may be of significance or a contributing factor. Involvements are generally combinations of certain environments, institutions, overdose agents or death factors that are of particular interest to the Coroner's office. Examples of involvements are alcohol and drug use, a fall in a nursing home or helmet use. The Office of the Chief Coroner codes more than 60 involvements; up to 3 can be coded per case.

C. Reporting Guidelines

This report:

- is created by fiscal year and contains information on all injury deaths occurring in fiscal year 2000–2001 and trend analysis for injury deaths occurring between 1996–1997 through 2000–2001;
- is created by region based on the primary municipality documented by the Office of the Chief Coroner. Primary municipality refers to the geographic location where the injury occurred rather than the place of residence, which is not available from the Office of the Chief Coroner for all injury deaths;
- is created using data downloaded from the Office of the Chief Coroner of Ontario as of September 2003; 2000–2001 data are considered preliminary. Historical data dating back to 1995 have also been updated as of September 2003;
- uses the population of the area in which the injury occurred for the denominator for rate calculations;
- generally presents causes of death by E Code categories, although death factors are reported in Appendix E—Table 3;
- has changed all references to “accident” reported by the ICD or Office of the Chief Coroner definitions to “incident” or “collision” to reinforce injury prevention efforts; “accidental” (as in accidental death type) has been changed to “unintentional”;
- data may not always reconcile with data presented in previous or future reports; discrepancies may exist due to on-going and necessary data updates;
- may present percentages that do not sum to 100% because of rounding;
- does not include deaths with sex not documented in reports based on sex (e.g. single year of age by sex);
- does not include suicide deaths resulting from poisoning as determined by the definition of trauma unless otherwise stated. Only Tables 10 and 11 in Appendix E report all suicide deaths *including* poisoning;
- the sum of deaths in each region does not equal the Ontario total due to cases where the region of incident are unknown; and
- injury death rates are per 10,000 and are age standardized using 1991 population estimates from Statistics Canada.

3. Provincial Analysis

A. Trend Analysis

i) 1991–1992 Through 2000–2001

Between 1991–1992 and 2000–2001 the number of injury-related deaths decreased by 22%, from 3,985 to 3,115. This represents an average annual decrease of 3% over the past 10 years.




Figure 1. Injury Deaths in Ontario, 1991–1992 Through 2000–2001

ii) 1996–1997 Through 2000–2001

For the five fiscal years from 1996–1997 Through 2000–2001:

- the number of deaths decreased by 10% from 3,459 in 1996–1997 to 3,115 in 2000–2001, representing an average annual decrease of 2.5%;
- the age standardized injury death rate decreased from 3.0 per 10,000 population in 1996–1997 to 2.5 per 10,000 in 2000–2001, representing a five-year reduction of 17% and an average annual decrease of 4%.
- mean ages ranged between 51 and 55 years; median ages ranged between 49 and 53 years.
- males accounted for 64% to 67% of injury deaths.

- the proportion of deaths due to motor vehicle collisions ranged between 25% and 27%;
- the proportion of deaths due to unintentional falls ranged between 27% and 34%;
- deaths due to suicide (excluding poisoning) ranged between 21% and 25%, homicides (excluding poisoning) ranged between 3% and 5%; and
- the proportion of injury deaths reported as dead on arrival (DOA) at hospital emergency departments ranged from 15% to 17%.

B. Demographics

In 2000–2001, the mean age of injury deaths was 55 years. Figure 2 shows that the majority of injury deaths occurred among persons aged 65 years and over (41%, n = 1,266), followed by those aged 35 to 64 years (32%, n = 986).




Figure 2. Injury Deaths by Age Group, 2000–2001

Males represented approximately two-thirds (64%, n = 1,999) of all injury deaths. Figure 3 illustrates that the distribution of injury deaths by age among males differed markedly from that of females. Over one-half (60%, n = 654) of female injury deaths occurred among those 65 years and over, whereas a little over one-quarter (30%, n = 604) of injury deaths among males occurred in this age group. The majority (37%, n = 740) of injury deaths among males were among those aged 35 to 64 years and 22% (n = 435) occurred in the 20 to 34 year old age group.




Figure 3. Injury Deaths by Sex and Single Year of Age, 2000–2001*

* Note: 20 cases with unknown sex

C. Causes of Death

i) Overall Causes

Figure 4 shows that the leading three causes of injury-related death in 2000–2001 were unintentional falls (34%, n = 1,063), motor vehicle collisions (25%, n = 769), and suicide (excluding poisoning) (21%, n = 659). The “All other causes” group, which represented 9% (n = 271) of the total, included (but was not limited to):

- deaths due to injuries in which intentionality is undetermined (n = 70);
- natural and environmental factors (n = 42);
- suffocation (n = 32);
- pedal cycle incidents (n = 12); and
- railway incidents (n = 10).




Figure 4. Causes of Injury Death, 2000–2001

ii) Causes by Age Group

Under 20 Years

In 2000–2001, 10% (n=319) of all injury deaths occurred among children and teens under the age of 20 years. The percentage of all injury deaths among this age group is lower than its representation in the general population of Ontario estimated by Statistics Canada as of October 1, 2000, which was 26%.

Figure 5 shows that motor vehicle collisions (45%, n = 143), suicide and self-inflicted injuries (excluding poisoning) (17%, n = 55), and drowning (14%, n = 44) were the leading causes of injury-related death among persons under the age of 20. The majority of suicides (82%, n = 45) occurred among those between the ages of 15 and 19 years. Similarly, the majority (69%, n = 99) of motor vehicle collision deaths in this age group were also between 15 and 19 years of age.




Figure 5. Causes of Injury Deaths Among Persons Under 20 Years, 2000–2001

20 to 34 Years

In 2000–2001, 17% (n=544) of all injury deaths occurred among persons between the ages of 20 and 34 years. By comparison, this age group represented 21% of the general population of Ontario on October 1, 2000.

Figure 6 illustrates that motor vehicle collisions (38%, n=208), suicide and self-inflicted injury (excluding poisoning) (35%, n=190), and homicide and injury intentionally inflicted by another person (excluding poisoning) (6%, n=31) were the leading specific causes of injury death among 20 to 34 year olds. Nearly three-quarters (70%, n=133) of suicide and self-inflicted injuries in this age group were between 25 and 34 years of age.




Figure 6. Causes of Injury Deaths Among Persons Aged 20 to 34 Years, 2000–2001

35 to 64 Years

Over one-third (32%, n=986) of all injury-related deaths in 2000–2001 occurred among persons between the ages of 35 and 64 years. The percentage of all injury deaths experienced by this group is lower than its 40% representation in the general population of Ontario in 2000.

Figure 7 shows that one-third (33%, n=329) of injury deaths in this age group were attributed to suicide and self-inflicted injury (excluding poisoning), followed by motor vehicle collisions (28%, n=274), and unintentional falls (14%, n=138). Nearly one-half (48%, n=66) of deaths due to falls in this age group were among persons aged 55 to 64 years.




Figure 7. Causes of Injury Deaths Among Persons Aged 35 to 64 Years, 2000–2001

65 Years and Over

More than one-third (41%, n = 1,266) of all injury deaths in 2000–2001 occurred among persons 65 years of age and over. The proportion of all injury deaths in the province attributed to this age group is greater than its 12% representation in the Ontario general population as of October 1, 2000.

Figure 8 illustrates that unintentional falls accounted for nearly three-quarters (71%, n = 902) of the injury-related deaths among those 65 years of age and over. Motor vehicle collisions (11%, n = 144) and suicide and self-inflicted injury (excluding poisoning) (7%, n = 85) were also leading causes of injury death in this age group. Over one-half (51%, n = 460) of deaths due to falling in this age group were among persons 85 years of age and over.




Figure 8. Causes of Injury Deaths Among Persons Aged 65 Years and Over, 2000–2001

iii) Falls

Unintentional falls included here are mapped to the ICD-9 External Cause of Injury Code category E880–E888. In 2000–2001, unintentional falls represented 34% (n = 1,063) of all injury-related deaths in the province. Figure 9 shows that of these deaths, the majority (85%, n = 902) occurred among persons aged 65 years and over. Fifty-two percent (n = 552) of all fall cases were female.




Figure 9. Deaths Due to Unintentional Falls by Age Group, 2000–2001

iv) Motor Vehicle Collisions

Motor vehicle collisions presented here are mapped to the ICD-9 External Cause of Injury Code category E810–E825. In 2000–2001, 25% (n = 769) of all injury deaths were due to motor vehicle collisions. More than one-third (36%, n = 274) of these cases were between the ages of 35 and 64 years, followed by those in the 20 to 34 year old age group (27%, n = 208). Persons 65 years of age and over accounted for 19% (n = 144) of motor vehicle collision deaths, with a further 19% (n = 143) of cases under the age of 20 years.

Males represented 65% (n = 501) of these deaths. Figure 10 shows that there is a peak in the number of motor vehicle collisions deaths around the age of 20 years among males.




Figure 10. Motor Vehicle Collision Injury Deaths by Sex and Single Year of Age, 2000–2001*

* **Note:** 7 cases with unknown sex

Of the 769 motor vehicle collision deaths in 2000–2001, nearly one-half (49%, n = 378) were drivers and one-quarter (27%, n = 207) were passengers. The remaining motor vehicle collision injury deaths (24%, n = 184) included motorcyclists, pedestrians and snowmobile riders.

Seatbelt use was not documented for 42% (n = 247) of motor vehicle driver and passenger deaths. Of the 338 motor vehicle occupant deaths where seatbelt use was documented, 60% (n = 202) wore seatbelts and 39% (n = 132) did not. Seatbelts were not present for 4 cases.

Among the 378 motor vehicle driver deaths:

- more than one-third (35%, n = 133) were wearing seatbelts;
- 18% (n = 69) were not wearing seatbelts;
- seatbelt use was not documented for 46% (n = 174); and
- seatbelts were not present in the vehicle in two cases.

Of the 207 motor vehicle passenger deaths, 33% (n = 69) were wearing seatbelts and 30% (n = 63) were not. Seatbelt use was not documented for 35% (n = 73) of motor vehicle passenger deaths, and in two cases seatbelts were not present.

Drugs and/or alcohol were involved in nearly one-quarter (23%, n = 177) of all motor vehicle collision injury deaths. The Office of the Chief Coroner codes drug and/or alcohol use through specific involvement codes. These codes are activities or circumstances that did not directly lead to death but that may have been contributing factors.

v) Drowning

Four percent (n = 138) of injury-related deaths in Ontario were due to drowning in 2000–2001. Note that only deaths due to unintentional drowning are included in this category. Figure 11 shows that the majority of cases occurred among persons less than 20 years (32%, n = 44) and between the ages of 35 and 64 years (32%, n = 44). Males represented 83% (n = 115) of all drowning-related deaths.

Nearly one-quarter (23%, n = 32) of all drowning injury deaths involved alcohol and/or drugs. Of these cases, the majority (88%, n = 28) involved alcohol only.




Figure 11. Deaths Due to Drowning by Age Group, 2000–2001

Drowning by Death Factor

In 2000–2001, there were 188 deaths due to drowning as defined by death factors. Of these cases:

- 24% (n = 45) were under the age of 20 years;
- 22% (n = 42) were between the ages of 20 and 34 years;
- 38% (n = 72) were between the ages of 35 and 64 years; and
- 15% (n = 29) were 65 years of age and over.

Figure 12 shows that the majority of drowning deaths occurred on open water (67%, n = 126), followed by deaths in the bathtub (11%, n = 21), and ponds (8%, n = 15). Pools and other types of water accounted for the remaining 14% (n = 26) of recorded drowning death factors.




Figure 12. Deaths Due to Drowning by Death Factor, 2000–2001

D. Intentionality of Trauma Deaths

Of the injury deaths in Ontario in 2000–2001:

- 73% (n = 2,250) were unintentional;
- 21% (n = 657) were due to suicide and self-inflicted injury (excluding poisoning);
- 3% (n = 105) were due to homicide and injury purposely inflicted by another person; and
- 3% (n = 83) were of undetermined intent.

Intentionality was determined by death type, a component of the classification system used by the Office of the Chief Coroner.

i) Suicide

Suicide (excluding poisoning)

Deaths due to suicide (excluding poisoning) accounted for 21% (n = 659) of all injury-related deaths in 2000–2001. As shown in Figure 13, one-half (50%, n = 329) of suicide deaths (excluding poisonings) occurred among those between the ages of 35 and 64 years. Males accounted for the majority (81%, n = 529) of suicide deaths reported. Firearms were used in 19% (n = 128) of suicides and self-inflicted injury deaths, and 14% (n = 90) involved drugs and/or alcohol.




Figure 13. Deaths Due to Suicide (Excluding Poisoning) by Age Group, 2000–2001

Suicide (including poisoning)

A large portion of all injury is caused by mechanical energy. As a result, poisonings are often excluded in injury reports. In general, this report excludes poisonings because they do not fit the definition of trauma as *injury resulting from the transfer of energy such as mechanical, thermal or electric energy*. Appendix E—Tables 10 and 11 report all suicide deaths *including* poisoning by sex and age to provide a more complete representation of suicide deaths in Ontario. In 2000–2001, suicides *including* poisonings, accounted for 914 deaths. Males represented 78% (n = 710) of these deaths.

Figure 14 shows that when analyzed by age group, over half (56%, n = 511) of all suicide deaths occurred among persons in the 35 to 64 year old age group. Of the suicide deaths in this age group, 44% (n = 224) were between the ages of 35 and 44 years.




Figure 14. Deaths Due to Suicide (Including Poisoning) by Age Group, 2000–2001

In 2000–2001 males comprised 78% (n = 709) of the all suicide deaths including poisonings. Figure 15 shows that among males, hanging (35%, n = 249) was the leading means of suicide followed by the use of firearms (17%, n = 124). The most common methods of suicide among females were the use of drugs and alcohol (32%, n = 66) and hanging (31%, n = 64).

The “all other” category accounted for 13% (n = 123) of all suicide deaths including poisoning. This category included vehicle collision trauma, suffocation, cuts and stabs, cuts from hand tools, setting fire to oneself, other fires, electrocution, and the use of gases, fumes and other poisons.




Figure 15. Deaths Due to Suicide (Including Poisoning) by Sex and Suicide Method, 2000–2001

ii) Homicide

In 2000–2001, 3% (n = 105) of all injury-related deaths were attributed to homicide and injury purposely inflicted by another person (excluding poisoning). Figure 16 shows that the greatest proportion (39%, n = 41) of these deaths was among persons between the ages of 35 and 64 years, followed by those aged 20 to 34 years (30%, n = 31).

There were 34 firearm-related homicides in 2000–2001, accounting for 32% of all homicides. Nearly three-quarters (74%, n = 25) of all firearm-related homicides were inflicted upon males. Drugs and/or alcohol were involved in 25% (n = 26) of injury deaths due to homicide.




Figure 16. Deaths Due to Homicide (Excluding Poisonings) by Age Group, 2000–2001

E. Contextual Factors

i) Firearm-Related Deaths

Firearm related deaths represented 5% (n = 168) of all injury-related deaths in Ontario in 2000–2001. Of these, males accounted for 92% (n = 154).

Of the 168 firearm-related deaths in 2000–2001:

- 76% (n = 128) were related to suicide;
- 20% (n = 34) were related to homicide;
- 3% (n = 5) were unintentional; and
- 1% (n = 1) were of undetermined intent.

Figure 17 shows that persons aged 35 to 64 years accounted for the greatest proportion (53%, n = 89) of firearm-related deaths, followed by persons aged 20 to 34 years (23%, n = 39) and those under 20 years (14%, n = 23).




Figure 17. Firearm-Related Deaths by Age Group, 2000–2001

ii) Work-Related Deaths

The environment component of the classification system used by the Office of the Chief Coroner allows the identification of occupational deaths. There are 20 occupations identified by environment codes. In 2000–2001:

- there were 85 work-related deaths;
- 80 of these deaths (94%) were among males; and
- the mean age was 46 years.

Figure 18 shows the most common environments for work-related death. The leading specified environments in which work-related deaths occurred were:

- farming (25%, n = 21);
- factory, plant or warehouse (inside work) (14%, n = 12); and
- construction (13%, n = 11).




Figure 18. Work-Related Deaths by Work Environment, 2000–2001

iii) Drug and Alcohol Involvement

Figure 19 summarizes the involvement of drugs and/or alcohol in external causes of injury death. Use of drugs and/or alcohol is coded through specific involvement codes defined by the Office of the Chief Coroner. Involvements are activities or circumstances that did not directly lead to death, but may have been contributing factors. The greatest number of deaths involving drugs and/or alcohol was among motor vehicle collisions (n = 177). As a proportion within E Code group, the highest percentage of drug and/or alcohol involvement was reported among railway incident deaths (50%, n = 5), although there were only 10 cases in total reported in this category.




Figure 19. Injury Deaths with Drug/Alcohol Involvement by Cause of Injury, 2000–2001

4. Regional Analysis

A. Trend Analysis, 1996–1997 Through 2000–2001

Figure 20 shows that between 1996–1997 and 2000–2001 the age-adjusted rates of injury death in the seven health planning regions of Ontario declined. In each of the last 5 years the North region was characterized by the highest age-adjusted death rate of all regions, while Toronto experienced the lowest. Regional analyses were based on where the injury occurred rather than the person's place of residence.




Figure 20. Age-Adjusted Injury Death Rates by Ontario Health Planning Region, 1996–1997 Through 2000–2001

B. 2000–2001

Figure 21 illustrates regional variation in the number and rates of injury death in 2000–2001. The North region had 14% (n=423) of injury deaths and was characterized by an injury death rate of 4.6 per 10,000 population. In contrast, the Toronto region had 18% of injury deaths (n=570) and had the lowest injury death rate of 2.0 per 10,000 population.




Figure 21. Age-Adjusted Injury Death Rates and Number of Injury Deaths by Ontario Health Planning Region, 2000–2001

C. Demographics

Figure 22 illustrates the distribution of injury deaths by age group in each health planning region. For cases under the age of 20 years, the range was from 5% in Toronto to 15% in the North region. The proportion of cases between the ages of 20 and 34 years ranged from 15% in the Toronto and Central East regions to 22% in the Central West region. Among cases aged 35 to 64 years, the proportion ranged from 26% in the Central West region to 37% in the North region. Finally, the proportion of cases 65 years of age and over ranged from 28% in the North region to 44% in the Central South and Central East regions.




Figure 22. Trauma Deaths by Age Group and Ontario Health Planning Region, 2000–2001*

* **Note:** 11 deaths are excluded for at least one of the following reasons: injury occurred out-of-province or place unknown, or case had no indication of “primary environment”, or case had unknown age.

D. Causes of Death

Table 1 below shows injury deaths by cause for each health planning region:

- falls ranged from 21% in the North region to 46% in the Toronto region;
- motor vehicle collisions ranged from 9% in the Toronto region to 32% in the Central East region;
- suicide (excluding poisoning) ranged from 18% in each of the Central West and Central East regions to 27% in the Toronto region;
- drowning ranged from 1% in the Toronto to 11% in the North region; and
- homicide (excluding poisoning) ranged from 1% in the Central South region to 7% in the Toronto region.

Table 1. Injury Deaths by Cause of Injury and Health Planning Region of Ontario, 2000–2001**

	SW	CS	CW	CE	T	E	N	TOTAL
Falls	160 (33%)	108 (35%)	163 (38%)	151 (32%)	260 (46%)	130 (32%)	89 (21%)	1,061 (34%)
MVC	139 (28%)	87 (28%)	104 (24%)	151 (32%)	53 (9%)	105 (26%)	125 (30%)	764 (25%)
Suicide*	99 (20%)	69 (22%)	76 (18%)	88 (18%)	156 (27%)	86 (21%)	84 (20%)	658 (21%)
Drowning	17 (3%)	14 (5%)	17 (4%)	20 (4%)	7 (1%)	17 (4%)	46 (11%)	138 (4%)
Homicide*	12 (2%)	4 (1%)	17 (4%)	11 (2%)	42 (7%)	11 (3%)	8 (2%)	105 (3%)
All Other	63 (13%)	26 (8%)	52 (12%)	57 (12%)	52 (9%)	57 (14%)	71 (17%)	378 (12%)
TOTAL	490 (100%)	308 (100%)	429 (100%)	478 (100%)	570 (100%)	406 (100%)	423 (100%)	3,104 (100%)

* Excluding poisoning

** **Note:** 11 deaths are excluded for at least one of the following reasons: injury occurred out-of-province or place unknown, or case had no indication of "primary environment". Therefore, sum of regional totals does not equal the provincial total for each cause of injury.

Appendix A

Definitions of Terms

Note: All references to “accident” according to the International Classification of Diseases (ICD) or Office of the Chief Coroner definitions have been changed to “incident” or “collision” to reinforce injury prevention efforts; “accidental” (as in accidental death type) has been changed to “unintentional”.

CIHI

The Canadian Institute for Health Information (CIHI) is an independent, pan-Canadian, not-for-profit organization working to improve the health of Canadians and the health care system by providing quality health information.

Comprehensive Data Set

One of three data sets held by the Ontario Trauma Registry (OTR). The Comprehensive Data Set (CDS) consists of detailed information on patients hospitalized with major trauma in eleven participating hospitals in the province. These lead/trauma hospitals have been funded by the Ministry of Health and Long-Term Care for hardware, software and dedicated trauma staff including a Medical Director, Trauma Coordinator, Data Analyst and Administrative Assistant. The definition of trauma in the Comprehensive Data Set is based on the Injury Severity Score (ISS), an international scoring system created to calculate the severity of injury, and an appropriate E Code.

Death Data Set

One of three data sets held by of the Ontario Trauma Registry (OTR). Data comprising the Death Data Set (DDS) come from the Office of the Chief Coroner. The OTR DDS contains information on all deaths in the province due to injury, including demographics, cause of death and factors contributing to death such as alcohol use. Reporting on all injury deaths rather than only in-hospital deaths provides a more complete representation of trauma in the province.

Death Factors*

A death factor is an action, force, instrument or disease occurring in an environment that led directly to death. At least one, and up to four, death factors may be documented for each death. The primary death factor refers to the most significant circumstances or events leading to death.

* Definitions are reproduced from the Coroners System Manual, Office of the Chief Coroner, Ministry of the Solicitor General.

Death Type*

Death type is the classification of the intent of the action, force, instrument or disease that caused death. One death type is documented for each death. The following are the 6 death types defined by the Office of the Chief Coroner.

1. Natural
2. Unintentional (defined as accidental by the Office of the Chief Coroner)
3. Suicide
4. Homicide
5. Undetermined
6. Skeletal/archaeological/animal remains

E Codes (External Cause of Injury Codes)

The External Cause of Injury chapter of the ICD-9 coding system allows the classification and analysis of environmental events, circumstances, and conditions as the cause of injury. Examples include Falls (E880–E888) and Motor Vehicle Traffic Incidents (E810–E819).

Environment*

An environment is a combination of the location where and the activity of the deceased when an action, force, instrument or disease was applied that led toward death. Any investigation may involve several different environments.

ICD (International Classification of Diseases)

The International Classification of Diseases is a World Health Organization (WHO) publication that classifies morbidity and mortality information for statistical purposes, and for the indexing of hospital records by disease and operations, for data storage and retrieval.

Injury

Injury and trauma are used synonymously. Please see the definition of trauma.

Injury Deaths

Injury deaths are defined by the Ontario Trauma Registry using components of the classification system of the Chief Coroner that meet the definition of injury as the transfer of energy.

* Definitions are reproduced from the Coroners System Manual, Office of the Chief Coroner, Ministry of the Solicitor General.

Involvements*

An involvement is an activity or circumstance related to the deceased that did not directly lead to death but which may be of significance, or a contributing factor. Involvements are generally combinations of certain environments, institutions, overdose agents or death factors which are of particular interest to the Coroner's Office.

Mean

A measure of central tendency of a set of observations; the average.

Median

A measure of central tendency of a set of observations; 50th percentile (the value such that half of the data points fall above it and half below it).

Minimal Data Set

One of three data sets held by of the Ontario Trauma Registry (OTR). The Minimal Data Set (MDS) contains demographic, diagnostic and procedural information on all acute care hospitalizations due to injury in Ontario. These hospitalizations are selected from the Discharge Abstract Database (DAD) at CIHI and downloaded to the OTR data processing system. Selection is based on specific External Cause of Injury Codes (E Codes) within the ICD coding system.

Motor Vehicle Collision

A transport incident involving a motor vehicle. It is defined as a motor vehicle traffic incident or as a motor vehicle non-traffic incident according to whether the incident occurred on a public highway (traffic) or elsewhere (non-traffic).

Motor Vehicle Non-traffic Incident

Any motor vehicle incident that occurs entirely in any place other than a public highway.

Motor Vehicle Traffic Incident

Any motor vehicle incident occurring on a public highway (e.g. originating, terminating, or involving a vehicle partially on the highway). A motor vehicle incident is assumed to have occurred on the highway unless another place is specified, except in the case of incidents involving only off-road motor vehicles, which are classified as non-traffic incidents unless the contrary is stated.

Municipalities*

The province has been divided into municipalities by the Office of the Chief Coroner each of which has a four-digit code. Major cities or towns have their own code while smaller towns may be included under the township number. A primary municipality is documented by the Office of the Chief Coroner to indicate where the injury occurred.

* Definitions are reproduced from the Coroners System Manual, Office of the Chief Coroner, Ministry of the Solicitor General.

Other Incidents

Refers to the “Other Accidents” ICD-9 E Code category for the E Code range of E916–E928. Unintentional gunshot wounds, injury caused by machinery and explosions are included in this E Code category.

Regions

Regions are identified in this report based on the primary municipality (i.e. where the injury occurred). There are 7 health planning regions in Ontario (Southwest, Central South, Central West, Central East, Toronto, East, and North) as defined by the Ministry of Health and Long-Term Care.

Single Year of Age

Individual values for ages less than one year through one hundred years. This provides more detail than age groups.

Suicide

For the purposes of this report, suicide is defined as intentionally self-inflicted injuries (excluding poisoning) that result in death. Poisonings are excluded from the definition of trauma, and are therefore excluded from trauma reports. Information is provided in this report on all suicide deaths to provide a more complete representation of suicide in the province.

Trauma

Trauma is defined as injury resulting from the transfer of energy (i.e. kinetic, thermal). The OTR Death Data Set defines trauma by forty death factors for unintentional, suicide, homicide or undetermined deaths types. Trauma deaths that are coded with a natural death type that have an involvement code indicating a fall in an institution such as a nursing home are also included in the definition of trauma.

Trauma Registry Advisory Committee (TRAC)

The multidisciplinary group responsible for guiding the implementation and operation of the OTR.

Appendix B

Trauma Definition: ICD-9 E Code Inclusions and Exclusions

Trauma Definition: ICD-9 E Code Inclusions

The following lists the ICD-9 E Code categories reported by the Ontario Trauma Registry (OTR) based on the trauma definition of injury resulting from the transfer of energy.

“Incident” and “unintentional” have been substituted for the terms “accidents” and “accidental” used in the ICD-9 definitions.

Definition	E Code Category
Railway incidents	E800–E807
Motor vehicle traffic incidents	E810–E819
Motor vehicle nontraffic incidents	E820–E825
Pedal cycles	E826
Other road vehicle incidents	E827–E829
Water transport incidents	E830–E838
Air and space transport incidents	E840–E845
Vehicle incidents not elsewhere classifiable	E846–E848
Unintentional falls	E880–E888
Incidents caused by fire and flame	E890–E899
Incidents due to natural and environmental factors	E900–E902, E906–E909
Incidents caused by drowning and suffocation	E910, E913
Foreign bodies (excluding choking)	E914–E915
Other incidents	E916–E928
Suicide and self inflicted injury (excluding poisoning)	E953–E958
Homicide and injury purposely inflicted by other persons	E960–E961, E963–E968
Legal intervention	E970–E976, E978
Injury undetermined whether unintentionally or purposely inflicted	E983–E988
Injury resulting from operations of war	E990–E998

Trauma Definition: ICD-9 E Code Exclusions

The following lists the ICD-9 E Code categories that are excluded from the Ontario Trauma Registry (OTR) definition of trauma.

Definition	E Codes
Poisonings by drugs	E850–E858
Poisoning by gases	E860–E869
Misadventures	E870–E876
Complications	E878–E879
Travel and motion	E903
Hunger, thirst, exposure, neglect	E904
Venomous animals and plants	E905
Inhalation and ingestion of food causing obstruction	E911
Inhalation and ingestion of other objects causing obstruction	E912
Late effects	E929
Drugs, medicinal and biological substances causing adverse effects	E930–E949
Suicide and self inflicted injury (poisonings)	E950–E952
Late effects of self inflicted injury	E959
Assault by poisoning	E962
Late effects of injury purposely inflicted by other person	E969
Injury due to legal intervention	E977
Poisoning undetermined whether unintentionally or purposely inflicted	E980–E982
Late effects intentionality undetermined	E989
Late effects due to war	E999

Appendix C

Trauma Definition: Mapping Methodology

Mapping Methodology

The mapping methodology developed by OTR is divided into the following five sections:

- i. Unintentional deaths (other than air or vehicle crash)
- ii. Unintentional deaths for motor vehicle and air crashes
- iii. Natural deaths
- iv. Intentional and undetermined intentionality deaths
- v. Other E Code categories

i. Unintentional Deaths (Other Than Air or Vehicle Crashes)

A death type indicating an unintentional death and a primary death factor other than a vehicle crash (620) or air crash (636) were mapped to the ICD-9 E Code categories based on primary death factors as shown in Table 1.

Table 1. Mapping Unintentional Deaths (Other Than Air or Vehicle Crashes)

ICD-9 E Code Category	Primary Death Factor
Unintentional Falls (E880–E888)	Fall, jump—different level (660) Fall, jump—same level (665)
Fire and Flames (E890–E899)	Burns—heat (655) Fire—forest, grass (652) Fire—self (651) Fire—structural (650) Fire—vehicle (653)
Natural and Environmental Factors (E900–E902, E906–E909)	Animal bites, kicks (615) Exposure to heat (657) Exposure to cold (658) Lightning (645) Natural disasters (646)
Submersion, Suffocation (E910, E913)	Asphyxia (674) Drowning—bathtub (601) Drowning—open water (600) Drowning—other (605) Drowning—pond, quarry (604) Drowning—private pool (603) Drowning—public pool (602) Hanging (678) Sexual Asphyxia (672) Strangulation (643) Suffocation (670)

Table 1. Mapping Unintentional Deaths (Other Than Air or Vehicle Crashes) (continued)

ICD-9 E Code Category	Primary Death Factor
Other Incidents (E916–E928)	Caught in machinery (622) Cuts from hand tools (611) Cuts and stabs (610) Shooting—air rifle (633) Shooting—handgun (632) Shooting—rifle (630) Shooting—shotgun (631) Shooting—unspecified (634) Electrocution (640) Explosion (640) Crushed and/or buried (621) Burns—acid, caustic (656) Blunt trauma—beating (623) (Death type unintentional only) Blunt trauma—unintentional (625)

ii. Unintentional Deaths for Motor Vehicle and Air Crashes

A death type indicating an unintentional death and a primary death factor of a vehicle crash (620) or air crash (636) were mapped to ICD-9 E Code categories based on environments as shown in Table 2.

Table 2. Mapping Unintentional Deaths for Motor Vehicle and Air Crashes

ICD-9 E Code Category	Environment
Railway Incidents (E800–E807)	Railway worker, employee of railroad (155) Railway shipping, consignor employee (156) Railway, not on board, trespasser (535)
Motor Vehicle Traffic Incidents (E810–E819)	Motor vehicle driver (520) Motor vehicle passenger (521) Motorcycle driver (522) Motorcycle passenger (523) Pedestrian (525) Snowmobiling on road (244) Ambulance (570)
Motor Vehicle Nontraffic Incidents (E820–E825)	Land vehicle—off road (540) Motorcycle, dirt bike, ATV (241) Dunebuggy (245) Go Kart racing on track (242) Snowmobiling off road (243)
Pedal Cyclist (E826)	Bicycle (528)
Other Road Vehicle Incidents (E827–E829)	Horse back riding (220) Public transit—bus, streetcar (526)

Table 2. Mapping Unintentional Deaths for Motor Vehicle and Air Crashes (continued)

ICD-9 E Code Category	Environment
Water Transport Incidents (E830–E838)	Non-power boating (214) Sailboating, sailboarding (213) Power boating (212) Water Skiing (211) Swimming (210) Water vehicle—not recreational (550)
Air and Space Transport Incidents (E840–E845)	Ski diving, parachuting (225) Hang glider (226) Ultra light aircraft (227)
Vehicle Incidents Not Elsewhere Classifiable (E846–E848)	Non-motorized vehicle (541) Travelling—other (599)

iii. Natural Deaths

There are a number of deaths resulting from falls in institutions that are documented with a natural death type. The Trauma Registry Advisory Committee has decided that these deaths should be included in the Injury Deaths in Ontario. Natural deaths with one of the involvements listed below were mapped to the ICD-9 E Code category E880–E888 (unintentional falls).

Involvements:

- Fall in Nursing Home (985)
- Fall in Home for the Aged (986)
- Fall in Private Residence (987)
- Fall, Other (984)

iv. Intentional and Undetermined Intentionality Deaths

The ICD-9 E Code categories in Tables 1 and 2 correspond to deaths that are documented with an unintentional death type. The following points outline the mapping methodology for deaths with death types indicating suicide, homicide and undetermined.

- a) All deaths with a death type of suicide, excluding those deaths due to poisoning as defined by death factors, were mapped to the E Code category of E953–E958 (suicide and self-inflicted injury, excluding poisoning) for trauma reports. Figures 14 and 15 and Tables 10 and 11 in this report include all suicide deaths to show the complete picture of suicide in Ontario.

- b) All deaths with a death type of homicide were mapped to the ICD-9 E Code category of E960–E961, E963–E968 (homicide and injury purposely inflicted by others). Cases with death factors of child abuse (730) and blunt trauma—beating (623) were also mapped to this E Code category. A small number of cases with a death factor indicating blunt trauma—beating who are documented with an unintentional death type were mapped to E916–E928 (other incidents).
- c) All deaths with a death type of undetermined were mapped to the E Code category of E983–E988 (injury undetermined whether accidentally or purposely inflicted).

v. Other E Code Categories

No injury deaths were mapped to the ICD-9 E Code categories listed below because there are no corresponding primary death factors, environments or involvements.

- E914–E915—Foreign bodies
- E970–E976, E978—Legal intervention
- E990–E998—Injury resulting from operations of war

These reporting categories were therefore excluded from all tables in Appendix E of this report.

Appendix D

Trauma Definition: Death Factors

Trauma Definition: Death Factors

The following are the forty Coroner's Death Factors used by the Ontario Trauma Registry to define trauma in the Death Data Set. The death factors are listed alphabetically and include the numeric code (in brackets) assigned by the Office of the Chief Coroner.

Death Factors	
Animal bites, kicks (615)	Exposure to cold (658)
Asphyxia (674)	Fall or jump—different level or height (660)
Blunt Trauma—beating (623)	Fall or jump—same level (665)
Blunt trauma—unintentional (625)	Fire—forest, grass (652)
Burns—heat (655)	Fire—self (651)
Burns—acid, caustic (656)	Fire—structural (650)
Caught in machinery (622)	Fire—vehicle (653)
Child abuse (730)	Hanging (678)
Crushed and/or buried (621)	Lightning (645)
Cuts from hand tools (611)	Natural disasters (i.e. tornado) (646)
Cuts, stabs (610)	Sexual asphyxia (672)
Drowning—open water (600)	Shooting—air rifle (633)
Drowning—bathtub (601)	Shooting—handgun (632)
Drowning—public pool (602)	Shooting—rifle (630)
Drowning—private pool (603)	Shooting—shotgun (631)
Drowning—pond, quarry, casual water (604)	Shooting—unspecified (634)
Drowning—other (605)	Strangulation (643)
Electrocution (640)	Suffocation (670)
Explosion (624)	Trauma of air crash (636)
Exposure to heat (657)	Trauma of vehicle collision (620)

Appendix E

List of Tables

Appendix E—List of Tables

Table of Contents

1.	Trauma Injury Death Highlights—Ontario, 1996–1997 Through 2000–2001	E-1
2.	External Causes of Injury (E Codes) For Trauma Deaths—Ontario, 1996–1997 Through 2000–2001	E-2
3.	Death Factors for Trauma Deaths—Ontario, 1996–1997 through 2000–2001	E-4
4.	Trauma Death Rate Per 10,000 Population by County/Regional Municipality/District of Injury, 1996–1997 Through 2000–2001	E-7
5.	External Causes of Injury (E Codes) By Age Group For Trauma Deaths— Ontario, 2000–2001	E-11
6.	Intentionality (Death Type) by Age Group and Sex For Trauma Deaths— Ontario, 2000–2001	E-13
7.	Drug and Alcohol Involvements by External Causes of Injury (E Codes) for Trauma Deaths—Ontario, 2000–2001	E-14
8.	Reported Seatbelt Use for Motor Vehicle Occupant Deaths—Ontario, 2000–2001	E-16
9.	Firearm Related Deaths by Death Type and Sex—Ontario, 2000–2001	E-17
10.	Suicide (Including Poisoning) by Sex—Ontario, 1996–1997 Through 2000–2001	E-18
11.	Mechanism of Suicide (Including Poisoning) by Sex and Age, 2000–2001	E-19
12.	Trauma Deaths by Age Group and Sex—by Region, 2000–2001	E-21
13.	Motor Vehicle Trauma Deaths by Age Group and Sex— by Region, 2000–2001	E-22
14.	External Causes of Injury by Sex, 2000–2001	E-23

TRAUMA INJURY DEATH HIGHLIGHTS - ONTARIO, 1996-1997 Through 2000-2001

	1996-1997		1997-1998		1998-1999		1999-2000		2000-2001	
No. of Trauma Deaths	3,459		3,452		3,381		3,428		3,115	
Mean Age (Years)	51.4		51.6		52.1		53.1		54.5	
Median Age (Years)	49		50		49		51		53	
Death Rate per 10,000*	3.0		3.0		2.9		2.8		2.5	
	No.	%								
Males	2,300	66.5	2,287	66.3	2,263	66.9	2,247	65.5	1,999	64.2
D.O.A	600	17.3	574	16.6	510	15.1	568	16.6	504	16.2
MVC Deaths	930	26.9	898	26.0	898	26.6	912	26.6	769	24.7
Seatbelt Worn**	230	24.7	186	20.7	246	27.4	257	28.2	206	26.8
Firearm Injuries	279	8.1	266	7.7	258	7.6	237	6.9	168	5.4
Unintentional Falls	955	27.6	914	26.5	919	27.2	1,022	29.8	1,063	34.1
Farming Deaths	13	0.4	14	0.4	12	0.4	10	0.3	21	0.7
Pediatric Deaths	168	4.9	197	5.7	167	4.9	140	4.1	139	4.5
Cycling Deaths	18	0.5	26	0.8	41	1.2	15	0.4	12	0.4

* Death rates are per 10,000 population and are age standardized using Canada 1991 population estimates from Statistics Canada.

** Count includes all motor vehicle occupants, regardless of whether they were drivers or passengers. Denominator for percentage is the total number of MVC deaths.

**EXTERNAL CAUSES OF INJURY (E CODES)
FOR TRAUMA DEATHS - ONTARIO, 1996-1997 Through 2000-2001**

E CODE	1996-1997		1997-1998		1998-1999		1999-2000		2000-2001	
	No.	%								
	3,459	100.0	3,452	100.0	3,381	100.0	3,428	100.0	3,115	100.0
E800-807 RAILWAY	14	0.4	14	0.4	13	0.4	12	0.4	10	0.3
E810-819 MOTOR VEHICLE TRAFFIC	919	26.6	883	25.6	883	26.1	900	26.3	748	24.0
E820-825 MOTOR VEHICLE NON TRAFFIC	11	0.3	15	0.4	15	0.4	12	0.4	21	0.7
E826 PEDAL CYCLE	18	0.5	26	0.8	41	1.2	15	0.4	12	0.4
E827-829 OTHER ROAD VEHICLE	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
E830-838 WATER TRANSPORT	0	0.0	1	0.0	0	0.0	0	0.0	0	0.0
E840-845 AIR AND SPACE TRANSPORT	11	0.3	8	0.2	12	0.4	23	0.7	7	0.2
E846-848 VEHICLE INCIDENTS NOT ELSEWHERE CLASSIFIED	1	0.0	0	0.0	0	0.0	0	0.0	3	0.1
E880-888 UNINTENTIONAL FALLS	955	27.6	914	26.5	919	27.2	1,022	29.8	1,063	34.1
E890-899 FIRE AND FLAMES	108	3.1	134	3.9	103	3.0	96	2.8	95	3.0
E900-902 & E906-909 NATURAL AND ENVIRONMENTAL FACTORS	34	1.0	26	0.8	44	1.3	30	0.9	42	1.3
E910 DROWNING	169	4.9	173	5.0	157	4.6	169	4.9	138	4.4

**EXTERNAL CAUSES OF INJURY (E CODES)
FOR TRAUMA DEATHS - ONTARIO, 1996-1997 Through 2000-2001**

E CODE	1996-1997		1997-1998		1998-1999		1999-2000		2000-2001	
	No.	%								
	3,459	100.0	3,452	100.0	3,381	100.0	3,428	100.0	3,115	100.0
E913 SUFFOCATION	44	1.3	36	1.0	46	1.4	29	0.8	32	1.0
E916-928 OTHER INCIDENTS	123	3.6	110	3.2	114	3.4	86	2.5	110	3.5
E953-958 SUICIDE & SELF INFLICTED INJURY (EXCL.POISONINGS)	792	22.9	851	24.7	801	23.7	793	23.1	659	21.2
E960-961 & E963-968 HOMICIDE AND INJURY PURPOSELY INFLICTED (EXCL.POISONINGS)	179	5.2	175	5.1	142	4.2	148	4.3	105	3.4
E983-988 UNDETERMINED WHETHER UNINTENTIONALLY OR PURPOSELY INFLICTED	81	2.3	86	2.5	91	2.7	93	2.7	70	2.2

NOTE: Trauma deaths are mapped to ICD E-Codes using components of the Office of the Chief Coroner's classification system.

**DEATH FACTORS FOR
TRAUMA DEATHS - ONTARIO, 1996-1997 Through 2000-2001**

DEATH FACTOR	1996-1997		1997-1998		1998-1999		1999-2000		2000-2001	
	No.	%								
	3,455	100.0	3,443	100.0	3,374	100.0	3,417	100.0	3,096	100.0
Animal Bites (615)	2	0.1	4	0.1	3	0.1	1	0.0	2	0.1
Asphyxia (674)	28	0.8	14	0.4	17	0.5	11	0.3	28	0.9
Blunt Trauma, Accidental (625)	58	1.7	45	1.3	53	1.6	54	1.6	74	2.4
Blunt Trauma - Beating(623)	28	0.8	28	0.8	32	0.9	28	0.8	22	0.7
Burns, Acid (656)	1	0.0	2	0.1	1	0.0	1	0.0	1	0.0
Burns, Heat (655)	12	0.3	13	0.4	12	0.4	13	0.4	15	0.5
Caught in Machinery (622)	9	0.3	9	0.3	8	0.2	11	0.3	4	0.1
Child Abuse (730)	9	0.3	6	0.2	2	0.1	5	0.1	3	0.1
Crushed and/or Buried (621)	33	1.0	38	1.1	27	0.8	23	0.7	29	0.9
Cuts from Hand Tools (611)	2	0.1	1	0.0	1	0.0	1	0.0	1	0.0
Cuts, Stabs (610)	84	2.4	82	2.4	84	2.5	67	2.0	58	1.9
Drowning, Bathtub (601)	22	0.6	24	0.7	20	0.6	26	0.8	21	0.7
Drowning, Open Water (600)	183	5.3	165	4.8	146	4.3	168	4.9	126	4.1
Drowning, Other (605)	4	0.1	6	0.2	14	0.4	7	0.2	10	0.3
Drowning, Pond/Quarry/Casual Water (604)	23	0.7	19	0.6	15	0.4	16	0.5	15	0.5

**DEATH FACTORS FOR
TRAUMA DEATHS - ONTARIO, 1996-1997 Through 2000-2001**

DEATH FACTOR	1996-1997		1997-1998		1998-1999		1999-2000		2000-2001	
	No.	%								
	3,455	100.0	3,443	100.0	3,374	100.0	3,417	100.0	3,096	100.0
Drowning, Private Pool (603)	8	0.2	14	0.4	19	0.6	14	0.4	14	0.5
Drowning, Public Pool (602)	2	0.1	2	0.1	1	0.0	3	0.1	2	0.1
Electrocution (640)	18	0.5	6	0.2	18	0.5	13	0.4	13	0.4
Explosion (624)	6	0.2	7	0.2	7	0.2	4	0.1	2	0.1
Exposure to Cold (658)	37	1.1	25	0.7	47	1.4	27	0.8	44	1.4
Exposure to Heat (657)	2	0.1	2	0.1	2	0.1	2	0.1	1	0.0
Fall or Jump, Diff.Level (660)	312	9.0	294	8.5	299	8.9	313	9.2	292	9.4
Fall or Jump, Same Level (665)	743	21.5	718	20.9	725	21.5	797	23.3	834	26.9
Fire, Forest or Grass (652)	1	0.0	0	0.0	1	0.0	0	0.0	0	0.0
Fire, Self (651)	32	0.9	28	0.8	21	0.6	21	0.6	15	0.5
Fire, Structural (650)	89	2.6	124	3.6	92	2.7	68	2.0	76	2.5
Fire, Vehicle (653)	6	0.2	11	0.3	6	0.2	16	0.5	10	0.3
Hanging (678)	333	9.6	400	11.6	364	10.8	384	11.2	322	10.4
Lightning (645)	1	0.0	3	0.1	4	0.1	4	0.1	1	0.0
Natural Disasters, Tornado (646)	0	0.0	0	0.0	0	0.0	0	0.0	1	0.0

**DEATH FACTORS FOR
TRAUMA DEATHS - ONTARIO, 1996-1997 Through 2000-2001**

DEATH FACTOR	1996-1997		1997-1998		1998-1999		1999-2000		2000-2001	
	No.	%								
	3,455	100.0	3,443	100.0	3,374	100.0	3,417	100.0	3,096	100.0
Sexual Asphyxia (672)	10	0.3	6	0.2	7	0.2	7	0.2	5	0.2
Shooting, Air Rifle (633)	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Shooting, Handgun (632)	85	2.5	73	2.1	63	1.9	59	1.7	55	1.8
Shooting, Rifle (630)	100	2.9	99	2.9	101	3.0	89	2.6	64	2.1
Shooting, Shotgun (631)	94	2.7	93	2.7	90	2.7	82	2.4	48	1.6
Shooting, Unspecified (634)	0	0.0	1	0.0	4	0.1	7	0.2	1	0.0
Strangulation (643)	17	0.5	15	0.4	7	0.2	13	0.4	10	0.3
Suffocation (670)	43	1.2	58	1.7	41	1.2	27	0.8	28	0.9
Trauma of Vehicle Collision (620)	1,007	29.1	1,000	29.0	1,008	29.9	1,009	29.5	842	27.2
Trauma, Air Crash (636)	11	0.3	8	0.2	12	0.4	26	0.8	7	0.2

NOTE: This table excludes deaths where there is no indication of a Primary Death Factor or the Primary Death Factor is not included in the above categories.

**TRAUMA DEATH RATE PER 10,000* POPULATION
BY COUNTY/REGIONAL MUNICIPALITY/DISTRICT OF INJURY, 1996-1997 Through 2000-2001**

Region - County/R.M./District Name**	1996-1997		1997-1998		1998-1999		1999-2000		2000-2001	
	No. Deaths	Rate								
Southwestern - Bruce	26	3.6	31	4.3	31	5.3	23	3.5	43	6.0
- Elgin	22	2.8	36	4.3	28	3.3	27	3.3	32	3.7
- Essex	101	2.7	106	2.9	107	2.8	127	3.2	101	2.5
- Grey	32	3.3	24	3.0	33	3.5	29	2.8	37	3.9
- Huron	29	4.2	28	4.4	23	3.5	27	4.2	28	4.0
- Kent	44	3.9	35	3.0	49	4.2	31	2.8	31	2.5
- Lambton	39	2.9	41	2.9	60	4.3	33	2.3	53	4.0
- Middlesex	132	3.2	127	3.0	84	2.0	96	2.2	119	2.6
- Oxford	41	3.9	51	4.8	47	4.1	51	4.4	18	1.7
- Perth	48	5.8	29	3.0	45	5.4	33	3.7	28	3.0
Southwestern Total	514	3.3	508	3.3	507	3.2	477	2.9	490	2.9
Central South - Brant	43	3.3	36	2.8	43	3.4	47	3.6	36	2.7
- Haldimand-Norfolk R.M.	40	3.5	36	3.5	38	3.4	43	3.7	36	3.1
- Hamilton-Wentworth R.M.	179	3.3	156	2.9	154	2.8	153	2.8	132	2.3
- Niagara R.M.	139	3.0	130	2.8	131	2.9	147	3.1	104	2.3
Central South Total	401	3.2	358	2.8	366	3.0	390	3.1	308	2.4

**TRAUMA DEATH RATE PER 10,000* POPULATION
BY COUNTY/REGIONAL MUNICIPALITY/DISTRICT OF INJURY, 1996-1997 Through 2000-2001**

Region - County/R.M./District Name**	1996-1997		1997-1998		1998-1999		1999-2000		2000-2001	
	No. Deaths	Rate								
Central West										
- Dufferin	17	3.7	29	6.1	24	5.1	23	5.1	14	3.0
- Halton R.M.	87	2.7	121	3.5	102	2.9	94	2.5	95	2.5
- Peel R.M.	208	3.0	173	2.4	187	2.4	187	2.4	172	2.1
- Waterloo R.M.	86	2.0	84	2.0	88	2.1	68	1.5	101	2.3
- Wellington	58	3.2	45	2.4	44	2.3	69	3.6	47	2.5
Central West Total	456	2.7	452	2.6	445	2.4	441	2.3	429	2.2
Central East										
- Durham R.M.	121	3.0	122	2.8	126	3.0	127	2.8	131	2.8
- Haliburton	14	10.1	8	5.2	9	5.4	17	8.8	9	3.9
- Northumberland	35	3.8	32	3.6	31	3.4	39	3.8	36	3.8
- Peterborough	53	3.5	63	4.2	56	3.3	57	3.8	48	3.0
- Simcoe	113	3.3	121	3.5	133	3.6	121	3.2	122	3.1
- Victoria	20	2.8	19	2.6	25	2.9	26	3.5	30	4.1
- York R.M.	106	2.0	118	2.0	92	1.6	123	1.9	102	1.6
Central East Total	462	2.8	483	2.8	472	2.7	510	2.8	478	2.5

**TRAUMA DEATH RATE PER 10,000* POPULATION
BY COUNTY/REGIONAL MUNICIPALITY/DISTRICT OF INJURY, 1996-1997 Through 2000-2001**

Region - County/R.M./District Name**	1996-1997		1997-1998		1998-1999		1999-2000		2000-2001	
	No. Deaths	Rate								
Toronto	647	2.4	664	2.4	649	2.4	645	2.3	570	2.0
Eastern										
- Frontenac	54	3.7	44	3.0	52	3.6	57	4.1	52	3.4
- Hastings	60	4.6	49	3.9	49	3.8	52	4.3	41	3.1
- Lanark	23	4.2	27	4.6	25	3.9	31	4.6	19	3.0
- Leeds & Grenville	40	2.6	47	2.6	38	2.6	35	2.3	38	2.1
- Lennox & Addington	13	3.4	12	3.0	14	3.7	11	2.6	11	2.7
- Ottawa-Carleton R.M.	171	2.3	156	2.0	154	2.0	178	2.2	138	1.7
- Prescott & Russell	22	2.3	17	1.7	19	1.5	21	1.5	27	1.6
- Prince Edward	8	3.3	8	3.6	12	4.7	13	5.4	5	1.4
- Renfrew	36	3.7	39	3.8	36	3.3	45	4.4	40	3.8
- Stormont, Dundas & Glen.	39	1.7	49	1.9	54	2.3	36	1.2	35	1.6
Eastern Total	466	3.0	448	2.8	453	2.8	479	2.9	406	2.4

**TRAUMA DEATH RATE PER 10,000* POPULATION
BY COUNTY/REGIONAL MUNICIPALITY/DISTRICT OF INJURY, 1996-1997 Through 2000-2001**

Region - County/R.M./District Name**	1996-1997		1997-1998		1998-1999		1999-2000		2000-2001	
	No. Deaths	Rate								
North										
- Algoma District	55	4.1	60	4.5	50	3.6	46	3.3	49	3.5
- Cochrane District	51	5.5	52	5.6	43	4.6	54	6.0	29	3.1
- Manitoulin District	14	11.7	6	4.1	13	10.2	15	11.3	3	2.2
- Muskoka District	34	6.0	40	7.5	36	6.7	32	5.5	45	8.4
- Nipissing District	33	3.6	46	4.9	48	5.5	42	4.9	45	4.9
- Parry Sound District	38	9.8	38	8.9	31	8.7	27	6.1	38	8.5
- Sudbury R.M.	71	4.3	82	4.8	64	3.8	58	3.5	48	2.9
- Sudbury District	22	8.2	20	7.4	24	9.1	26	9.9	20	8.3
- Timiskaming District	14	3.6	17	4.6	21	5.5	13	2.8	20	5.4
- Kenora District	74	11.6	71	11.0	53	8.2	60	9.0	63	9.6
- Rainy River District	11	4.3	20	7.9	16	6.2	15	6.8	12	4.7
- Thunder Bay District	77	4.8	75	4.7	76	4.5	83	5.0	51	3.0
North Total	494	5.4	527	5.8	475	5.2	471	5.2	423	4.6
Ontario	3,440	3.0	3,440	3.0	3,367	2.9	3,413	2.8	3,104	2.5

* Death rates are per 10,000 population and are age standardized using Canada 1991 population estimates from Statistics Canada.

** Based on 'Primary Municipality', which indicates where an injury occurred.

NOTE: This table excludes deaths where there is no indication of 'Primary Environment' or the death occurred out of province.

**EXTERNAL CAUSES OF INJURY (E CODES) BY AGE GROUP
FOR TRAUMA DEATHS - ONTARIO, 2000-2001**

	<1	1-4	5-9	10-14	15-19	20-24	25-34	35-44	45-54	55-64	65-74	75-84	85+	Unk	Total	%
E CODE	16	27	30	51	195	203	341	375	350	261	282	462	522	0	3,115	100.0
% of DEATHS	0.5	0.9	1.0	1.6	6.3	6.5	10.9	12.0	11.2	8.4	9.1	14.8	16.8	0.0	100.0	
E800-807 RAILWAY	0	0	0	0	2	0	2	2	3	0	0	0	1	0	10	0.3
E810-819 MOTOR VEHICLE TRAFFIC	5	7	13	16	95	97	103	104	96	68	58	61	25	0	748	24.0
E820-825 MOTOR VEHICLE NON TRAFFIC	0	0	0	3	4	2	6	2	0	4	0	0	0	0	21	0.7
E826 PEDAL CYCLE	0	0	1	2	0	0	3	0	3	2	1	0	0	0	12	0.4
E827-829 OTHER ROAD VEHICLE	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0
E830-838 WATER TRANSPORT	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0
E840-845 AIR AND SPACE TRANSPORT	0	0	0	0	0	0	0	2	1	1	1	2	0	0	7	0.2
E846-848 VEHICLE INCIDENTS NOT ELSEWHERE CLASSIFIED	0	0	1	0	1	0	0	0	0	0	0	1	0	0	3	0.1
E880-888 UNINTENTIONAL FALLS	0	1	0	1	5	5	11	23	49	66	120	322	460	0	1,063	34.1
E890-899 FIRE AND FLAMES	0	3	6	3	2	4	9	11	12	8	14	14	9	0	95	3.0
E900-902 & E906-909 NATURAL AND ENVIRONMENTAL FACTORS	0	0	0	0	0	2	4	9	3	3	6	10	5	0	42	1.3
E910 DROWNING	3	10	2	11	18	8	26	15	18	11	9	6	1	0	138	4.4
E913 SUFFOCATION	2	3	2	2	0	1	6	5	6	1	0	2	2	0	32	1.0

**EXTERNAL CAUSES OF INJURY (E CODES) BY AGE GROUP
FOR TRAUMA DEATHS - ONTARIO, 2000-2001**

	<1	1-4	5-9	10-14	15-19	20-24	25-34	35-44	45-54	55-64	65-74	75-84	85+	Unk	Total	%
E CODE	16	27	30	51	195	203	341	375	350	261	282	462	522	0	3,115	100.0
% of DEATHS	0.5	0.9	1.0	1.6	6.3	6.5	10.9	12.0	11.2	8.4	9.1	14.8	16.8	0.0	100.0	
E916-928 OTHER INCIDENTS	1	0	2	2	7	7	15	21	15	20	13	4	3	0	110	3.5
E953-958 SUICIDE & SELF INFLICTED INJURY (EXCL.POISONINGS)	1	0	0	9	45	57	133	146	121	62	46	28	11	0	659	21.2
E960-961 & E963-968 HOMICIDE AND INJURY PURPOSELY INFLICTED (EXCL.POISONINGS)	2	3	2	1	13	16	15	23	12	6	6	4	2	0	105	3.4
E983-988 UNDETERMINED WHETHER UNINTENTIONALLY OR PURPOSELY INFLICTED	2	0	1	1	3	4	8	12	11	9	8	8	3	0	70	2.2

**INTENTIONALITY (DEATH TYPE) BY AGE GROUP AND SEX
FOR TRAUMA DEATHS - ONTARIO, 2000-2001**

	<1	1-4	5-9	10-14	15-19	20-24	25-34	35-44	45-54	55-64	65-74	75-84	85+	UNK	Total	%
TOTAL																
- FEMALES	5	9	11	21	50	44	64	83	75	80	105	214	335	0	1,096	35.4
- MALES	10	18	19	30	143	158	277	286	274	180	174	246	184	0	1,999	64.6
No. of DEATHS	15	27	30	51	193	202	341	369	349	260	279	460	519	0	3,095	100.0
% of DEATHS	0.5	0.9	1.0	1.6	6.2	6.5	11.0	11.9	11.3	8.4	9.0	14.9	16.8	0.0	100.0	
UNINTENTIONAL																
- FEMALES	4	8	9	17	36	24	34	50	48	61	85	205	331	0	912	29.5
- MALES	6	16	18	23	96	100	150	140	154	120	132	212	171	0	1,338	43.2
No. of DEATHS	10	24	27	40	132	124	184	190	202	181	217	417	502	0	2,250	72.7
% of DEATHS IN AGE GRP	66.7	88.9	90.0	78.4	68.4	61.4	54.0	51.5	57.9	69.6	77.8	90.7	96.7	0.0		
SUICIDE*																
- FEMALES	0	0	0	4	12	12	24	22	22	13	14	3	1	0	127	4.1
- MALES	1	0	0	5	33	45	109	121	100	49	32	25	10	0	530	17.1
No. of DEATHS	1	0	0	9	45	57	133	143	122	62	46	28	11	0	657	21.2
% of DEATHS IN AGE GRP	6.7	0.0	0.0	17.6	23.3	28.2	39.0	38.8	35.0	23.8	16.5	6.1	2.1	0.0		
HOMICIDE*																
- FEMALES	0	1	1	0	1	6	5	8	2	3	1	2	0	0	30	1.0
- MALES	2	2	1	1	12	10	10	15	10	3	5	2	2	0	75	2.4
No. of DEATHS	2	3	2	1	13	16	15	23	12	6	6	4	2	0	105	3.4
% of DEATHS IN AGE GRP	13.3	11.1	6.7	2.0	6.7	7.9	4.4	6.2	3.4	2.3	2.2	0.9	0.4	0.0		
UNDETERMINED																
- FEMALES	1	0	1	0	1	2	1	3	3	3	5	4	3	0	27	0.9
- MALES	1	0	0	1	2	3	8	10	10	8	5	7	1	0	56	1.8
No. of DEATHS	2	0	1	1	3	5	9	13	13	11	10	11	4	0	83	2.7
% of DEATHS IN AGE GRP	13.3	0.0	3.3	2.0	1.6	2.5	2.6	3.5	3.7	4.2	3.6	2.4	0.8	0.0		

* Excluding poisoning.

This table excludes deaths where the Death Type is not included in the above categories.

Note: 20 cases have an unknown sex.

**DRUG AND ALCOHOL INVOLVEMENTS BY EXTERNAL CAUSES OF INJURY (E CODES)
FOR TRAUMA DEATHS - ONTARIO, 2000-2001**

E CODE	INVOLVEMENT			TOTAL No. of DEATHS W/INVOL.	TOTAL No. of DEATHS	% of DEATHS*
	DRUGS ONLY	ALCOHOL ONLY	DRUGS & ALCOHOL			
	54	416	36	506	3,115	16.2
E800-807 RAILWAY	0	5	0	5	10	50.0
E810-819 MOTOR VEHICLE TRAFFIC	9	154	4	167	748	22.3
E820-825 MOTOR VEHICLE NON TRAFFIC	1	7	2	10	21	47.6
E826 PEDAL CYCLE	1	3	0	4	12	33.3
E827-829 OTHER ROAD VEHICLE	0	0	0	0	0	0.0
E830-838 WATER TRANSPORT	0	0	0	0	0	0.0
E840-845 AIR AND SPACE TRANSPORT	0	0	0	0	7	0.0
E846-848 VEHICLE INCIDENTS NOT ELSEWHERE CLASSIFIED	0	0	0	0	3	0.0
E880-888 UNINTENTIONAL FALLS	6	68	3	77	1,063	7.2
E890-899 FIRE AND FLAMES	3	28	0	31	95	32.6
E900-902 & E906-909 NATURAL AND ENVIRONMENTAL FACTORS	1	15	2	18	42	42.9
E910 DROWNING	1	28	3	32	138	23.2

**DRUG AND ALCOHOL INVOLVEMENTS BY EXTERNAL CAUSES OF INJURY (E CODES)
FOR TRAUMA DEATHS - ONTARIO, 2000-2001**

E CODE	INVOLVEMENT			TOTAL No. of DEATHS W/INVOL.	TOTAL No. of DEATHS	% of DEATHS*
	DRUGS ONLY	ALCOHOL ONLY	DRUGS & ALCOHOL			
	54	416	36	506	3,115	16.2
E913 SUFFOCATION	0	6	1	7	32	21.9
E916-928 OTHER INCIDENTS	1	16	1	18	110	16.4
E953-958 SUICIDE & SELF INFLICTED INJURY (EXCL.POISONINGS)	19	58	13	90	659	13.7
E960-961 & HOMICIDE AND INJURY E963-968 PURPOSELY INFLICTED	8	14	4	26	105	24.8
E983-988 UNDETERMINED WHETHER UNINTENTIONALLY OR PURPOSELY INFLICTED	4	14	3	21	70	30.0

Involvements for alcohol use are:

- Alcohol Involvement, Driver (815)
- Alcohol Involvement, Passenger (820)
- Alcohol Involvement, Pedestrian (825)
- Alcohol Involvement, Other Driver (830)
- Alcohol Involvement, Accidents not M.V.C (835)
- Alcohol Involvement (980)

* Involving drugs and/or alcohol.

Involvements for drug use are:

- Cocaine Detected in System (811)
- Drug Involvement (970)

**REPORTED SEATBELT USE FOR
MOTOR VEHICLE OCCUPANT DEATHS - ONTARIO, 2000-2001**

	SEATBELTS PRESENT				SEATBELTS NOT PRESENT		SEATBELT USE NOT DOCUMENTED		TOTAL No. of CASES	
	WORN		NOT WORN		TOTAL	%	TOTAL	%	TOTAL	%
	TOTAL	%	TOTAL	%						
TOTAL	202	34.5	132	22.6	4	0.7	247	42.2	585	100
MOTOR VEHICLE DRIVER	133	35.2	69	18.3	2	0.5	174	46.0	378	100
MOTOR VEHICLE PASSENGER	69	33.3	63	30.4	2	1.0	73	35.3	207	100

Note: Of the 769 MVC Deaths in 2000-2001, 184 of these deaths are not documented as motor vehicle occupants (i.e. drivers or passengers). These 184 Deaths include motorcyclists, pedestrians and snowmobilers.

Primary Environments reported in this table are:

- Motor Vehicle, Driver (520)
- Motor Vehicle, Passenger (521)
- Commercial Driver, Truck/taxi/Etc. (163)

Seatbelt use is documented as involvements:

- Seatbelt, Present But Not Worn (906)
- Seatbelt, Present Worn By Deceased (907)
- Seatbelt, Not Present (908)

**FIREARM RELATED DEATHS
BY DEATH TYPE AND SEX - ONTARIO, 2000-2001**

	UNINTENTIONAL	SUICIDE	HOMICIDE	UNDETERMINED	TOTAL	%
TOTAL						
No. OF DEATHS	4	127	34	1	166	100.0
% OF DEATHS*	2.4	76.5	20.5	0.6	100.0	
Males						
NO. OF DEATHS	4	124	25	1	154	92.8
% OF DEATHS*	2.6	80.5	16.2	0.6	100.0	
Females						
NO. OF DEATHS	0	3	9	0	12	7.2
% OF DEATHS*	0.0	25.0	75.0	0.0	100.0	

* Percentage calculation based on row totals.

Note: 2 cases have an unknown sex and were therefore excluded from this table.

**SUICIDE* (INCLUDING POISONING) BY SEX
ONTARIO, 1996-1997 Through 2000-2001**

		1996-1997	1997-1998	1998-1999	1999-2000	2000-2001
Males	# of Deaths	849	913	856	824	710
	Mean Age	45	44	44	45	43
	Median Age	41	42	41	43	41
Females	# of Deaths	292	287	231	237	204
	Mean Age	44	45	45	45	43
	Median Age	43	44	43	44	43
Total	# of Deaths	1,141	1,200	1,087	1,061	914
	Mean Age	45	44	44	45	43
	Median Age	42	42	41	43	42

* Suicide deaths due to poisoning are excluded from the OTR definition of trauma. However, all suicide deaths are reported in this Table to provide a complete picture of suicide in the province.

Note: This table excludes cases where sex is not documented.

**MECHANISM OF SUICIDE* (INCLUDING POISONING)
BY SEX AND AGE GROUP, 2000-2001**

	<1	1-4	5-9	10-14	15-19	20-24	25-34	35-44	45-54	55-64	65-74	75-84	85+	UNK	Total	%**
Hanging																
Males	1	0	0	5	20	22	55	55	45	21	14	6	5	0	249	27.3
Females	0	0	0	4	9	6	11	7	9	5	10	3	0	0	64	7.0
Total	1	0	0	9	29	28	66	62	54	26	24	9	5	0	313	34.3
Firearm Injury																
Males	0	0	0	0	10	6	18	25	31	18	7	7	2	0	124	13.6
Females	0	0	0	0	0	0	0	3	0	0	0	0	0	0	3	0.3
Total	0	0	0	0	10	6	18	28	31	18	7	7	2	0	127	13.9
Drugs & Alcohol																
Males	0	0	0	0	1	3	12	28	29	10	5	1	1	0	90	9.9
Females	0	0	0	0	1	3	9	19	19	8	3	4	0	0	66	7.2
Total	0	0	0	0	2	6	21	47	48	18	8	5	1	0	156	17.1
Vehicle Exhaust																
Males	0	0	0	0	2	4	14	25	19	4	1	3	0	0	72	7.9
Females	0	0	0	0	1	0	1	3	2	1	0	0	0	0	8	0.9
Total	0	0	0	0	3	4	15	28	21	5	1	3	0	0	80	8.8
Jumping																
Males	0	0	0	0	1	7	14	11	7	4	4	5	1	0	54	5.9
Females	0	0	0	0	1	5	9	5	6	2	0	0	0	0	28	3.1
Total	0	0	0	0	2	12	23	16	13	6	4	5	1	0	82	9.0

**MECHANISM OF SUICIDE* (INCLUDING POISONING)
BY SEX AND AGE GROUP, 2000-2001**

	<1	1-4	5-9	10-14	15-19	20-24	25-34	35-44	45-54	55-64	65-74	75-84	85+	UNK	Total	%**
Drowning																
Males	0	0	0	0	1	0	4	9	4	1	4	2	0	0	25	2.7
Females	0	0	0	0	0	0	0	2	1	1	3	0	0	0	7	0.8
Total	0	0	0	0	1	0	4	11	5	2	7	2	0	0	32	3.5
All Other																
Males	0	0	0	0	2	11	21	23	20	7	3	6	2	0	95	10.4
Females	0	0	0	0	2	1	5	6	7	5	1	0	1	0	28	3.1
Total	0	0	0	0	4	12	26	29	27	12	4	6	3	0	123	13.5
Total																
Males	1	0	0	5	37	53	138	176	155	65	38	30	11	0	709	77.7
Females	0	0	0	4	14	15	35	45	44	22	17	7	1	0	204	22.3
Total	1	0	0	9	51	68	173	221	199	87	55	37	12	0	913	100.0

* Mechanism of suicide is defined using Death Factors.

** Percents are of the total number of suicides for the year.

Note: 1 case is excluded because its "Environment Primary Indicator" is null.

Note: 6 cases have an unknown sex and were therefore excluded from this table.

**TRAUMA DEATHS BY AGE GROUP AND SEX
BY REGION, 2000-2001**

Age (Years)	Southwestern			Central South			Central West			Central East			Toronto			Eastern			North			Ontario		
	M	F	Total	M	F	Total	M	F	Total	M	F	Total	M	F	Total	M	F	Total	M	F	Total	M	F	Total
<1	0	1	1	1	0	1	4	1	5	2	2	4	3	0	3	0	1	1	0	0	0	10	5	15
1-4	2	0	2	1	1	2	3	1	4	7	3	10	2	1	3	2	1	3	1	2	3	18	9	27
5-9	4	1	5	2	0	2	3	3	6	1	3	4	2	0	2	5	3	8	2	1	3	19	11	30
10-14	8	5	13	4	1	5	1	4	5	6	4	10	1	1	2	7	1	8	3	5	8	30	21	51
15-19	23	11	34	11	5	16	19	7	26	22	6	28	15	3	18	21	1	22	32	17	49	143	50	193
20-24	24	8	32	15	3	18	33	7	40	21	5	26	23	9	32	20	4	24	21	8	29	158	44	202
25-34	44	7	51	24	5	29	44	10	54	38	10	48	45	11	56	35	10	45	45	11	56	277	64	341
35-44	44	21	65	33	6	39	30	11	41	40	9	49	51	16	67	43	12	55	43	7	50	286	83	369
45-54	45	18	63	30	8	38	35	9	44	36	14	50	48	10	58	35	8	43	45	8	53	274	75	349
55-64	25	13	38	16	3	19	18	7	25	25	14	39	27	18	45	35	10	45	34	15	49	180	80	260
65-74	28	12	40	15	17	32	23	17	40	31	20	51	33	17	50	22	11	33	22	9	31	174	105	279
75-84	35	35	70	27	31	58	27	35	62	36	38	74	57	47	104	36	13	49	28	14	42	246	213	459
85+	30	46	76	20	26	46	26	51	77	25	58	83	44	84	128	20	43	63	18	27	45	184	335	519
Unknown	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Total	312	178	490	199	106	305	266	163	429	290	186	476	351	217	568	281	118	399	294	124	418	1,999	1,095	3,094

Note: There are 20 cases with no documented sex. The sum of the regional totals does not equal the Ontario total due to missing data on the region of incident. Further, 1 case is excluded because its "Environment Primary Indicator" is null.

MOTOR VEHICLE TRAUMA DEATHS BY AGE GROUP AND SEX BY REGION, 2000-2001

Age (Years)	Southwestern			Central South			Central West			Central East			Toronto			Eastern			North			Ontario		
	M	F	Total	M	F	Total	M	F	Total	M	F	Total	M	F	Total	M	F	Total	M	F	Total	M	F	Total
<1	0	0	0	1	0	1	1	0	1	0	1	1	0	0	0	0	1	1	0	0	0	2	2	4
1-4	0	0	0	0	0	0	0	0	0	2	0	2	1	0	1	2	1	3	1	0	1	6	1	7
5-9	2	1	3	1	0	1	1	2	3	1	0	1	2	0	2	0	1	1	1	1	2	8	5	13
10-14	3	3	6	2	1	3	0	2	2	3	2	5	0	0	0	3	0	3	0	0	0	11	8	19
15-19	15	8	23	3	3	6	12	3	15	13	6	19	2	0	2	6	1	7	13	12	25	64	33	97
20-24	15	6	21	8	1	9	17	4	21	14	3	17	3	2	5	9	1	10	10	4	14	77	21	98
25-34	14	2	16	9	4	13	11	4	15	14	5	19	5	2	7	18	3	21	15	1	16	88	21	109
35-44	13	10	23	12	3	15	9	5	14	13	6	19	8	3	11	5	5	10	10	3	13	70	35	105
45-54	9	9	18	7	1	8	10	2	12	13	8	21	2	0	2	10	4	14	14	6	20	65	30	95
55-64	5	7	12	5	1	6	4	6	10	7	8	15	0	2	2	11	3	14	9	4	13	41	31	72
65-74	1	2	3	2	9	11	2	3	5	6	10	16	6	1	7	5	1	6	3	5	8	25	32	57
75-84	4	4	8	2	8	10	2	2	4	6	5	11	4	5	9	5	3	8	5	6	11	28	33	61
85+	2	4	6	0	1	1	2	0	2	3	2	5	4	1	5	3	1	4	1	0	1	16	9	25
Unknown	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Total	83	56	139	52	32	84	71	33	104	95	56	151	37	16	53	77	25	102	82	42	124	501	261	762

Note: There are 7 cases with no documented sex. The sum of the regional totals does not equal the Ontario total due to missing data on the region of incident.

**EXTERNAL CAUSES OF INJURY
BY SEX, 2000-2001**

E CODE	FEMALES			MALES			TOTAL			
	No.	%**	MEAN AGE	No.	%**	MEAN AGE	No.	%**	MEAN AGE	
	1,096	100	64.3	1,999	100	49.1	3,095	100	54.5	
E800-807	RAILWAY	3	0.3	60.3	7	0.4	34.1	10	0.3	42.0
E810-819	MOTOR VEHICLE TRAFFIC	259	23.6	46.3	482	24.1	38.9	741	23.9	41.4
E820-825	MOTOR VEHICLE NON TRAFFIC	2	0.2	36.5	19	1.0	30.5	21	0.7	31.0
E826	PEDAL CYCLE	1	0.1	*	11	0.6	38.1	12	0.4	37.2
E827-829	OTHER ROAD VEHICLE	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0
E830-838	WATER TRANSPORT	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0
E840-845	AIR AND SPACE TRANSPORT	1	0.1	*	6	0.3	58.7	7	0.2	60.6
E846-848	VEHICLE INCIDENTS NOT ELSEWHERE CLASSIFIED	0	0.0	0.0	3	0.2	34.3	3	0.1	34.3
E880-888	UNINTENTIONAL FALLS	552	50.4	83.2	505	25.3	73.3	1,057	34.2	78.5
E890-899	FIRE AND FLAMES	33	3.0	57.4	61	3.1	47.4	94	3.0	50.8
E900-902 &	NATURAL AND ENVIRONMENTAL FACTORS	16	1.5	64.5	26	1.3	56.0	42	1.4	59.2
E910	DROWNING	23	2.1	26.3	115	5.8	36.0	138	4.5	34.4

**EXTERNAL CAUSES OF INJURY
BY SEX, 2000-2001**

E CODE		FEMALES			MALES			TOTAL		
		No.	%**	MEAN AGE	No.	%**	MEAN AGE	No.	%**	MEAN AGE
		1,096	100	64.3	1,999	100	49.1	3,095	100	54.5
E913	SUFFOCATION	11	1.0	32.8	21	1.1	37.9	32	1.0	36.2
E916-928	OTHER INCIDENTS	17	1.6	50.0	90	4.5	43.9	107	3.5	45.3
E953-958	SUICIDE & SELF INFLICTED INJURY (EXCL.POISONINGS)	127	11.6	41.3	529	26.5	42.4	656	21.2	42.2
E960-961 &	HOMICIDE AND INJURY PURPOSELY INFLICTED	30	2.7	37.5	75	3.8	37.0	105	3.4	37.2
E983-988	UNDETERMINED WHETHER UNINTENTIONALLY OR PURPOSELY INFLICTED	21	1.9	51.1	49	2.5	47.3	70	2.3	48.5

Note: There are 20 cases with no documented sex.

* Data suppressed due to small cell sizes

** Denominator for percentage is column total.



taking health information further
à l'avant-garde de l'information sur la santé