

# Ecosystem Health

## Science-Based Solutions



**Canadian Tissue Residue Guidelines  
for the Protection of Wildlife  
Consumers of Aquatic Biota:  
Methylmercury**


**Report No. 1-4**



Environment  
Canada

Environnement  
Canada

Canada



Science-Based Solutions  
Solutions fondées sur la science

Prepared and published by  
National Guidelines and Standards Office  
Environmental Quality Branch  
Environment Canada  
Ottawa

March, 2002

ISSN 1497-2689  
ISBN 0-662-29602-8  
Cat. no. En1-34/2-2002E

Scientific Supporting Document

Canadian Tissue Residue Guidelines  
for the Protection of Consumers of Aquatic Life:  
Methylmercury

Report No. 1-4

© Her majesty the Queen in Right of Canada, represented by the Minister of the Environment, 2001.  
All rights reserved. Reproduction authorized if source is acknowledged. The reproduction must be  
presented within its proper context and must not be used for profit.

## NOTE TO READERS

The *Ecosystem Health: Science-based Solutions* series is dedicated to the dissemination of scientific knowledge, information and tools for monitoring, assessing, and reporting on ecosystem health to support Canadians in making sound decisions. Documents published in this series include the scientific basis, methods, approaches and frameworks for environmental guidelines and their implementation; monitoring, assessing, and rehabilitating environmental quality in Canada; and, indicator development, environmental reporting and data management. Issues in this series are published *ad libitum*.

This particular issue provides the scientific supporting information and rationale for the development of Canadian Tissue Residue Guidelines for the protection of consumers of aquatic life for methylmercury. The information in this document is current as of 2000, when the document was originally prepared. Minor revisions and editorial changes have been made for publication in 2002. For additional information regarding these guidelines, please contact:

Environment Canada  
Environmental Quality Branch  
National Guidelines and Standards Office  
351 Saint-Joseph Boulevard  
Hull, QC K1A 0H3

phone: 819-953-1550  
fax: 819-953-0461  
[ceqg-rcqe@ec.gc.ca](mailto:ceqg-rcqe@ec.gc.ca)  
<http://www.ec.gc.ca>

Canadian Water Quality Guidelines are approved by the Water Quality Guidelines Task Group of the Canadian Council of Ministers of the Environment (CCME). Environment Canada acts as the federal representative and serves as the technical secretariat to this Task Group. These guidelines are included as a 2001 update in the *Canadian Environmental Quality Guidelines*, which was published by the CCME in October of 1999. For CCME publications, please contact:

CCME Documents  
c/o Manitoba Statutory Publications  
200 Vaughan Street  
Winnipeg, MB R3C 1T5

tollfree: 800-805-3025  
fax: 204-945-7172  
[spccme@gov.mb.ca](mailto:spccme@gov.mb.ca)  
<http://www.ccme.ca>

This scientific supporting document is available in English and in French. A factsheet is also available in English under the title *Canadian Environmental Quality Guidelines* (CCME 1999). Ce document scientifique du soutien est disponible en français et en anglais. Un feuillet d'information est aussi disponible en français sous le titre *Recommandations canadiennes pour la qualité de l'environnement* (CCME 1999).

### Reference listing:

Environment Canada. 2002. Canadian Tissue Residue Guidelines for the Protection of Consumers of Aquatic Life: Methylmercury. Scientific Supporting Document. Ecosystem Health: Science-based Solutions Report No. 1-4. National Guidelines and Standards Office, Environmental Quality Branch, Environment Canada. Ottawa.

## **ACKNOWLEDGEMENTS**

Brenda Miskimmin is gratefully acknowledged for her major scientific contribution to this document. The Guidelines and Standards Division extends its appreciation to members of the CCME Water Quality Task Group: Bijan Aidun, Jerry Choate, Sam Ferris, Isabelle Guay, Francis Jackson, Haseen Khan, Clair Murphy, Doug Spry, Les Swain, Darrell Taylor, Gerry Whitley, and Dwight Williamson for their expert review of this document. We would like to also thank Erin Burns-Flett, C. Dumont, Gary Ironside, N.P. McEwen, Greg Mierle, Alain Plouffe, Leo-Guy de Repentigny, Mark Richardson, Luke Trip, Rudy Wagemann, Kim Westcott, and Chris Wren for supplying valuable information. Elizabeth Roberts and Susan Roe provided much appreciated technical and editorial support.

**TABLE OF CONTENTS**

|   |      |
|---|------|
| NOTE TO READERS .....   | I    |
| ACKNOWLEDGEMENTS.....   | II   |
| TABLE OF CONTENTS .....   | III  |
| LIST OF TABLES .....  | V    |
| LIST OF FIGURES .....   | VI   |
| LIST OF APPENDICES.....   | VII  |
| ABSTRACT.....   | VIII |
| RÉSUMÉ .....  | X    |
| 1.0 INTRODUCTION .....  | 1    |
| 2.0 PHYSICAL AND CHEMICAL PROPERTIES .....  | 2    |
| 2.1 Properties.....   | 2    |
| 2.2 Sample Collection and Analytical Methods.....   | 3    |
| 2.2.1 Water sample collection.....  | 3    |
| 2.2.2 Analysis of Total Mercury and Methylmercury in Water .....                            | 3    |
| 2.2.3 Analysis of Methylmercury in Sediment and Tissue Samples .....                        | 4    |
| 3.0 PRODUCTION AND USES .....   | 6    |
| 3.1 Mining of Mercury .....   | 6    |
| 3.2 Uses.....   | 6    |
| 3.2.1 Use Reduction .....   | 8    |
| 4.0 SOURCES AND PATHWAYS INTO THE ENVIRONMENT.....  | 11   |
| 4.1 Natural Sources and Pathways .....  | 11   |
| 4.2 Anthropogenic Sources.....  | 12   |
| 5.0 ENVIRONMENTAL FATE AND BEHAVIOUR.....   | 17   |
| 5.1 The Mercury Cycle.....  | 17   |
| 5.2 Mercury Methylation and Demethylation .....   | 20   |
| 5.3 Factors Affecting Partitioning and Speciation .....                                     | 21   |
| 6.0 BIOACCUMULATION OF MERCURY AND METHYLMERCURY IN AQUATIC AND SEMI-AQUATIC ORGANISMS..... | 23   |
| 6.1 Aquatic Plants .....  | 23   |
| 6.2 Invertebrates .....   | 25   |
| 6.3 Freshwater Fish .....   | 27   |
| 6.4 Accumulation by Semi-Aquatic Animals and Birds .....                                    | 30   |
| 7.0 OVERVIEW OF DISTRIBUTION AND LEVELS OF METHYLMERCURY IN THE CANADIAN ENVIRONMENT.....   | 33   |
| 7.1 Aquatic Plants (App. B-1).....  | 33   |
| 7.2 Aquatic Invertebrates (App. B-2) .....  | 34   |
| 7.3 Fish (App. B-3).....  | 35   |
| 7.3.1 Freshwater fish (App. B-3a).....  | 35   |
| 7.3.2 Marine fish (App. B-3b).....  | 38   |
| 7.4 Amphibians (App. B-4).....  | 38   |
| 7.5 Reptiles (App. B-4).....  | 39   |
| 7.6 Mammals .....   | 39   |
| 7.6.1 Semi-aquatic Mammals (App. B-5).....  | 39   |
| 7.6.2 Marine Mammals (App. B-6).....  | 40   |
| 7.7 Birds (App. B-7) .....  | 41   |
| 7.8 Non-biological media (App. B-8).....  | 43   |
| 8.0 TOXICITY OF METHYLMERCURY TO PISCIVOROUS ORGANISMS.....                                 | 45   |
| 8.1 Mode of Toxicity.....   | 45   |
| 8.2 Toxicity to Mammals .....   | 47   |

## Canadian Tissue Residue Guidelines: Methylmercury

|        |   |    |
|--------|---|----|
| 8.2.1  | Acute and Chronic Toxicity (Appendix C-1)                     | 47 |
| 8.2.2  | Reproductive and Developmental Toxicity (Appendix C-2)        | 51 |
| 8.2.3  | Carcinogenic and Mutagenic Effects (Appendix C-3)             | 51 |
| 8.3    | Toxicity to Avian Species                                     | 53 |
| 8.3.1  | Acute and Chronic Toxicity                                    | 54 |
| 8.3.2  | Reproductive and Developmental Toxicity                       | 54 |
| 8.4    | Toxicity to Piscivorous Fish                                  | 57 |
| 9.0    | EXISTING TISSUE RESIDUE GUIDELINES                            | 59 |
| 9.1    | Human Health  | 59 |
| 9.2    | Wildlife  | 60 |
| 10.0   | CANADIAN TISSUE RESIDUE GUIDELINES FOR METHYLMERCURY          | 63 |
| 10.1   | Protocol  | 63 |
| 10.2   | Guideline Derivation and Rationale                            | 64 |
| 10.2.1 | Avian Species   | 64 |
| 10.2.2 | Mammalian Species   | 65 |
| 10.3   | Methylmercury Guideline                                       | 67 |
| 11.0   | DATA GAPS   | 71 |
| 12.0   | GUIDELINE IMPLEMENTATION                                      | 72 |
| 12.1   | Use of Appropriate Wildlife Species                           | 72 |
| 12.2   | Application of TRG to Appropriate Size Class of Dietary Items | 72 |
| 12.3   | Trophic Level Considerations                                  | 73 |
| 12.3.1 | Uncertainties with Establishing Distinct Trophic Levels       | 73 |
| 12.3.2 | Basic Aquatic Trophic Levels                                  | 73 |
| 12.3.3 | Prey Trophic Levels of Representative Species                 | 74 |
| 13.0   | REFERENCES  | 80 |

**LIST OF TABLES**

|           |  |    |
|-----------|--|----|
| Table 1.  | Physical and chemical properties of mercury. ....  | 5  |
| Table 2.  | Worldwide production of mercury from ores and concentrates in metric tonnes... 9   |    |
| Table 3.  | Metric tonnes of mercury used per year in various categories in Canada and the United States (OECD 1994). ....   | 10 |
| Table 4.  | Estimates of Fluxes of Mercury to the Global Atmosphere .....  | 11 |
| Table 5.  | Estimates of worldwide emissions of mercury (metric tonnes per year) to the atmosphere, soil, and water in 1983 (from Nriagu and Pacyna 1988 and Nriagu 1989).....   | 15 |
| Table 6.  | U.S. nation-wide discards <sup>a</sup> (tonnes per year) of mercury in products in municipal solid wastes (MSW), 1970 to 2000 .....  | 16 |
| Table 7.  | Variables associated with mercury concentrations in fish .....   | 30 |
| Table 8.  | Frequency of clinical signs and symptoms of the Minimata Bay disease where humans consumed fish and shellfish containing 10-35 µg·g <sup>-1</sup> mercury for an extended period. ....   | 47 |
| Table 9.  | Consumption parameters for mink, otter, kingfisher, loon, osprey, and eagle ....   | 58 |
| Table 10. | Worldwide survey of mercury tissue residue guidelines (concentrations or daily limits) for mercury in fish and fisheries products.....   | 62 |
| Table 11. | TRG derivation and calculations.....   | 68 |
| Table 12. | Reference concentrations (RCs; µg·kg <sup>-1</sup> diet, ww) <sup>a,b</sup> for Canadian wildlife species derived from the lowest avian (31 µg·kg <sup>-1</sup> bw per day) or mammalian (22 µg·kg <sup>-1</sup> bw per day) tolerable daily intakes (TDIs) for methylmercury and food intake (FI) to body weight (BW) ratios <sup>c</sup> ..... | 70 |
| Table 13. | Basic freshwater aquatic trophic levels.....   | 75 |
| Table 14. | Feeding habits and prey trophic levels of representative freshwater amphibian, reptilian, avian, and mammalian species.....  | 77 |

**LIST OF FIGURES**

Figure 1. Mercury emissions in Canada by sector for 1990 and 1995 as estimated by the Transboundary Air Issues Branch of Environment Canada. .... 106

Figure 2. Tonnes of mercury emitted by each Canadian province in 1990 as estimated by the Transboundary Air Issues Branch of Environment Canada. .... 107

Figure 3. Global and North American total mercury emissions for 1990 as reported by the Transboundary Air Issues Branch of Environment Canada. .... 108

Figure 4. The Mercury Cycle..... 109



**LIST OF APPENDICES**

Appendix A: Canadian strategies under a bilateral strategy to eventually eliminate anthropogenic toxic substances in the Great Lakes Basin. .... 110

Appendix B-1: Levels of total mercury and methylmercury in freshwater aquatic plants in the Canadian environment. .... 111

Appendix B-2a: Levels of total mercury and methylmercury in freshwater invertebrates in the Canadian environment. .... 116

Appendix B-2b: Levels of total mercury and methylmercury in marine invertebrates in the Canadian environment. .... 126

Appendix B-3a: Levels of total mercury and methyl mercury in freshwater fish in the Canadian environment. .... 127

Appendix B-3b: Levels of total mercury in marine fish in the Canadian environment. .... 144

Appendix B-4: Levels of total mercury and methylmercury in amphibians and reptiles in the Canadian environment. .... 145

Appendix B-5: Levels of total mercury and methylmercury in freshwater mammals in the Canadian environment. .... 147

Appendix B-6: Levels of total mercury and methylmercury in marine mammals in the Canadian environment. .... 152

Appendix B-7: Levels of total mercury and methylmercury in birds in the Canadian environment. .... 156

Appendix B-8: Levels of total mercury and methylmercury in selected non-biological media in the Canadian environment. .... 167

Appendix C-1: Summary of data on the acute or chronic toxicity of orally-administered methylmercury in mammals. .... 170

Appendix C-2: Summary of data on the reproductive and developmental toxicity of orally-administered methylmercury in mammals. .... 177

Appendix C-3: Summary of data on the carcinogenic and mutagenic effects of orally-administered methylmercury in mammals. .... 179

Appendix C-4: Summary of data on the acute, chronic or reproductive toxicity of methylmercury in birds. .... 180

Appendix D: Human consumption restriction categories for methylmercury in fish from Canadian waters. .... 185

## ABSTRACT

This document reviews the background information and rationale for the derivation of the Canadian Tissue Residue Guideline (TRG) for methylmercury (MeHg) for the protection of wildlife consumers of aquatic biota. It outlines information on the physical and chemical properties of MeHg, the sources and uses of mercury, production of MeHg, environmental fate and behaviour, bioaccumulation in biota, levels of MeHg in Canadian biota and various environmental media, and describes the toxicological effects of MeHg in mammals and birds. This abstract will briefly outline the information to be found in each chapter of the report. Tables are located either within the text or following each chapter, and all figures and Appendices are at the end of the report following the References.

To protect Canadian wildlife that consume fish or shellfish from any toxicological effects of MeHg, aquatic biota should contain no more than **33  $\mu\text{g}\cdot\text{kg}^{-1}$**  (0.033  $\text{mg}\cdot\text{kg}^{-1}$ ) MeHg on a wet weight basis.

The derivation and rationale for the wet-weight based TRG are fully described in chapter 10. It is important to note that the TRG was derived using the reference concentration (RC) of the Wilson's storm petrel (susceptible to MeHg bioaccumulation because it consumes almost its entire body weight in aquatic biota per day). In regions where this species is not found, RC values for resident wildlife may be applied (Table 12). The use of these other RC values will result in allowable concentrations that are higher than the TRG. **Monitoring programs in Canada typically measure mercury concentrations in large sport fish that generally contain higher levels than small fish (<35 cm), which are consumed by wildlife.** As such, mercury levels reported herein are biased toward those for large fish and may not represent actual risks to wildlife.

Mercury can exist in many forms, both inorganic and organic. Elemental mercury ( $\text{Hg}^0$ ) is very volatile, and is the only metal that is liquid at room temperature. The most toxicologically relevant form of mercury is MeHg ( $\text{CH}_3\text{Hg}$ ). Methylmercury is produced through the biological and chemical methylation of inorganic mercury. It is a potent neurotoxicant for animals and humans. Methylmercury is not very lipid soluble compared to many organochlorine contaminants, but it binds strongly with sulphhydryl groups in proteins and is therefore readily accumulated and retained in biological tissues (Clarkson 1994).

There are many sources of mercury to the environment, both natural and human related. Natural sources include geological mercury deposits, forest fires and other wood burning activities, volcanoes, and volatilisation from the oceans. Humans may accelerate the natural weathering process of mercury by removing it from areas rich in geological deposit and introducing or re-introducing it to the mercury cycle. Consumer and industrial products, coal and other fossil fuel combustion, chlorine alkali processing, mercury waste in landfills or storage, waste incineration, cement manufacture, metal smelting, and other activities have all altered the natural Hg cycle.

While world production (mining) and use of mercury is declining, substantial quantities are still used in some countries for gold mining. Mercury has been used in dental amalgams, pesticides, thermometers, barometers, and electrical products such as dry-cell batteries, fluorescent lights, switches, and other control equipment. It is also used in substantial quantities

in the electrolytic preparation of chlorine and caustic soda (chlor-alkali industry) (AEP 1992).

Generally, the aquatic mercury cycle may be characterised by high atmospheric fallout, removal by deposition to sediments, re-emission to the atmosphere as elemental Hg (Hg<sup>0</sup>), methylation and demethylation transformation processes within the lake or watershed, and exchanges through tributaries and groundwater (runoff and seepage) (Hurley et al. 1994). Some portion of the total mercury (THg) pool is accumulated by biota.

Methylmercury has a high potential for bioaccumulation as well as biomagnification with increasing trophic levels. Mercury and MeHg may be taken up by aquatic organisms directly from water or through their diet. Uptake by animals directly from water is the result of dissolved mercury adsorption or absorption through the body surface and respiratory organs. Available data indicate that MeHg usually represents less than 50% of total mercury in freshwater aquatic plants. The proportion of MeHg (MeHg/THg) increases through the food web, being lowest in aquatic plants, intermediate in invertebrates, and highest in fish and birds.

Fish, and piscivorous mammals and birds accumulate most of their mercury body burdens through their diet. Mercury in fish is comprised almost exclusively of MeHg regardless of the composition of diet sources and exposure water. Methylmercury levels in wildlife consumers of fish are elevated relative to the fish or other aquatic biota they consume indicating its biomagnification potential. Selenium (Se) is known to reduce the accumulation of mercury and to exert a measure of protection against MeHg toxicity. The Hg-Se ratio has shown a tendency to be higher in freshwater fish than in marine fish, which may be an indication of the higher relative mercury levels in certain freshwater environments (Pelletier 1985).

With respect to animal uptake, about 95% of an ingested dose of MeHg is efficiently absorbed into the bloodstream from the gastrointestinal tract, whereas inorganic Hg is less readily absorbed (7 to 15%; Wolfe et al. 1998). Methylmercury distributes via the bloodstream to all parts of the body, readily crosses placental barriers, may enter the foetal brain, and is readily accumulated in growing animal fur, bird feathers or human hair. The accumulation of mercury in the brain is greater in mammals than in fish (Zillioux et al. 1993). Some early clinical signs of MeHg toxicity in mammals include constricted vision, loss of co-ordination and nervous tremors. Birds often suffer reproductive impairment without an indication of obvious neurological effects.

Many countries have mercury or MeHg guidelines to protect human consumers of fish and shellfish. Only the United States and Canada, however, are in the process of establishing guidelines to protect wildlife that consume fish and other aquatic biota. The US EPA "Mercury Report to Congress" is a series of comprehensive reviews of all aspects of mercury and MeHg in the environment, including the derivation of water concentrations and tolerable consumption levels to protect wildlife consumers of fish (US EPA 1997b,c).

## RÉSUMÉ

Le présent document expose les informations et les critères qui ont servi à élaborer la Recommandation canadienne visant les résidus dans les tissus (RRT) pour le méthylmercure (MeHg), dont l'objectif est de protéger les espèces fauniques qui consomment des organismes aquatiques. Il présente dans ses grandes lignes l'information sur les propriétés chimiques et physiques du MeHg, les sources et les utilisations du mercure, la production du MeHg, le devenir et le comportement du mercure dans l'environnement ainsi que sa bioaccumulation dans les biotes, et les concentrations de MeHg dans le biote canadien et divers compartiments de l'environnement; il décrit aussi les effets toxiques du MeHg chez les mammifères et les oiseaux. Le présent résumé décrit brièvement l'information présentée dans chaque chapitre du rapport. Certains tableaux ont été placés au fil du texte et d'autres, à la fin des chapitres; toutes les figures et annexes se trouvent à la fin du rapport, après la section des références bibliographiques.

Afin de protéger les espèces fauniques canadiennes qui consomment des poissons ou des crustacés et coquillages contre tout effet toxique du MeHg, les organismes aquatiques ne devraient pas contenir plus de **33  $\mu\text{g}\cdot\text{kg}^{-1}$**  (0,033  $\text{mg}\cdot\text{kg}^{-1}$ ) de MeHg (poids frais).

La méthode et les critères qui ont servi à élaborer la RRT basée sur le poids frais sont décrits en détail au chapitre 10. Il est important de remarquer que la RRT a été établie à partir de la concentration de référence (CR) de l'Océanite de Wilson (oiseau particulièrement susceptible de bioaccumuler le MeHg, étant donné qu'il consomme quotidiennement presque l'équivalent de son poids en organismes aquatiques). Dans les régions où cette espèce n'est pas présente, les CR des espèces résidentes peuvent être appliquées (tableau 12). L'utilisation de ces autres CR entraînera des concentrations admissibles plus élevées que la RRT. **Au Canada, les programmes de surveillance mesurent généralement les concentrations de mercure dans les grands poissons de sport, qui contiennent habituellement des concentrations plus élevées que les petits poissons (<35 cm) consommés par la faune.** Ainsi, les concentrations de mercure mentionnées dans le présent document correspondent plutôt à celles mesurées chez les grands poissons et peuvent n'être pas représentatives des risques réels pour la faune.

Le mercure peut exister sous plusieurs formes, tant inorganiques qu'organiques. Le mercure élémentaire ( $\text{Hg}^0$ ) est très volatil et est aussi le seul métal liquide à la température de la pièce. La forme de mercure la plus importante sur le plan toxicologique est le MeHg ( $\text{CH}_3\text{Hg}$ ). Le méthylmercure est produit par méthylation biologique et chimique du mercure inorganique. Il constitue un neurotoxique puissant pour les animaux et les humains. Le méthylmercure n'est pas très liposoluble par rapport à de nombreux contaminants organochlorés, mais il se lie solidement aux groupes sulfhydryles des protéines et est donc facilement accumulé et retenu dans les tissus biologiques (Clarkson, 1994).

Le mercure présent dans l'environnement provient de nombreuses sources naturelles et anthropiques. Les sources naturelles comprennent les gisements mercurifères, les feux de forêts et toute autre combustion de bois, les volcans et la volatilisation du mercure présent dans les océans. Les humains peuvent accélérer le processus naturel de mobilisation du mercure

sous l'effet de l'altération météorique des roches mercurifères, en le déplaçant des régions riches en gisements géologiques et en l'introduisant ou en le réintroduisant dans le cycle du mercure. Certains produits de consommation et de l'industrie, la combustion du charbon et d'autres combustibles fossiles, la production de chlore et de soude caustique, les déchets de mercure dans les décharges ou les entrepôts, l'incinération des déchets, les cimenteries et les fonderies sont au nombre des facteurs qui ont altéré le cycle naturel du mercure.

Bien que la production (exploitation minière) et l'utilisation mondiales du mercure diminuent, des quantités importantes sont toujours utilisées dans certains pays dans l'exploitation de l'or. Le mercure a été utilisé dans les amalgames dentaires, les pesticides, les thermomètres, les baromètres et divers produits électriques tels que les piles sèches, les tubes fluorescents, les interrupteurs et d'autres dispositifs de contrôle. Le mercure est également utilisé en quantités importantes dans la production électrolytique de chlore et d'hydroxyde de sodium (soude caustique) (AEP, 1992).

De façon générale, le cycle du mercure dans le milieu aquatique peut être caractérisé de la façon suivante : dépôt atmosphérique important, dépôt dans les sédiments, réémission dans l'atmosphère sous forme de mercure élémentaire ( $Hg^0$ ), méthylation et déméthylation dans le lac ou le bassin hydrographique, et transport dans les cours d'eau et les eaux souterraines (ruissellement et infiltration) (Hurley *et al.*, 1994). Une partie du mercure total (HgT) s'accumule dans le biote.

Le méthylmercure a un important potentiel de bioaccumulation, ainsi que de bioamplification d'un niveau trophique à l'autre. Le mercure et le MeHg peuvent être absorbés par les organismes aquatiques directement de l'eau, ou par leur alimentation. Dans le premier cas, le mercure dissous est adsorbé ou absorbé au niveau de la surface corporelle et des organes respiratoires. Les données disponibles indiquent que le MeHg représente habituellement moins de 50 % du mercure total présent dans les plantes aquatiques d'eau douce. La proportion de MeHg (MeHg/HgT) augmente d'un niveau à l'autre du réseau trophique : elle est au plus bas chez les plantes aquatiques, intermédiaire chez les invertébrés et au plus haut chez les poissons et les oiseaux.

Les poissons, les mammifères piscivores et les oiseaux accumulent la plus grande partie de leur charge corporelle de mercure par leur alimentation. Le mercure présent dans les poissons est presque exclusivement sous forme de MeHg, quelles que soient la composition du régime alimentaire des poissons et de l'eau dans laquelle ils se trouvent exposés au mercure. Les concentrations de méthylmercure dans les animaux piscivores sont élevées par rapport à celles mesurées chez les poissons ou d'autres organismes aquatiques qu'ils consomment, ce qui témoigne de son potentiel de bioamplification. On sait que le sélénium (Se) réduit l'accumulation du mercure et offre une protection contre la toxicité du MeHg. Le ratio Hg:Se tend à être plus élevé chez les poissons dulcicoles que chez les poissons marins, ce qui pourrait indiquer des concentrations relatives de mercure plus élevées dans certains milieux d'eau douce (Pelletier, 1985).

En ce qui a trait à l'absorption du mercure par les animaux, environ 95 % d'une dose ingérée de MeHg est efficacement absorbée par le tractus gastro-intestinal et se retrouve dans le sang, alors que le Hg inorganique est moins facilement absorbé (7 % à 15 %; Wolfe *et al.*, 1998). Le méthylmercure est distribué dans toutes les parties du corps par le sang, traverse facilement les barrières placentaires, peut pénétrer le cerveau du fœtus, et s'accumule facilement dans la fourrure, les plumes ou les cheveux humains durant leur croissance. L'accumulation de mercure dans le cerveau est plus importante chez les mammifères que chez les oiseaux (Zillioux *et al.*,

1993). Certains des premiers signes cliniques de l'intoxication par le MeHg chez les mammifères comprennent une vision réduite, une perte de la coordination et des tremblements nerveux. Chez les oiseaux, on observe souvent des problèmes de reproduction, sans signes évidents de troubles neurologiques.

De nombreux pays ont des lignes directrices pour le mercure ou le MeHg qui visent à protéger les personnes qui consomment du poisson et des coquillages et crustacés. Cependant, seuls les États-Unis et le Canada sont en train d'élaborer des lignes directrices visant à protéger les espèces fauniques qui consomment des poissons ou d'autres organismes aquatiques. Le « Mercury Report to Congress » de l'EPA des États-Unis présente des examens détaillés de tous les aspects du mercure et du MeHg dans l'environnement, dont les méthodes d'établissement des concentrations dans l'eau et des concentrations tolérables dans le biote permettant d'assurer la protection des animaux piscivores (US EPA, 1997b,c).

## 1.0 INTRODUCTION

Methylmercury ( $\text{CH}_3\text{HgX}$ , hereafter abbreviated as MeHg) is one of many forms of mercury that may be found in natural environments. It is approximately ten times more toxic than inorganic mercury to biota and is a potent neurotoxicant. Methylmercury is the form of mercury that is most readily accumulated by biota, and is only slowly excreted because of strong binding with proteins (Clarkson 1994).

As a mercury species, MeHg is not produced by humans, but rather it is produced a) biologically by micro-organisms, and b) chemically in natural aquatic environments under certain circumstances. Inorganic mercury occurs naturally in rocks all over the world. Humans cause an accelerated weathering of inorganic mercury when they extract it from the earth for their use, and may extend its life in the global mercury cycle when they re-use or incinerate mercury-containing products. Since MeHg is produced from inorganic mercury, high MeHg concentrations may be found where inorganic mercury occurs. Excessive inorganic mercury is released to the environment where humans use it during mining, ore smelting, fossil fuel and solid waste incineration, chlor-alkali plants (OECD 1994), and other uses that will be described in this report. Once released to the atmosphere, mercury may be carried hundreds or thousands of miles to be deposited in watersheds and surface waters. It is under certain aquatic conditions that MeHg is produced from the deposited or *in situ* inorganic mercury, and enters the food web.

Levels of mercury are higher in natural environments now and in recent history than in pre-industrial times. This has been demonstrated by changes in mercury flux rates in lake sediment cores collected in North America and Europe (e.g., Lucotte et al. 1995; Engstrom and Swain 1997; Landers et al. 1998). Recognising the environmental and health-related problems that mercury may cause, many governments have placed restrictions on human fish consumption because of mercury contamination, have banned or are phasing out the use of mercury in consumer and industrial products, and are curtailing industrial emissions of mercury (WHO 1990; OECD 1994; Environment Canada and US EPA 1997; US EPA 1997b,c).


Fish monitoring programs have shown that piscivorous fish may have elevated MeHg in their tissues. Likewise, avian and mammalian wildlife that are dependent on aquatic biota such as fish or shellfish are susceptible to accumulating toxic levels of MeHg (reviewed by Wolfe et al. 1998). This report will comprehensively examine the evidence that indicates that avian wildlife species exposed to excessive MeHg may incur reproductive and behavioural deficits, while mammalian piscivores are more likely to suffer neurological effects. Based on this evidence, a Canadian Tissue Residue Guideline (TRG) for MeHg was developed to protect piscivorous birds and mammals.

Development of the Canadian TRG for MeHg involved a review of the scientific literature on the production, uses, sources, and levels of MeHg in biota in Canada. The chemical properties, bioaccumulation, fate and behaviour, including the important processes of mercury methylation and demethylation were evaluated to characterise the types of sites in Canada most likely to develop MeHg contamination. Diet-based acute and chronic toxicity data for avian receptors such as loons, herons and eagles, and for piscivorous mammals such as mink and otter, were evaluated according to the protocol for deriving Canadian TRGs for the protection of wildlife consumers of aquatic biota (CCME 1998).

## 2.0 PHYSICAL AND CHEMICAL PROPERTIES

### 2.1 Properties

Mercury ( $\text{Hg}^0$ ) belongs to group IIb of the periodic table of elements, and is chemically quite different from the other two members of the group, Zn and Cd. For example, of the three metals, only Hg may become methylated in the environment to form alkylmercury compounds [such as mono methylmercury ( $\text{CH}_3\text{Hg}^+$ ) and dimethylmercury ( $(\text{CH}_3)_2\text{Hg}$ )] and aryl compounds (phenyl-mercury) (AEP 1992). The mechanism involves the nonenzymatic transfer of methyl groups ( $\text{CH}_3$ ) from methylcobalamin ( $\text{CH}_3\text{B}_{12}$ ) to  $\text{Hg}^{2+}$  (Robinson and Tuovinen 1984), a process that will be described in more detail in chapter 5:



Unlike the other metals, mercury has a strong tendency toward covalent interactions, and has a high chemical affinity for thiol (S-H) groups. Compared to cadmium or zinc, covalent complexes of  $\text{Hg}^{2+}$  are many orders of magnitude more stable. For example, the stability constants for  $\text{HgCl}_2$ ,  $\text{CdCl}_2$  and  $\text{ZnCl}_2$  are  $10^{16}$ ,  $10^3$  and  $10^{-1}$ , respectively (AEP 1992). Unlike mercury, cadmium and zinc compounds are not volatile nor are they reduced or methylated by micro-organisms.

Mercury has a very low melting point ( $-37^\circ\text{C}$ ) compared with cadmium ( $321^\circ\text{C}$ ) and zinc ( $420^\circ\text{C}$ ). Elemental mercury ( $\text{Hg}^0$ ) is the only metal that is liquid at room temperature. It has a high surface tension, forming spherical droplets when the liquid is released. It is also of low solubility in water, making precipitation and aquatic systems a poor sink for  $\text{Hg}^0$ . Elemental Hg is volatile; with volatility increasing with temperature. Its high surface tension and uniform volume of expansion make mercury ideal for use in thermometers, barometers and other measuring devices (Smith and Rowan-West 1996).

Mercury can theoretically exist in various valence states (0, +1, and +2), and forms inorganic as well as organic compounds. The chemical compounds of Hg (II) are much more numerous than those of Hg (I) (OECD 1994). The inorganic forms include the very stable, and relatively water-soluble halides,  $\text{HgCl}_2$ ,  $\text{HgBr}_2$ , and  $\text{HgI}_2$ , plus the water-soluble, dissociated form,  $\text{HgF}_2$  (AEP 1992). Among the inorganic anions,  $\text{Hg}^{2+}$  forms the strongest covalent bond with chloride ions, so mercuric-chloride species may dominate when chloride salts are abundant (AEP 1992).

In addition to simple salts, such as chloride, nitrate and sulphate, mercury (II) forms an important class of organometallic compounds. These are characterised by the attachment of mercury to either one or two carbon atoms to form compounds of the type  $\text{RHgX}$  and  $\text{RHgR}'$  where R and R' represent the organic moiety (most commonly  $\text{CH}_3$ ). The most numerous are those of the type  $\text{RHgX}$ , where X may be an anion such as chloride ( $\text{Cl}^-$ ), sulphide, or sulphate ( $\text{SO}_4^-$ ) (WHO 1989).

The most toxicologically relevant form of mercury is methylmercury ( $\text{CH}_3\text{Hg}$ ). Hg is easily transported across cell membranes, resulting in toxicity to biota. Of the mercury compounds, methylmercury has the highest octanol-water partition coefficient ( $K_{ow} = [\text{Hg}]_{\text{octanol}}/[\text{Hg}]_{\text{water}}$ ;  $\log K_{ow}$  1.7-2.5; Table 1), which is an indication of its affinity for body fats. While this  $\log K_{ow}$  is moderate relative to many organochlorine substances (e.g., toxaphene 6.44; DDT compounds



>5.5), it is the combination of lipophilicity and especially its binding with sulphhydryl groups in proteins which fosters retention of MeHg (Saouter et al. 1993).

## 2.2 Sample Collection and Analytical Methods

Until the last decade or so, many mercury measurements in water samples were artificially high because of sample contamination (Driscoll et al. 1994). For example, mercury concentrations in water from Vandercook Lake, Wisconsin, saw an apparent decrease from over 200 ng·L<sup>-1</sup> in 1983 to ~50 ng·L<sup>-1</sup> in 1985 to 0.5 ng·L<sup>-1</sup> in 1986 as investigators used progressively cleaner techniques for sample collection and handling (Fitzgerald and Watras 1989). Coincidentally, formerly undetectable levels of mercury and MeHg (such as in filtered water samples) are now commonly reported because of improvements in analytical techniques. Notably, the detection limits for MeHg are often in the pg·L<sup>-1</sup> range (Lee 1987; Bloom 1989). At the Third International Conference on “Mercury as a Global Pollutant” in July 1994, an entire session was devoted to mercury collection and measurement methods [see Vol. 80 (1-4), Water Air Soil Pollut. 1995].

### 2.2.1 Water sample collection

The use of trace-metal clean collection techniques is increasingly being recognised as necessary for samples used for mercury analyses. Typically, Teflon bottles are first washed with hot HNO<sub>3</sub>, filled with a 1% ACS grade HCl solution, and stored in new sealable polyethylene bags (St. Louis et al. 1994). Bottles should be transported to the field within clean containers that may also be placed inside plastic bags. All handlers must wear new polyethylene gloves. At the site, the HCl solution is drained and the bottle rinsed three times with sample water, filled, and then placed back into the bags. If a sample pump is used, acid-rinsed Teflon tubing should be used throughout. Within 2 h of sample collection, a small quantity (250 µL per 250 mL water) of concentrated *trace metal* grade HCl is added as a preservative (St. Louis et al. 1994).

Unfiltered samples may then be used for mercury analysis. For dissolved mercury analyses, filtering should be done with either an acid-cleaned glass (Ramamoorthy 1982) or Teflon filtration apparatus using acid-cleaned polycarbonate filters (e.g., 0.4 µm nucleopore filters; Gill and Bruland 1990; Hurley et al. 1995).

### 2.2.2 Analysis of Total Mercury and Methylmercury in Water

The methods of analysis used worldwide for mercury and its compounds are too numerous to detail in this report, however, the recently developed and widely preferred method will be detailed. Descriptions of some other methods in use are tabulated from pages 248 to 255 in US DOH and HS (1994).

Bloom (1989) and Watras et al. (1995)<sup>1</sup> analysed for total mercury (THg), dissolved mercury, MeHg, plus Hg<sup>0</sup>. They used a cryogenic gas chromatograph with a highly sensitive cold vapour atomic fluorescence detector (CVAFS). Methylmercury is determined by aqueous-phase ethylation followed by purging of volatile methyl ethyl mercury onto graphitized carbon black (Carbotrap). Hg<sup>0</sup> is determined by sparging 500 mL of unfiltered, untreated water for 20 min with Hg-free N<sub>2</sub> and trapping the volatile Hg on gold. Methylmercury and Hg<sup>0</sup> are both thermally desorbed from their respective “traps” prior to CVAFS detection. Detection limits range from

---

<sup>1</sup> Both of these first authors were directly involved in developing US EPA Method 1631 (1996).

## Canadian Tissue Residue Guidelines: Methylmercury

0.01 to 0.06 ng·L<sup>-1</sup> for THg, from 0.01-0.02 ng·L<sup>-1</sup> for MeHg, and from 0.004 to 0.02 ng·L<sup>-1</sup> for Hg<sup>0</sup> (Watras et al. 1995).

An excerpted summary of this recent standard technique developed by scientists for the US EPA (1996) follows. This method is increasingly being used in Canada, such as at the Freshwater Institute (DFO) in Winnipeg (St. Louis et al. 1994).

### Summary of EPA Method 1631 (US EPA 1996).

A 100–2000 mL sample is collected directly into specially cleaned, pretested, fluoropolymer bottle(s) using sample handling techniques specially designed for collection of mercury at trace levels (US EPA 1995a).

For dissolved Hg, samples are filtered through a 0.45-µm capsule filter.

Samples are preserved by adding 5 mL·L<sup>-1</sup> of pretested 12 N HCl (to allow both total and methyl Hg determination) or 5 mL·L<sup>-1</sup> BrCl solution, if total mercury only is to be determined.

Prior to analysis, a 100 mL sample aliquot is placed in a specially designed purge vessel, and 0.2 N BrCl solution is added to oxidize all Hg compounds to Hg(II).

After oxidation, the sample is sequentially prerduced with NH<sub>2</sub>OH·HCl to destroy the free halogens, and then reduced with SnCl<sub>2</sub> to convert Hg(II) to volatile Hg(0).

The Hg(0) is separated from solution by purging with nitrogen onto a gold-coated sand trap.

The trapped Hg is thermally desorbed from the gold trap into an inert gas stream that carries the released Hg(0) into the cell of a cold-vapor atomic fluorescence spectrometer (CVAFS) for detection.

### 2.2.3 Analysis of Methylmercury in Sediment and Tissue Samples

The analysis for tissue or sediment samples would be the same as for water samples except the sample is either 1) digested with a mixture of hot (180°C) nitric and sulphuric acids, or 2) treated with 60°C nitric and sulphuric acids and oxidized with a solution of potassium permanganate, with excess permanganate titrated with 30% peroxide, or 3) microwave digested in the presence of nitric acid. In all cases, the oxidized mercury is converted to the elemental state with a reducing solution containing stannous chloride (or stannous sulphate), hydroxylamine sulphate, and sodium chloride (Hendzel and Jamieson 1976; Scruton et al. 1994). The mercury is partitioned into vapour and determined by GC/CVAA or GC/CVAFS as described above.

A recent modification to these procedures is that methylmercury is first distilled from all matrix types into a Teflon vessel, ethylated with sodium tetraethyl borate and sorbed onto a Tenax (Enka Research Institute, Anaheim CA) resin column. Methylmercury is then separated using gas-liquid chromatography, pyrolyzed, and detected as Hg<sup>0</sup> using CVAFS (Liang et al. 1994).

The Canadian Food Inspection Agency, which is responsible for monitoring mercury concentrations in fish for domestic use and for export, analyses tissue for total mercury with the assumption that essentially all mercury in fish tissue is methylmercury (E. Burns-Flett 1999, DFO, pers. com.).

**Table 1. Physical and chemical properties of mercury.**

| Property <sup>a</sup>                         | Mercuric Hg (II)<br>(HgCl <sub>2</sub> ) | Methylmercury<br>(CH <sub>3</sub> Hg <sup>+</sup> )       | Elemental Mercury<br>(Hg <sup>0</sup> )                              |
|---|--|---|--|
| Atomic number                                 | --                                       | --  | 80   |
| Molecular Weight                              | 271.5                                    | 215.63  | 200.59   |
| Boiling point                                 | --                                       | Hg(CH <sub>3</sub> ) <sub>2</sub> :<br>92°C/98.66 kPa (4) | 356.9°C/<br>101.32 kPa (1)   |
| CAS No.                                       | 7487-94-7                                | 22967-92-6  | 7439-97-6  |
| Conductivity                                  | --                                       | --  | 0.022 cal·s <sup>-1</sup> ·cm <sup>-3</sup> ·<br>C <sup>-1</sup> (1) |
| Density (spec. gravity)                       | 5.4 (4)                                  | Hg(CH <sub>3</sub> ) <sub>2</sub> :<br>3.1874 (4)         | 13.534 g·cm <sup>-3</sup> at<br>25°C (4)                             |
| Melting point                                 | --                                       | --  | 38.87°C  |
| K <sub>OC</sub> (organic carbon) <sup>b</sup> | Not avail.                               | DOC increases<br>solubility (6)                           | Not avail.   |
| log K <sub>sed</sub>                          | 3.4-4.1 (3)                              | 2.9-3.2 (reduced by<br>DOC in water; 6)                   | --   |
| log K <sub>susp.part.</sub>                   | 5.35 (2); 4.8-<br>4.9 (10)               | 5.73(2)   | --   |
| log K <sub>ow</sub>                           | 0.61 (5)                                 | 1.7(7)-2.54(5)  | --   |
| Surface tension                               | --                                       | --  | 475 dynes·cm <sup>-1</sup> (1)                                       |
| Vapour pressure, Pa                           | 1.87·10 <sup>-2</sup><br>@ 34°C (4)      | 1.13 @ 20°C MeHgCl<br>(11)                                | 0.16 @ 20°C (11)<br>2·10 <sup>-3</sup> at 25°C (10)                  |
| Water Solubility                              | 69 g·L <sup>-1</sup> (4)                 | 1 µg·L <sup>-1</sup> (8);                                 | 25-30 µg·L <sup>-1</sup> (9)   |

Numbers in parentheses refer to the following references: (1) AEP 1992; (2) Hurley et al. 1989; (3) Hurley et al. 1994; (4) RSC 1994; (5) Halbach 1985; (6) Miskimmin 1991; (7) Major et al. 1991; (8) Eisler 1987; (9) Krenkel 1974; (10) Hurley et al. 1995; and (11) WHO 1990.

Notes:

<sup>a</sup> With the exception of molecular weight, most of these properties are dependent on pH, temperature, carbon content and other characteristics, and are given as general values only.

<sup>b</sup> K refers to a partition coefficient of a substance between indicated phase:water (Hg concentrations in each medium). For example, K<sub>ow</sub>=octanol/water partition coefficient; K<sub>susp.part.</sub>=suspended particle/water; K<sub>sed.</sub>=sediment/water; K<sub>oc</sub>=organic carbon/water.

### **3.0 PRODUCTION AND USES**

#### **3.1 Mining of Mercury**

Mercury cannot be created or destroyed. Human activities introduce excess mercury into the global mercury cycle that would otherwise have remained buried. The mining of mercury essentially results in an accelerated weathering process, by which much more mercury than normal is released from rocks.

Relatively few countries are involved in the production of mercury from mining cinnabar or other deposits. Important international mining locations include Algeria, Russia, China, Finland, Spain, and Mexico (OECD 1994; Table 2). Canada has no mercury mines that are currently active. Two mines were formerly active in central British Columbia along the Pinchi Fault zone. The Pinchi Mine was operational in the early 1940s and briefly in the early 1970s, and the Bralorne Takla Mine was operational for one year in the 1940s (Armstrong 1946; A. Plouffe 1999, GSC, pers. com.).

Mercury production from mining has declined steadily in the last 20 years (OECD 1994). Worldwide production decreased from 10 000 t in 1970 to 5 500 t in 1990. There is significant secondary production of mercury, which includes recycling, recovery, and industrial reprocessing of mercury. In the United States, secondary industrial production varied between 10 and 20% of total mercury consumption in 1985-89 (Minerals Yearbook 1989).

From 1972 to 1984, Mexico was the main Hg supplier to the largest gold mining nation, Brazil who requires about 340 t of Hg per year mainly for its gold mining processes (Veiga et al. 1995). Worldwide, the total number of gold miners using mercury amalgamation to extract gold in the early 1990s was estimated to be  $4 \pm 1$  million persons (Brazil 650 000; Tanzania 250 000; Indonesia 250 000; Vietnam 150 000) (Jernelöv and Ramel 1994).

Stockpiles of mercury in the United States are maintained for strategic defence purposes by the National Defense Stockpile (NDS) and the Department of Energy (DOE). NDS legislation enacted in 1992 authorised the disposal of the entire mercury stock from the NDS. At the end of 1993, there was an estimated 4466 t of mercury in the NDS groups (US Department of Commerce 1993).

#### **3.2 Uses**

Mercury has been used in dental amalgams, pesticides, thermometers, barometers, and electrical products such as dry-cell batteries, fluorescent lights, switches, and other control equipment. It is also used in substantial quantities in the electrolytic preparation of chlorine and caustic soda (chlor-alkali industry) (AEP 1992). Smaller quantities are used in industrial catalyst manufacture, general laboratory use, and pharmaceuticals. (US EPA Factsheet; [www.epa.gov/OGWDW](http://www.epa.gov/OGWDW))

According to an Organisation for Economic Co-operation and Development (OECD) questionnaire submitted to member countries (including Canada) the general range of consumption patterns for all countries for 1988-92 were as follows:

- batteries, range 3 to 69% (average 25%);
- chlor-alkali industry, range 2 to 78% (average 28%);

## Canadian Tissue Residue Guidelines: Methylmercury

- electrical equipment and measurement equipment, range 1 to 50% (average 16%);
- paint, range 0.1 to 16% (average 10%);
- tooth fillings, range 2 to 51% (average 7%);
- other uses, such as thermometers and laboratory uses (average 14%) (OECD 1994).

Fehr and Dennis (1975) estimated that 75% of seeds for cereal crops in Alberta were formerly treated with mercury compounds and that up to 120 t of organic mercury compounds were applied to Alberta and Saskatchewan soils from 1930 to 1972. During this period, seed-eating and predatory birds, as well as their eggs, were recognised to have elevated methylmercury concentrations that originated from the fungicide treatments of the seeds. Some of the mercury used in the 95 agricultural products that were registered for use may still be a component of the mercury cycle. By the early 1990s, there were four remaining mercurial fungicides registered for use in Canada, all of which were for turf application (Sang and Lourie 1997). Since December 31 1995, no further registration of mercury-containing fungicides was permitted although the remaining retail stocks could be used until depleted. These retail stocks were estimated to be depleted by the end of 1997.

According to the OECD survey, Canada used an average of 56.7 t Hg·a<sup>-1</sup> from 1982 to 1991. Of this amount 24 t was reported to be used in measurement equipment, 10 t in dental amalgams, with approximately 4 t for each of batteries, thermometers, electrical switches, and in laboratory uses. Chlor-alkali industries, paints, and pesticides were each reported at approximately 2 t Hg·a<sup>-1</sup> in Canada (OECD 1994; Table 3). A summary of Canadian mercury use by sector is also found in Pilgrim and EMAN (1998).

The remaining mercury cell chlor-alkali plants in Canada are regulated federally by the Chlor-Alkali Mercury Liquid Effluent Regulation<sup>2</sup> and the Chlor-Alkali Mercury Release Regulations<sup>3</sup> with respect to the release of mercury to the environment. Chlor-alkali plants produce chlorine and caustic soda (which is used by the pulp and paper industry) by passing a brine solution over mercury cells. A status report prepared by Environment Canada summarised the loss of mercury from mercury cell chlor-alkali plants to effluents, emissions, products, and solid wastes for the five mercury cell plants that were operating in Canada from 1986-1989. During 1986-1989, total mercury losses to effluent decreased from 88.07 kg·a<sup>-1</sup> (1986) to 46.29 kg·a<sup>-1</sup> (1989). Total mercury emissions over the four year period decreased from 680.56 kg·a<sup>-1</sup> (1986) to 547.18 kg·a<sup>-1</sup> (1989). Total mercury losses to products and solid wastes for all plants during 1986-1989 were 302.64 kg and 3318.13 kg, respectively (Paine 1994).

In the U.S., chlor-alkali plants used the most mercury (381 t), followed by batteries (250 t), paint (192 t), electrical switches (141 t) and measurement equipment (87 t; OECD 1994). These figures were as of 1989, and were reported in the 1995 OECD survey (Table 3).

Because mercury is an important long-range atmospheric pollutant, the decline in use of mercury by the U.S. (former world's largest Hg user; 1214 t used in 1989), may have important long term implications for mercury cycling in Canada. In the U.S., the use of mercury-containing products in 1990 was reduced by a factor of 3.6 compared to the maximum in 1968. One important reason for this decline is that mercury is no longer used in agriculture, amalgamation, the pulp and paper

---

<sup>2</sup> Chlor-Alkali Mercury Liquid Effluent Regulations, C.R.C., c.811.

<sup>3</sup> Chlor-Alkali Mercury Release Regulations, SOR/90-130

industry, and the pharmaceutical industry. Electrical equipment, batteries and the chlor-alkali industry each account for one third of consumption. Paint accounted for up to 18% of mercury consumption until 1990, when a decrease occurred following new restrictions on mercury use in paints. Battery use showed a substantial downward trend from 1989 to 1990, whereas other uses showed smaller changes since 1984 (OECD 1994).

### **3.2.1 Use Reduction**

Mercury and its compounds are now ranked as Level I persistent toxic substances on a list prepared by Environment Canada and the United States Environmental Protection Agency, in a bilateral strategy to eventually eliminate anthropogenic toxic substances in the Great Lakes Basin (Environment Canada and US EPA 1997). As such, goals concerning reductions in the use of mercury compounds have been set by each country.

The U.S. goal is to seek by the year 2006, a 50% reduction nationally in the deliberate use of mercury and a 50% reduction in the release of mercury from sources resulting from human activity. The release challenge will apply to the aggregate of releases to the air nation-wide and of releases to the water within the Great Lakes Basin. This target is considered as an interim reduction target and, in consultation with stakeholders, will be revised if warranted, following circulation of the Mercury Study Report to Congress (US EPA 1997b,c).

The Canadian goal is to seek by the year 2000, a 90% reduction in the release of mercury, or where warranted, the use of mercury, from polluting sources resulting from human activity in the Great Lakes Basin. This target is considered as an interim reduction target and, in consultation with stakeholders in the Great Lakes Basin, will be revised if warranted, following completion of the 1997 COA (Canada-Ontario Agreement Respecting the Great Lakes Basin Ecosystem, established in 1994).

Activities by Canadian companies in the Great Lakes Basin to date have resulted in significant reductions in mercury content in batteries (60 – 90%), fluorescent lamps (44%) and switches, while further reductions are planned, such as 70% by fluorescent lamp manufacturers by 2000 (Environment Canada and US EPA 1997). Other Canadian strategies are summarised in Appendix A.

Canadian Tissue Residue Guidelines: Methylmercury

**Table 2. Worldwide production of mercury from ores and concentrates in metric tonnes**

| 1981 | 1982 | 1983 | 1984 | 1985 | 1986             | 1987             | 1988             | 1989             | 1990             | 1991             | Country/continent      |
|------|------|------|------|------|------------------|------------------|------------------|------------------|------------------|------------------|------------------------|
| 76   | 53   | -    | -    | -    | -                | -                | -                | -                | -                | -                | Germany, FR            |
| 252  | 159  | -    | -    | -    | -                | -                | -                | -                | -                | -                | Italy                  |
| 1560 | 1540 | 1416 | 1520 | 1539 | 1471             | 1533             | 1499             | 967              | 962              | 52               | Spain                  |
| 67   | 71   | 65   | 80   | 130  | 147              | 144              | 135              | 160              | 140              | 74               | Finland                |
| -    | -    | 52   | 72   | 88   | 75               | 67               | 70               | 51               | 37               | 35 <sup>a</sup>  | Yugoslavia             |
| 1955 | 1823 | 1533 | 1672 | 1757 | 1693             | 1764             | 1704             | 1178             | 1139             | 161              | Europe <sup>a</sup>    |
| 204  | 246  | 162  | 182  | 226  | 262              | 211              | 97               | 202              | 47               | 25               | Turkey                 |
| 204  | 246  | 162  | 182  | 226  | 262              | 211              | 97               | 202              | 47               | 25               | Asia <sup>a</sup>      |
| 877  | 386  | 828  | 586  | 801  | 764              | 756              | 662              | 587              | 637              | 431              | Algeria                |
| 877  | 386  | 828  | 586  | 801  | 764              | 756              | 662              | 587              | 637              | 431              | Africa                 |
| 3    | 2    | 4    | 2    | 1    | -                | -                | -                | -                | -                | -                | Dominican Republic     |
| 240  | 295  | 221  | 384  | 394  | 185              | 124              | 345              | 651              | 735              | 720 <sup>b</sup> | Mexico                 |
| 962  | 888  | 864  | 657  | 570  | 510 <sup>b</sup> | 140 <sup>b</sup> | 520 <sup>b</sup> | 470 <sup>b</sup> | 500 <sup>b</sup> | 40 <sup>b</sup>  | United States          |
| 1205 | 1185 | 1089 | 1043 | 965  | 695              | 264              | 865              | 1121             | 1235             | 760              | America                |
| 4241 | 3640 | 3612 | 3483 | 3749 | 3414             | 2995             | 3328             | 3088             | 3058             | 1377             | Western countries      |
| 153  | 151  | 144  | 152  | 158  | 168              | 164              | 168              | 131              | 126              | 75               | Czechoslovakia         |
| 1700 | 1700 | 1700 | 1600 | 1600 | 1500             | 1650             | 1650             | 1500             | 1400             | 1200             | USSR <sup>b</sup>      |
| 800  | 800  | 850  | 800  | 800  | 850              | 900              | 900              | 1200             | 930              | 1000             | China, PR <sup>b</sup> |
| 2653 | 2651 | 2694 | 2552 | 2558 | 2518             | 2714             | 2718             | 2831             | 2456             | 2275             | Eastern countries      |
| 6894 | 6291 | 6306 | 6035 | 6307 | 5932             | 5709             | 6046             | 5919             | 5514             | 3652             | Total World            |

Source: Metallgesellschaft 1992; c.f. OECD 1994

<sup>a</sup> Excluding eastern countries (Czechoslovakia, USSR, and China)

<sup>b</sup> Estimated

**Table 3. Metric tonnes of mercury used per year in various categories in Canada and the United States (OECD 1994).**

| <b>Use Category</b>         | <b>Canada</b>  | <b>U.S.A.</b> |
|-----------------------------|----------------|---------------|
| <b>Year</b>                 | <b>1982/91</b> | <b>1989</b>   |
| Batteries                   | 4.3            | 250.0         |
| Thermometers                | 4.0            | NA            |
| Tooth fillings              | 10.0           | 39.0          |
| Pesticides as seed dressing | 2.0            | NA            |
| Laboratory                  | 4.0            | 18.0          |
| Hospital equipment          | 24.0           | NA            |
| Measurement equipment       | NA             | 87.0          |
| Electrical:                 |                |               |
| Switches                    | NA             | 141.0         |
| Light sources               | 4.5            | 31.0          |
| Paint                       | 2.0            | 192.0         |
| Industry:                   |                |               |
| chlor-alkali                | 1.9            | 381.0         |
| other                       | NA             | 40.0          |
| Other uses                  | NA             | 35.0          |
| <b>Total use</b>            | <b>56.7</b>    | <b>1214</b>   |

NA= not applicable



#### 4.0 SOURCES AND PATHWAYS INTO THE ENVIRONMENT

There are many sources of mercury to the environment, both natural and human related. Natural sources include geological mercury deposits, forest fires and other wood burning activities, volcanoes, and volatilisation from the oceans. The primary human-related sources include: coal and other fossil fuel combustion, chlorine alkali processing, mercury waste in landfills or storage, waste incineration, cement manufacturing and metal smelting. Current estimates suggest that human activities have at least doubled or tripled the amount of mercury in the atmosphere since pre-industrialised times (Krabbenhoft and Rickert 1995). These authors also reported that the atmospheric burden (from all sources) continues to increase by about 1.5% per year. A number of studies have estimated the relative contributions of natural and anthropogenic mercury fluxes to the global atmosphere (Table 4; OECD 1994):

**Table 4. Estimates of Fluxes of Mercury to the Global Atmosphere**

| Process                 | Metric tonnes of mercury per year |  |                 |                       |
|-------------------------|-----------------------------------|--|-----------------|-----------------------|
|                         | Anthropogenic emissions           | 2000-10,000                                      | 3560 (910-6200) | 2000                  |
| Natural emissions       | <15000                            | 2500 (100-4900)                                  | 3000-4000       | 3000 (2000-9000)      |
| Total present emissions | 2000-<25,000                      | 6060 (1010-11,100)                               | 5000-6000       | 7500 (5000-15,000)    |
| Reference               | Lindqvist et al. 1984             | Nriagu and Pacyna 1988, Nriagu 1989 <sup>a</sup> | Fitzgerald 1986 | Lindqvist et al. 1991 |

Notes:

<sup>a</sup> Specific sources are listed in Table 5.

#### 4.1 Natural Sources and Pathways

Various amounts of the mercury are found naturally in rocks and surface materials in all parts of the world. Weathering, leaching and runoff of mercury from soils directly contribute to the mercury in receiving waters. The processes are augmented by deforestation and other forms of land disturbance that promote the release of mercury (Veiga et al. 1994).

The most abundant mercury-containing ores are cinnabar (red HgS), metacinnabar (black HgS) and livingstonite (HgSb<sub>4</sub>S<sub>7</sub>) (D'Itri 1972). The mercury content of the vermilion-red sulphide mineral, cinnabar exceeds 86%. It occurs in high quantities in relatively few locations in the world (Mittra 1986), although not all countries that have large deposits of mercury are involved in its mining. Rocks and soils commonly have concentrations of between 5 and 1000 µg·kg<sup>-1</sup> with highest levels near cities and some of the lowest levels in cultivated chernozemic soils of the Canadian prairie provinces (Anderson 1979).

Major mercuric sulphide deposits in Canada are located mainly in western Canada (with one occurrence at Clyde Forks, Ontario). In British Columbia significant economic grade deposits of mercury are known (Nagpal 1989), with concentrations of more than 1 g·kg<sup>-1</sup> in the Kamloops, Port Clements, Bridge River and Yalakom River areas (AEP 1992). No mercury mines are currently operating in Canada (A. Plouffe 1999, GSC, pers. com.).

The volatile nature of mercury makes degassing of the earth's mantle and volcanic emissions important natural sources of mercury to the atmosphere (U.S. NAS 1977). Oceanic submarine

volcanoes (Varekamp and Buseck 1986), and seismic activities like earthquakes (Varshal et al. 1985) are other natural sources of mercury. Biological formation of volatile elemental mercury from methylmercury is also recognised as a source of Hg to the atmospheric pool (Lindqvist et al. 1991).

While the vast majority of natural mercury emissions is from the natural weathering of rocks and mineral deposits and degassing, mercury is also released to the atmosphere through forest fires and other wood burning activities. For example, annual releases to the atmosphere from natural fires in the boreal forest region of Manitoba was calculated to be  $20 \text{ g Hg}\cdot\text{ha}^{-1}$  representing 0.02% of the provincial annual emissions from natural sources which creates high short-term emissions in the form of a pulse. The evaluation assumed 0.4 ppm as the Hg concentration in timber, but about 0.08 ppm was considered transferred to the atmosphere during fires (Williamson 1986). Worldwide, forest fires are estimated to be responsible for releasing 20 t of Hg to the atmosphere each year, which is less than 1% of one researcher's estimate of natural emissions (Nriagu 1989). Intentional wood combustion was estimated to represent 60 to 300 t of Hg (about 5% of all anthropogenic emissions in 1983).

Different types of lakes are associated with proportionally different species and sources of mercury and methylmercury (MeHg). In seepage lakes (no surface inflow), essentially all MeHg is formed within the lake (Watras et al. 1994). In drainage lakes, significant inorganic mercury or organic mercury may be transported from the drainage basin to the lake complexed with dissolved organic carbon (DOC) or particulate matter (Lee and Hultberg 1990; Lee and Iverfeldt 1991), thereby supplementing direct atmospheric sources and in-lake net methylation. The processes of MeHg production and transport to aquatic systems will be discussed more fully in chapter 5.

Watersheds tend to be net sinks of mercury, with the amount of mercury retained often depending on the proportion of wetlands in the watershed (Mierle 1995). More mercury and methylmercury are exported downstream from watersheds dominated by wetlands because of complexation with dissolved organic matter (Mierle and Ingram 1991; Engstrom et al. 1994; Rudd 1995). Also, more mercury usually enters lakes that have a large drainage basin relative to the lake area. The same relationship to high  $A_{\text{dr}}/A_{\text{lake}}$  has been associated with increasing mercury levels in some Ontario fish (Suns and Hitchin 1990).

## 4.2 Anthropogenic Sources

Of the significant sources of mercury to the Canadian environment; smelting, municipal solid waste and sewage sludge incineration, fossil-fuel burning, and cement manufacturing are presently considered the most important.

From all Canadian sources, mercury emissions were estimated at 40 t during 1978, 31 t in 1982 (Jaques 1987), 33 t (Environment Canada 1997; L. Trip 1999, Environment Canada pers. com.), or 43 t (EPRI 1996) in 1990, and declined to 12 t in 1995 (Environment Canada 1997; L. Trip). The decline by 1995 was principally due to the closure of a smelter in Flin Flon, Manitoba that resulted in reducing overall smelter emissions to about 26% (<5 t) of total mercury emissions in Canada. It also reduced the estimated overall national emissions by more than 50%. Emissions of mercury to the atmosphere as of 1995 totalled approximately 143 t for the U.S. (Pilgrim and EMAN 1998), down from 280 t in 1990. In 1990, about two-thirds of Canadian emissions (22 t) were from non-ferrous metal smelting. The next highest emission sector in 1990 was the incineration of municipal solid waste and sewage sludge, at 9.4% (of 33 t), or 3.1 t (L. Trip, unpubl. data). Anthropogenic mercury emissions in Canada for all sectors in 1990

compared to 1995 are shown in Figure 1, and emissions for each province in 1990 are shown in Figure 2<sup>4</sup>.

Emissions reported to the National Pollutant Release Inventory (NPRI) in 1995 totalled 2.4 t of mercury (NPRI 1997). The highest reporting emitter was Cominco Ltd. in Trail, BC, which released 1.8 t to the air, and 60 kg to the Columbia River. All other industries in Canada combined, reported a total of 476 kg to air and 11 kg to water (NPRI 1997). Reported emissions are only a fraction of the total releases of mercury to the environment that actually occur. The discrepancy is mainly due to other sources including fossil fuel combustion that are not covered under the NPRI<sup>5</sup>, but which are included in the Canada-wide estimates discussed above (L. Trip 1999, Environment Canada, pers. com.).

On a global scale, the major air emissions of mercury originate from coal combustion and incineration of solid waste products, whereas the major emission to water is from manufacturing processes. Nriagu and Pacyna (1988) and Nriagu (1989) estimated the worldwide emissions of mercury to the atmosphere, soil and water for 1983, listing all anthropogenic plus total natural sources (Table 5). Even though the anthropogenic component has decreased as use has declined, some researchers agree that anthropogenic sources still comprise about half of the global emissions (see Flux estimate table above; Lindqvist et al. 1984, 1991; Fitzgerald 1986).

Of atmospheric mercury originating from anthropogenic sources, worldwide fossil fuel combustion is estimated to provide one-half, while the rest is released through volatilisation of fungicides, paints, manufacturing, mining, waste incineration, battery decay, fluorescent lamps and various other sources (Douglas 1991). Fewer chlor-alkali operations exist than in former times, but coal-fired power plants and petroleum product combustion (gasoline, oil, etc.) that release mercury in the vapour phase directly to the atmosphere, continue to be important sources. An estimate reported annual global emissions at 2214 t as of 1990. That estimate reported that Asian countries combined emit the highest tonnage of mercury to the atmosphere, at 44% of global emissions (976 T; L. Trip 1999, Environment Canada pers. com.). As of 1990, North America, western Europe, and eastern Europe plus the former USSR countries each emitted about 15%, or 330-349 t of mercury (Figure 3). Of the 1990 total, North American emissions of 330 t, the United States emitted 84.9%, Canada emitted 9.4%, and Mexico emitted 5.7% (Figure 3).

Mercury in solid waste is a potentially preventable source of environmental mercury contamination. For example, the overall quantity of mercury in the U.S. in municipal solid wastes was predicted by US EPA (1992) to decrease through the 1990's, falling from a total of 709 tons in 1989 to 173 tons in the year 2000 (Table 6)<sup>6</sup>. The amount of mercury contributed by batteries was also predicted to decrease, from approximately 621 tons in 1989 to 98 tons in the year 2000, largely due to reduced use of mercury in these products. In contrast, US EPA projected that mercury in solid waste from electrical lighting would slowly increase, rising from approximately 27 tons in 1989 to 41 tons in the year 2000 (Smith and Rowan-West 1996).

Mercury in solid waste in Canada has not been quantified (G. Ironside, Environment Canada, pers. com.). Assuming that Canadian usage of mercury is similar to the U.S, the values cited for the U.S. in the previous paragraph may be reduced by a factor of about ten to estimate

---

<sup>4</sup> 1995 provincial emission data were unavailable as of the time of writing (L. Trip).

<sup>5</sup> Appendix containing detailed a) national compilation of discharge and emission data; and b) control/treatment technology data and information is in preparation.

<sup>6</sup> Mercury mining operations in the U.S. ceased in 1991. In 1994, the U.S. suspended sales of its stockpile mercury.

amounts of mercury in solid waste in Canada.

Even though many point sources of mercury have been removed, long-range transport of mercury in the atmosphere remains elevated in North America in comparison with pre-industrial times (Engstrom et al. 1994). One global estimate suggested that about one-third of atmospheric mercury came from the oceans, and one-third resulted from natural terrestrial processes like degassing and geologic weathering (Fitzgerald 1986). The remaining atmospheric sources were estimated to be from the production, use and disposal of mercury by humans.

Mercury releases and usage are declining in Canada and the United States, but because of the persistence of mercury compounds in the environment, former discharges contribute to the current atmospheric loading and redistribution of mercury in the environment. Mercury sources, uses and recycling in the environment are reviewed in Nriagu (1979, 1989), IJC (1990), AEP (1992), Watras et al. (1994), and Rudd (1995).

At a recent meeting of the Scientific Committee on Problems of the Environment (SCOPE, Stockholm 1993), it was concluded that anthropogenic sources account for approximately 50% of atmospheric mercury (Jernelöv and Ramel 1994). The Expert Panel on Mercury Atmospheric Processes (EPMAP), which convened in March 1994, suggested that as much as 75% of total emissions were from anthropogenic sources (EPMAP 1994). EPMAP also reported that historical deposition in lake and peat sediments indicate that global atmospheric concentrations of mercury have increased by a factor of about three (rather than a factor of two as discussed above) since pre-industrial times. All experts agree that precise estimates are difficult to make because it is currently impossible to accurately assess the recycled portion of mercury compounds emitted from the earth's surface including the oceans.

**Table 5. Estimates of worldwide emissions of mercury (metric tonnes per year) to the atmosphere, soil, and water in 1983 (from Nriagu and Pacyna 1988 and Nriagu 1989).**

| Source Category                         | Atmosphere <sup>a</sup> |              | Water       |             | Soil <sup>b</sup> |              |
|---|-------------------------|--------------|-------------|-------------|-------------------|--------------|
|   | min.                    | max.         | min.        | max.        | min.              | max.         |
| Agricultural waste                      | no estimate             |              | no estimate |             | 0                 | 1700         |
| Atmospheric fall-out                    | no relevance            |              | 220         | 1800        | 630               | 4300         |
| Coal combustion                         | 650                     | 3500         | 0           | 3600        | 370               | 4800         |
| Dumping of sewage sludge                | no relevance            |              | 10          | 310         | no relevance      |              |
| Logging and other wood wastes           | no estimate             |              | no estimate |             | 0                 | 2200         |
| Manufacturing Processes                 | no estimate             |              | 20          | 2300        | no estimate       |              |
| Metal mining                            | Insignificant input     |              | 0           | 150         | no estimate       |              |
| Mine tailings                           | no estimate             |              | no estimate |             | 550               | 2800         |
| Non-ferrous metal production            | 45                      | 220          | 0           | 40          | 0                 | 80           |
| Phosphate fertilizer production and use | Insignificant input     |              | no estimate |             | no estimate       |              |
| Refuse incineration                     |                         |              | no estimate |             | no estimate       |              |
| Municipal                               | 140                     | 2100         |             |             |                   |              |
| Sewage sludge                           | 15                      | 60           |             |             |                   |              |
| Smelter slags and wastes                | no estimate             |              | no estimate |             | 50                | 280          |
| Urban refuse                            | no estimate             |              | no estimate |             | 0                 | 260          |
| Wastage commercial products             | no estimate             |              | no estimate |             | 550               | 820          |
| Wastewater                              | no relevance            |              | 0           | 600         | 10                | 800          |
| Wood combustion                         | 60                      | 300          | no estimate |             | no estimate       |              |
| Total inputs (Median)                   | 910                     | 6180 (3560)  | 250         | 8800 (4600) | 2140              | 18040 (8300) |
| Natural (Median)                        | 100                     | 4900<br>2500 | no estimate |             | no estimate       |              |

Notes:

<sup>a</sup> Insignificant contributions to the atmosphere from: oil combustion, zinc-cadmium production, secondary non-ferrous production, steel and iron manufacturing, cement production, and mobile sources.

<sup>b</sup> Landfills included

**Table 6. U.S. nation-wide discards<sup>a</sup> (tonnes per year) of mercury in products in municipal solid wastes (MSW), 1970 to 2000**

| <b>Products</b>        | <b>1970</b>  | <b>1980</b>  | <b>1989</b>  | <b>1995</b>  | <b>2000</b>  |
|------------------------|--------------|--------------|--------------|--------------|--------------|
| Dental Uses            | 9.3          | 7.1          | 4.0          | 2.9          | 2.3          |
| Electric lighting      | 19.1         | 24.3         | 26.7         | 33.6         | 40.9         |
| Fever Thermometers     | 12.2         | 25.7         | 16.3         | 16.9         | 16.8         |
| Film Pack Batteries    | 2.1          | 2.6          | 0.0          | 0.0          | 0.0          |
| Household batteries    | 310.8        | 429.5        | 621.2        | 176.6        | 98.5         |
| Mercury Light Switches | 0.4          | 0.4          | 0.4          | 1.9          | 1.9          |
| Paint residues         | 30.2         | 26.7         | 18.2         | 2.3          | 0.5          |
| Pigments               | 32.3         | 23.0         | 10.0         | 3.0          | 1.5          |
| Special Paper Coating  | 0.1          | 1.2          | 1.0          | 0.0          | 0.0          |
| Thermostats            | 5.3          | 7.0          | 11.2         | 8.1          | 10.3         |
| <b>Total discards</b>  | <b>421.8</b> | <b>547.5</b> | <b>709.0</b> | <b>245.3</b> | <b>172.7</b> |

Sources: US EPA 1992; Smith and Rowan-West 1996.

<sup>a</sup> Discards before recovery (some MSW may be recovered for recycling).

## 5.0 ENVIRONMENTAL FATE AND BEHAVIOUR

While chapter 4 described the sources of mercury and the “big picture” of its transport in the environment, this chapter will review the smaller scale processes involved in the behaviour of the various mercury species. In ecosystems, the form of mercury and its mobility is dependent on environmental characteristics. Site-specific factors that control the net production (balance of methylation and demethylation), photodegradation, and distribution of MeHg between gaseous, dissolved and solid phases are of particular importance because MeHg is much more toxic and more readily bioaccumulated than any other form of mercury.

The mercury cycle describes the fate, behaviour, and transport of mercury in the environment. For the purposes of this report, fate is defined as the quantification and movement of various mercury species among environmental phases (i.e., gaseous, water, colloids, suspended particulates, sediment, or biota) in lakes and watersheds. Sorption, desorption, diffusion, volatilisation and transformation (reduction, methylation, demethylation) are among the processes that the various mercury species undergo.

For inorganic mercury, the most toxicologically relevant process is methylation. For MeHg, processes that define its fate as a species include demethylation and photodegradation. Diffusion of MeHg through biological surfaces and bioaccumulation are components of the mercury cycle that will be discussed more fully in the following chapter.

### 5.1 The Mercury Cycle

Chemical cycling is defined as inputs to and outputs from a system, along with transport and transformation of chemical species within the system (Zillioux et al. 1993). Transport of mercury, and especially MeHg, among various abiotic and biotic compartments is a concern when it threatens the health of aquatic biota or humans.

The mercury cycle in aquatic ecosystems is complicated by numerous possible mercury species that are present, and by variables affecting speciation. In the last decade, there was a marked increase in our knowledge of environmental concentrations of mercury species because of the development of strict clean protocols for the sampling and improvements to the analysis of mercury and MeHg (Gill and Fitzgerald 1985;1987; Driscoll et al. 1994). Typical total mercury concentrations in ‘uncontaminated’ lake water (epilimnetic) in the northern U.S. are about 0.5-4 ng·L<sup>-1</sup> (Hurley et al. 1994; Watras et al. 1994), with MeHg making up a small proportion of the total.

Generally, the aquatic mercury cycle may be characterised by high atmospheric fallout, removal by deposition to sediments, re-emission to the atmosphere as elemental Hg (Hg<sup>0</sup>), methylation and demethylation transformation processes within the lake or watershed (Figure 4; Watras et al. 1994 1995) and exchanges through tributaries and groundwater (runoff and seepage; Hurley et al. 1994). Some portion of the total mercury pool is accumulated by biota, with biomagnification of MeHg in higher trophic levels of food webs.

Atmospheric fallout directly provides most of the mercury to watersheds that are uninfluenced by point sources or high natural source contamination (Figure 4). Wet and dry fallout of mercury to lakes occurs largely as inorganic mercury (mainly Hg<sup>2+</sup>; Winfrey and Rudd 1990), with insignificant inputs of MeHg (Bloom and Watras 1989). Long-range atmospheric transport of mercury results in mercury deposition to environments thousands of kilometres from emission sources, although about half of the fallout originates within 1000 km (EPMAP 1994).

In surface waters, mercuric ions ( $\text{Hg}^{2+}$ ) may be reduced microbially to form elemental mercury (Ramamoorthy et al. 1983). Photochemical reduction may also occur with exposure to  $\text{UV}_B$  in sunlight (Amyot et al. 1994). Most surface waters are oversaturated with  $\text{Hg}^0$  and because it is volatile, it is likely to be re-emitted to the atmosphere (Vandal et al. 1991; Driscoll et al. 1994). Small amounts of MeHg may also volatilise (especially dimethylmercury), but  $\text{Hg}^0$  is by far the largest component of the total gaseous mercury concentration in air (Schroeder et al. 1991). Re-emissions are important sources of mercury to the global cycle and can result in oxidation to  $\text{Hg}^{2+}$  and re-entry to surface waters in association with precipitation or dry fallout (Figure 4).

Oceanic production and subsequent emission of  $\text{Hg}^0$  is a particularly important facet of the global Hg cycle. This is particularly true near ocean margins (Cossa et al. 1996). The oceans receive about 90% of their mercury as  $\text{Hg}^{2+}$  from wet and dry atmospheric deposition. Elemental mercury forms about 5 to 30% of the total Hg in ocean waters (Mason et al. 1994). Biological production of  $\text{Hg}^0$  has been reported in the open ocean, coastal mesocosms and in permanently anoxic fjords in Norway (Vandal et al. 1991). While re-emissions of the volatile  $\text{Hg}^0$  result in a loss of reactive Hg species ( $\text{Hg}^{2+}$ ) from a location where they might become methylated, the continual recycling of Hg back to the atmosphere prolongs the impact of anthropogenic Hg on aquatic systems. This is because while most of the Hg deposited to the ocean is recycled to the atmosphere, the terrestrial environment and freshwater ecosystems become the principal atmospheric Hg sinks (Mason et al. 1994).

Once in lakes, most mercury binds to particulates and is quickly deposited to sediments, where it may be buried or transformed. Of all of the binding sites, particulates are considered to be most significant and may result in the efficient removal of mercury from the water column by sedimentation. Methylmercury, and especially  $\text{HgCl}_2$ , bind quite strongly with suspended particulates (Table 1). After deposition, the mercury is found complexed with the sulphide as  $\text{HgS}$ , or adsorbed on organics, clays, and Fe or Mn oxides (Miller 1975; Jackson 1989).

The deeper water near the sediment-water interface in lakes is often enriched in mercury compared with other zones in the lake. Because of the finding that particulate-associated mercury concentrations were higher in the deeper layers than the shallow layers of stratified lakes, it was concluded that allochthonous (source outside the lake) Hg was scavenged by particles in the upper waters and transported down with settling particulates (Watras et al. 1994). The alternative was that deeper waters were enriched in particle-associated mercury because of resuspension from bottom sediments. Resuspension can occur due to turbulence from water currents, or bioturbation. In sediments, tubificids, clams, chironomids, other invertebrates, and even fish have been known to reintroduce mercury to the water column by disturbing the sediments (Andersson et al. 1990).

Porewater diffusion (movement through sediments) of mercury is undocumented for many types of lakes. Measurements of sediment porewater in a seepage lake indicated that diffusional movement of total mercury averaged  $1.4 \cdot 10^{-8} \text{ ng} \cdot \text{cm}^{-2} \cdot \text{s}^{-1}$  (concentration gradient of  $15\text{-}35 \text{ ng} \cdot \text{L}^{-1}$ ; Hurley et al. 1994). The authors considered that the diffusional flux from sediments to the water column was minor compared to redissolution of recently fallen particulate material at the sediment surface. Compared with other metals like iron, mercury release from bottom sediments is not nearly as highly redox-dependent (Hurley et al. 1991).

Runoff and seepage of mercury associated with particulates or dissolved organic matter from watersheds may also introduce both inorganic and methylated forms of mercury to lakes and rivers. Substantial increases in total mercury and MeHg may occur when groundwater discharges through wetlands into streams or from rivulets that drain across the surface of



wetlands (Krabbenhoft et al. 1995). Once in surface waters, mercury may enter sediments as described above, be transformed (e.g., methylated or demethylated) and/or be transferred to biota. More so than inorganic mercury, MeHg has some affinity for organic matter and biological tissue<sup>7</sup> because of its attraction to sulphhydryl groups in proteins, thus may become preferentially concentrated in fish and other aquatic biota.

Numerous studies have indicated the importance of organic carbon in binding and transporting mercury compounds (Ramamoorthy and Kushner 1975; Lee and Iverfeldt 1991; Mierle and Ingram 1991; Miskimmin 1991; St. Louis et al. 1994). There is little question that dissolved organic carbon (DOC - often measured as "colour") can increase the apparent solubility of mercury. For example, the export of mercury from watersheds to lakes is closely tied to the export of humic material (DOC; Mierle and Ingram 1991). Hurley et al. (1995) found a positive correlation between the percent wetland in the watershed and MeHg yield to rivers. In northern Ontario, even though upland watersheds without wetlands yielded more total mercury per area to lakes, high DOC wetland portions of watersheds yielded 26-79 times more MeHg than uplands (St. Louis et al. 1994). Since the proportion of MeHg that is transported from watersheds is dependent on biotic and abiotic MeHg production, abundant mercury methylation probably occurs in wetland environments, where high organic carbon and bacterial activity are typical.

Seasonally, mercury in the water column is lower in winter than in summer because frozen lakes are sealed off from atmospheric sources of mercury and most particulate settling has occurred (Watras et al. 1994). There is also a seasonal component to methylmercury cycling. Methylmercury concentrations in water were found to be higher in the late summer than at other times of the year in both rivers (Hurley et al. 1995) and lakes (Watras et al. 1995). Ramlal et al. (1993) observed the same seasonal trend for measured net methylation rates in sediment cores from northern Ontario. Methylmercury concentrations were highest by late in the summer because of the high bacterial activity associated with higher temperatures as the summer season progressed, causing MeHg to accumulate. Methylation but not demethylation was stimulated by warm summer temperatures; while the opposite was true in cold winter conditions (Ramlal et al. 1993). In general, net MeHg production in lakes is increased by factors that enhance bacterial activity, and total Hg is reduced when lakes are sealed off from atmospheric sources. Reviews of the aquatic mercury cycle may be found in Driscoll et al. (1994) and Zillioux et al. (1993). Also, a mathematical model of mercury cycling in lakes was developed by Hudson et al. (1994).

---


<sup>7</sup> MeHg has relatively low log Kow values of  $1.6 \pm 0.2$  (Major et al. 1991) and 2.5 (Halbach 1985).

## 5.2 Mercury Methylation and Demethylation

Before the mid-1960s, inorganic mercury was believed to be relatively inert in the environment. Then it was demonstrated that inorganic mercury could be methylated by micro-organisms to form MeHg, which was biomagnified in food webs (Wood et al. 1968; Jensen and Jernelöv 1969). The production of methylmercury is a balance between methylation and demethylation. Wherever MeHg is detected in the environment, methylation has exceeded demethylation.

Since methylation and demethylation always co-occur, the balance between the two in the aquatic ecosystem is called net MeHg production. Several factors influence the rate of net MeHg production in water. These include the concentration of  $\text{Hg}^{2+}$ , composition of the microbial population, nutrient and mineral substrate, pH, temperature, redox potential, dissolved and particulate organic matter, iron, and sulphate. Methylation by naturally-occurring micro-organisms is probably limited by processes that alter the availability of inorganic mercury (Ramamoorthy et al. 1982; Winfrey and Rudd 1990). Early studies reported that micro-organisms produced MeHg under anaerobic conditions (Jensen and Jernelöv 1969). Subsequently, aerobic methylation was demonstrated (Rissanen et al. 1970), and it is now widely accepted that methylation occurs in both aerobic and anaerobic environments.

MeHg is formed principally through the microbial transfer of methyl groups ( $\text{CH}_3$ ) to inorganic mercury (e.g.,  $\text{Hg}^{2+}$  transformed to  $\text{CH}_3\text{Hg}^+$ ; Robinson and Tuovinen 1984). Methylcobalamin is thought to be the only non-enzymatic methylation agent capable of transferring a methyl group to an inorganic mercuric ion. It can transfer groups as a carbanion ( $\text{CH}_3^-$ ) and a methylradical ( $\text{CH}_3^\bullet$ ) to produce methylmercury and dimethylmercury (D'Itri 1991), as follows:



Under anaerobic conditions, methylcobalamin was shown to be present and the reaction to require ATP and hydrogen as the source of electrons (Wood et al. 1968; Robinson and Tuovinen 1984). Three major co-enzymes may participate in enzymatic methylation, but the process remains unvalidated (D'Itri 1991).

While biotic methylation is the dominant process, some abiotic mercury methylation also occurs with methyl derivatives of lead, tin and silicon (AEP 1992), and in the presence of high concentrations of humic matter ( $6380 \text{ mg}\cdot\text{L}^{-1}$ ; Nagase et al. 1982) along with metal ions (humics  $>170 \text{ mg}\cdot\text{L}^{-1}$ ; Lee et al. 1985).

Demethylation was first discovered by Furukawa et al. (1969), at about the same time that methylation was reported. Demethylation is the degradation of MeHg involving the microbial cleavage of the carbon-mercury linkage followed by the reduction of  $\text{Hg}^{2+}$  to  $\text{Hg}^0$  (Robinson and Tuovinen 1984). In addition to metallic mercury, methane is a by-product of demethylation. The biochemistry and genetics of the plasmids and associated enzymes involved in mercury demethylation were thoroughly reviewed by Summers and Silver (1978); Robinson and Tuovinen (1984); Summers (1986); and Foster (1987).

While non-biological demethylation mechanisms are unknown (Winfrey and Rudd 1990), a newly discovered process is the breakdown of MeHg by photodegradation (Sellers et al. 1996).

Microbial demethylation has long been accepted as the dominant route of methylmercury degradation in fresh waters. It is not known whether photodegradation results in the production of the volatile Hg<sup>0</sup>. Under the conditions of the experiments of Sellers et al. (1996), however, photodegradation of MeHg was about 350 times greater than microbial demethylation. These results suggest that MeHg degradation may be an important process where light penetration is significant (such as in clear and/or shallow water).

A number of environmental variables affect the net production of methylmercury. Most factors that increase rates of bacterial activity will enhance MeHg production. Bacterial activity increases with increasing temperature and available biodegradable organic carbon. Thus, methylation rates tend to be highest in surface sediments with freshly deposited organic matter, and in warm shallow sediments where abundant bacterial activity takes place (Ramlal et al. 1986; Winfrey and Rudd 1990). Methylation also occurs in the aerobic water column (Furutani and Rudd 1980; Xun et al. 1987), in anaerobic water and sediments (Ramlal et al. 1985), fish intestinal contents and surface biofilms (*in vitro* studies; Rudd et al. 1980).

Some lakes are more prone to MeHg problems than others. These include hydroelectric reservoirs (Abernathy and Cumbie 1977; Lodenius et al. 1983; Stokes and Wren 1987; Jackson 1988a; Hecky et al. 1991), low pH lakes (Wren and MacCrimmon 1983; Verta et al. 1986; McMurtry et al. 1989), and even some remote lakes with no apparent other stressors associated with high mercury. The explanation for high MeHg in fish from these lakes is different in each case, but usually relates to factors affecting net MeHg production and transfer to the food web. In the case of newly formed reservoirs, the flooding of fresh vegetation and subsequent high bacterial activity causes mercury methylation to increase substantially (Ramlal et al. 1987; Hecky et al. 1991). The elevated MeHg in fish may decline in a few years or may remain elevated for decades (Bodaly et al. 1984). Prediction of mercury problems associated with reservoirs may be possible using mathematical modelling (Johnston et al. 1991).

In low pH and other remote lakes, the explanation for high MeHg in fish is more complex. Mercury methylation rates are higher in low pH lake water and surface sediments (Xun et al. 1987). In fact, the methylcobalamin-dependent methylation process is optimised at pH 4.7 (D'Itri 1991). It is also possible that the uptake of MeHg by aquatic organisms in acidic waters is increased by calcium mediated changes in gill permeability (Rodgers and Beamish 1983). Further, the importance of sulphate reducing bacteria is enhanced in culturally acidified lakes; these bacteria are thought to be the most important methylators in lake sediments (Gilmour and Riedel 1995). In remote non-acidified lakes with no apparent point source of mercury, sometimes the 'source' is traced to wetlands which may have high methylation rates combined with high DOC-related export of MeHg, as discussed in section 5.1 (The Mercury Cycle).

### **5.3 Factors Affecting Partitioning and Speciation**

The partitioning of mercury between "dissolved" and "particulate" phases is directly related to aqueous speciation and to lake chemistry. Partitioning of a given species is the outcome of the competition for that species among ligands in solution (truly dissolved), small particles called colloids, and organic or inorganic material. Speciation describes whether the form of mercury is an alkylated form, or is one of the many inorganic forms (e.g., HgCl<sub>2</sub>, HgS, etc.). This document will only discuss MeHg and a few of the important inorganic forms. Full reviews and examples of mercury speciation research may be found in Gill and Bruland (1990), Aceto et al. (1995), and in several papers within the book, "Mercury Pollution: Integration and Synthesis" (Watras and Huckabee 1994).

The relative abundance of MeHg in environmental samples is of particular concern because of its high toxicity to biota. Methylmercury is the major form of mercury accumulated in biological tissues due to the ease with which it permeates cell membranes and its strong affinity for sulphhydryl groups in proteins (Faust 1992). It is not highly lipophilic, with moderate  $K_{ow}$  values of up to 2.5 found experimentally (Table 1). In water samples, MeHg is typically a smaller proportion of total mercury than is inorganic mercury. In low alkalinity lakes, MeHg has been reported to represent 10-20% (Adirondacks, Driscoll et al. 1994), and <1-13.3% (lake inflows, Precambrian Shield of Ontario; St. Louis et al. 1994) of total mercury in water samples. An unusual case was for moderately alkaline (up to  $3 \text{ meq}\cdot\text{L}^{-1}$  or  $150 \text{ mg}\cdot\text{L}^{-1} \text{ CaCO}_3$ ), high pH (8-9) lakes in California, where up to 89% of total Hg was MeHg (Gill and Bruland 1990).

The effect of pH on mercury speciation relates partly to its effect on the solubility of inorganic mercury and partly to the finding of higher rates of MeHg production with decreasing pH (Xun et al. 1987; Miskimmin et al. 1992). With decreasing pH, all forms of mercury show increased binding to sediments (where bacteria are abundant). A different relationship between pH and DOC may be important in regulating the availability of inorganic mercury for methylation. Reduction in pH changes the character of DOC by increasing protonation of anions and thus desorbing metals (Davis et al. 1985). If binding of inorganic mercury to some forms of DOC is reduced by low pH, more mercury may be available to micro-organisms for methylation (Miskimmin et al. 1992).

Redox conditions and chloride ions influence Hg speciation and methylation (Gill and Bruland 1990). In well-aerated water (redox  $\geq 0.5\text{V}$ ), mercuric (+2) species will predominate, whereas under reducing conditions, elemental ( $\text{Hg}^0$ ) species will occur (AEP 1992). Abundant MeHg may be produced by anaerobic (e.g., Ramlal et al. 1985), as well as aerobic bacteria. Among the inorganic anions,  $\text{Hg}^{2+}$  forms the strongest covalent bond with chloride ions (stability constant,  $K=10^{15}$ ), so mercuric-chloride species may dominate when chloride salts are abundant (AEP 1992). Chloride salts may be locally important where runoff from road salting operations occur.

Within anoxic zones, inorganic mercury forms strong aqueous complexes with sulphide and precipitates as  $\text{HgS}$  (Craig and Bartlett 1978; Leermakers et al. 1993). These sulphide complexes are generally assumed to control speciation of both  $\text{Hg}^{2+}$  and MeHg. While sulphate reducing bacteria are known to methylate mercury (Gilmour et al. 1992), sulphate reducers make up only a small proportion of the microbial community in most freshwater lakes, where methanogens are the primary anaerobic bacteria. Sulphide is considered important because it out competes humic acid (DOC) by factors of  $10^4$  for  $\text{CH}_3\text{Hg}^{2+}$  and  $10^{18}$  for  $\text{Hg}^{2+}$  (Hudson et al. 1994). Hurley et al. (1994) completed extensive calculations of the equilibrium speciation of mercury and sulphur at a range of redox and pH levels.


Potential sulphide binding that reduces bioavailable mercury may be particularly important in the saline lakes ( $>3 \text{ g}\cdot\text{L}^{-1}$  salt) of Alberta and Saskatchewan which have some of the highest sodium sulphate concentrations in the world ( $\text{Na}_2\text{SO}_4 > 125,000 \text{ }\mu\text{M}$ ; Waiser and Roberts 1995).  $\text{HgS}$  precipitation combined with high pH (associated with low methylation rates), decreased methylation with increasing salinity (Blum and Bartha 1980) and potential  $\text{Hg}^0$  re-emissions (high from shallow, wind-mixed waters) makes significant MeHg bioaccumulation an unlikely problem in prairie saline lakes. Perhaps an analogous situation is in the high pH, high-alkalinity desert lakes in California, where very little or no MeHg was detected using highly sensitive analytical methods ( $0.5 \text{ pg}\cdot\text{L}^{-1}$  detection limit; Gill and Bruland 1990).

## 6.0 BIOACCUMULATION OF MERCURY AND METHYLMERCURY IN AQUATIC AND SEMI-AQUATIC ORGANISMS

Methylmercury has a high potential for bioaccumulation as well as biomagnification with increasing trophic levels. Mercury and MeHg may be taken up by aquatic organisms directly from water or through their diet. Uptake directly from water is the result of dissolved mercury adsorption or absorption through the body surface and respiratory organs such as gills. Uptake via food is based on mercury transfers through the digestive tract. The relative importance of either pathway depends on trophic level of the organisms, duration and intensity of exposure, and environmental factors (Zillioux et al. 1993).

Organisms at lower trophic levels usually contain the lowest proportion of total mercury as MeHg. Aquatic plants contain the lowest MeHg as a percentage of total mercury. Invertebrates often contain about 50% MeHg and 50% inorganic mercury (Hildebrand et al. 1980). Organisms higher in the food chain, like piscivorous fish (e.g., walleye, lake trout), aquatic birds (loons, herons), piscivorous mammals (mink, otters) and marine mammals contain a high proportion of THg as MeHg in muscle tissue. Most piscivorous fish have essentially all MeHg in their muscle tissue (Bloom 1989). The main route of uptake for MeHg is consistent with a number of persistent organic chemicals like DDT, PCBs or dioxins. Once debated, it is now generally accepted that uptake through diet sources is the most important route of uptake for fish (Rodgers 1994) and, for wildlife consumers of fish. Methylmercury accumulates in organic tissue because of its affinity with sulphhydryl groups, the relative ease with which it passes through the digestive wall and slower depuration rate relative to inorganic mercury (Saouter et al. 1993).

### MeHg Biomagnification in the Food Web



### 6.1 Aquatic Plants

Direct adsorption of mercury is the critical pathway for phototrophic organisms like macrophytes and algae. Small algal cells may accumulate more mercury than large ones because of the higher surface area to volume ratio for small cells compared to large cells. Methylmercury is preferentially sequestered in algal cytoplasm while inorganic Hg is retained in the algal membrane (Mason et al. 1995). This has implications for the transfer of MeHg to zooplankton, to be discussed in the next section (6.2).

Because phytoplankton cells continually divide, the partitioning of Hg into live cells is a balance between uptake, depuration and growth dilution. This is fundamentally different from adsorption of Hg onto detrital or inorganic particles. Plankton may potentially accumulate more Hg than does detritus because the living cells have both passive and facilitated transport mechanisms,

as well as intracellular ligands that sequester Hg (Hudson et al. 1994).

The reported data on mercury uptake by aquatic macrophytes are sometimes contradictory, with some studies indicating that mercury accumulation takes place primarily in roots while others indicating accumulation is primarily through rhizomes or shoots. Given that rooted macrophytes accumulate the majority of their nutrients through their roots and the fact that mercury concentrations are higher in sediments than in water, it may seem intuitive that more mercury would be accumulated through the roots from sediments. Jana (1988) found twice as much mercury in roots compared to shoots of water hyacinth (*Eichhornia crassipes*) and Coquery and Welbourn (1995) found seven times more total mercury in the roots than in the shoots of a submergent perennial in Ontario. On the other hand, Thompson (1996) found five times and three times more total mercury in the petioles and leaves, respectively, than in the roots of a floating leaved macrophyte [the yellow pond lily (*Nuphar variegatum*)], further fuelling the debate over "root versus shoot" absorption. Since the majority of macrophyte growth is in the shoot, mercury may also be taken up into new cells directly from the water column. Mortimer (1985) and Crowder (1991) felt that the confusion over "roots versus shoots" may have come from the fact that mercury may be translocated from one portion of the macrophyte to another. Regardless, the important point is that if significant amounts of mercury are accumulated by plants, the mercury will be released to the environment upon senescence, or transferred to animals that consume them.

With MeHg, while high accumulation was measured in leaves, stems and roots of *Elodea densa*, root absorption was reported to be the dominant direct route of uptake, and the leaves were the principal storage organ (Maury et al. 1988).

Environmental variables can affect the bioaccumulation of mercury by plants. Coquery and Welbourn (1995) found that total mercury in macrophyte roots was related to mercury in sediments, but only after correction for sediment organic content. If organic content of the sediments was high, mercury concentrations of the sediments could also be elevated without significant uptake by the plant. In that study, concentrations of mercury in plants were greater than the concentration in sediments, and mercury in roots was consistently higher than in the shoots of *Eriocaulon septangulare* ( $0.22 \mu\text{g}\cdot\text{g}^{-1}$  and  $0.03 \mu\text{g}\cdot\text{g}^{-1}$  dw, respectively). Increasing temperature and photoperiod exert a strong positive influence of MeHg accumulation to macrophytes, with the effects amplified when both are increased concurrently (*Elodea densa*; Maury-Brachet et al. 1990). Differences in accumulation of mercury may also depend on the species of plant, type and amount of complexing ligands and seasonal growth rate changes (Czuba and Mortimer 1980).

Experimentally, the presence of DOC (in the form of humic acid or EDTA) significantly suppressed dissolved mercury uptake by duckweed (Mo et al. 1989). However, in a field survey of 34 Precambrian Shield lakes in Ontario and Québec, algal mercury concentrations showed a significant positive correlation with DOC and with fish mercury concentrations even though mercury in the water was at low to undetectable concentrations (Stokes et al. 1985). The difference between experimental and field results is probably because while DOC may suppress uptake, high DOC lakes often contain more mercury from terrestrial/wetland sources, as discussed section 5.1.

Furthermore, the total mercury concentration in fifty-five *Myriophyllum spicatum* plants has been shown to be negatively correlated with the pH of the water (Thompson-Roberts et al. 1999). In the same study, DOC, alkalinity, and pH were not related to the total mercury concentrations in three other aquatic macrophytes. The authors believed the lack of relationships were due to the

small range in environmental variables.

Bioaccumulation is proportional to dissolved concentrations assuming that the rates of uptake by plants are similar for inorganic and organic forms (Mortimer 1985). Aquatic plants are known to bioconcentrate mercury and MeHg up to several thousand times the water concentration (per g dw). Laboratory studies on four genera of submergent macrophytes from the Ottawa River had bioconcentration factors (BCFs) for total mercury ranging from 900-3300 ( $\mu\text{g}\cdot\text{g}^{-1}/\mu\text{g}\cdot\text{mL}^{-1}$ ; water concentration  $0.28 \mu\text{g}\cdot\text{mL}^{-1}$ , Mortimer 1985). Similarly, in the field, floating macrophytes (*Nuphar variegatum*) from forty-five Ontario wetlands had BCFs for total mercury ranging from 400 to 6000 (Thompson 1996; Thompson-Roberts et al. 1999). Calculations from Jana's (1988) data resulted in BCFs of 670 for an alga, and 800 for water hyacinth.

Mortimer (1985) proposed that concentrations of mercury in plants water could be used to estimate those in water; for example, if an *Elodea* plant contained  $300 \text{ ng}\cdot\text{g}^{-1}$  dw, then the water in which it was growing would contain  $20 \text{ ng}\cdot\text{L}^{-1}$  based on a BCF of 1500. Mortimer (1985) further stated that since approximately 30% of total Hg was MeHg, the water concentration of both MeHg and inorganic mercury would be  $6 \text{ ng}\cdot\text{L}^{-1}$  and  $14 \text{ ng}\cdot\text{L}^{-1}$ , respectively. This theory has not been tested over a wide range of environmental conditions, and is based on uptake from the water column (shoots) rather than from sediments (roots). Furthermore, the experiments preceded the use of trace-metal clean collection and analysis techniques, and should be repeated.

## 6.2 Invertebrates

Mercury is accumulated by aquatic invertebrates directly from water (including porewater), and actively, through ingestion of contaminated food. It may be bound on the outer membranes or chitinous exoskeleton, absorbed within gut contents, incorporated into body tissues and excreted. Mercury bound to outer membranes or exoskeletons may be lost with moulting (Zauke 1977). Also, egg-carrying females have higher concentrations than other individuals because MeHg is readily transferred across lipid bilayers into eggs. As previously stated, MeHg is more likely than inorganic mercury to be absorbed and retained in tissue because of its affinity for sulphhydryl groups and its slow rate of depuration (Saouter et al. 1993). The rate of MeHg uptake from water has been shown to be related to metabolic rate (Visman et al. 1995). Invertebrates tend to have a lower percentage of MeHg than fish because any forms of mercury taken up are either associated with gut contents or the exoskeleton rather than absorbed into tissue. Also, the sources of mercury to which they are exposed are comprised of relatively low percentages of MeHg.

Sediment-associated invertebrates or 'benthos' (e.g., chironomids, amphipods, mussels) generally contain more mercury than those in the water column (e.g., *Daphnia*). Mercury uptake by sediment-dwelling organisms is likely favoured by the combination of high mercury concentrations in their diet as well as in their physical environment (Parkman and Meili 1993). Some suggest that uptake for mussels changes from water to particulates depending on the concentration and composition of suspended particles (Gagnon and Fisher 1997). While laboratory experiments suggested water was the major route of mercury uptake for *Mytilus edulis*, under estuarine conditions, when inorganic Hg on suspended sediments was up to 80 times higher than dissolved Hg, the particulate pathway was dominant (King and Davies 1987). Malley et al. (1996) found that uptake of MeHg by the freshwater mussel, *Pyganodon grandis* was higher after reservoir flooding experiments when suspended particulate concentrations increased substantially.

Because predacious invertebrates belong to a higher trophic level, they will sometimes (but not always) bioaccumulate more MeHg than non-predaceous invertebrates (Parkman and Meili 1993). For example, a predatory mite, *Hydracarina* exhibited the highest bioconcentration of THg and MeHg at  $2.5 \cdot 10^5$  and  $12.6 \cdot 10^6$  times higher than water, respectively (BCF:  $\text{ng} \cdot \text{g}^{-1} \text{ dw}$  per  $\text{ng} \cdot \text{L}^{-1}$ ). In the same lakes, the predacious *Chaoborus* spp. bioconcentrated mercury to lower levels than most non-predacious taxa (Back and Watras 1995). Tremblay (1998) reported the mean proportion of MeHg in 16 Québec lakes increased from detritivores (35-45%; dipterans, ephemeropterans, trichopterans) to predacious species (70-85%; notonectids, coleopterans, odonates).

The proportion of accumulated mercury taken up either from water or food depends on degree of contamination of each compartment, as well as on the chemical forms of mercury present. Saouter et al. (1993) measured the distribution of mercury and MeHg in a common mayfly, *Hexagenia rigida*, after a nine-day exposure in laboratory microcosms containing both water and sediment. Mercury was added either to the overlying water or to the sediment. Methylmercury was taken up 60 times more readily than  $\text{HgCl}_2$  when it was added to sediments, and twice as readily when added to overlying water. Both forms of mercury tended to accumulate in the gut when taken up through ingestion of sediments, whereas they accumulated on the gills when taken up from water (Saouter et al. 1993). Crayfish that were caged with MeHg contaminated or uncontaminated food and water, absorbed virtually all of their MeHg from their diet and not from water (Parks et al. 1988). Invertebrates usually have mercury body burdens of less than 65% MeHg because 1) inorganic mercury concentrations in sediments, particulates and water are proportionally higher than MeHg, and 2) because of the shorter life spans of invertebrates that limits the potential absorption of MeHg into tissue (Parkman and Meili 1993).

Even though MeHg is proportionately less in invertebrates than it is in fish, the trophic transfer of MeHg is much more efficient than for inorganic Hg (62% and 15%, respectively). Mason et al. (1995) demonstrated this principle experimentally between zooplankton and an algal food source by finding that the outer membrane and cytoplasm contained 37% and 63% MeHg, respectively; and 91% and 9% inorganic Hg, respectively. Since zooplankton digest the cytoplasm but defecate the membrane material, MeHg becomes biomagnified at lower levels of the food web because the bulk of the digested material contains a high proportion of MeHg.

Water quality variables can influence the uptake of mercury by aquatic invertebrates. For zooplankton, total mercury content varies positively with mercury concentration in water (Sorensen et al. 1990). For benthic invertebrates (as with aquatic plants), mercury concentrations are not correlated with sediment mercury concentrations (Kristensen 1982; Parkman and Meili 1993). Higher concentrations of MeHg in zooplankton have been reported in acidified lakes (Bloom 1992), especially if they were also brown-water (high DOC) drainage lakes (Westcott and Kalff 1996). Laboratory experiments confirmed that increased uptake of  $\text{HgCl}_2$  and MeHgCl occurred with decreasing pH (Saouter et al. 1993) and that the diffusional fluxes of mercury over lipid bilayers were higher at pH 5.0 than at pH 8.5 (Boudou et al. 1991). There is some evidence that DOC suppresses uptake of both inorganic Hg and MeHg by invertebrates based on results that show a negative correlation of all forms of Hg in zooplankton with DOC in seepage lakes (no terrestrial sources of Hg; Watras et al. 1994; Back and Watras 1995).

Higher mercury concentrations are found in invertebrates that reside in the deepwater compared to shallow sediments, particularly in anoxic conditions (Särkkä 1979). Mercury concentrations may be ten times higher in deepwater than near shore taxa (Parkman and Meili 1993). The sediment-water interface is the zone of highest Hg concentrations, steep oxygen



gradients, and high bacterial activity (thus methylation potential), therefore minor differences in feeding strategy may result in large differences in Hg bioaccumulation.

Invertebrates that feed on plants and sediments may contain more THg than do predatory invertebrates (Särkkä 1979; Parkman and Meili 1993). This is attributed to the high proportion of inorganic mercury found in non-predatory animals; predators would not accumulate inorganic mercury because it is poorly retained compared with MeHg (Parkman and Meili 1993). Theoretically, the larger body size of predators may reduce the mercury concentration in their tissue by growth dilution (important for fish). More research is required in this area.

As animals take up mercury through both diet and water, it is difficult to separate a bioconcentration factor, which is the concentration of a contaminant in the animal relative to the water concentration ( $BCF = \frac{[animal]}{[water]}$ ), from a bioaccumulation factor (BAF), which theoretically includes uptake from both sources (i.e.,  $\frac{[animal]}{[water + food]}$ ; Metcalfe 1986). Whenever values are calculated from field results, a BCF is often reported, although the two uptake routes cannot truly be separated. When diet is reported to be the predominant route of uptake, comparing the contaminant concentration in the consumer to the concentration in their food is more representative. For example, crayfish had BAFs ( $\frac{[crayfish]}{[diet]}$ ) of 4.5 to 8.3 when fed a high and low MeHg diet of fish tissue over a period of ten weeks (Parks et al. 1988). Some older examples of wet weight-based total mercury BCFs are 900 for a dragonfly nymph from a contaminated lake in Ontario (Clay Lake, Smith et al. 1975) and 29 000 for a dragonfly nymph from a stream in South Carolina (Cox et al. 1975)<sup>8</sup>.

Because of the scarcity of MeHg measurements done on water, plants and invertebrates within individual studies, few bioconcentration factors and/or bioaccumulation factors for MeHg are available for invertebrates. Tremblay (1998) found a MeHg biomagnification of about three between the two adjacent trophic levels of phytoplankton and zooplankton. BCFs for zooplankton were reported to increase by 10 to 100 times more for MeHg compared to inorganic Hg (based on wet weights; Watras and Bloom 1992). Concentrations of MeHg and THg were higher in all compartments in the experimentally acidified basin (pH 4.7) of Little Rock Lake. The BCFs ( $\frac{[zooplankton]}{[water]}$ ) were an order of magnitude higher for MeHg than for inorganic Hg. BAFs ( $\frac{[zooplankton]}{[phytoplankton]}$ ) were the same for inorganic Hg in both the acidified and reference basins (BAF=1.9); BAFs for MeHg were 3.6 and 5.0 in the acidified and reference basins, respectively (Watras and Bloom 1992). These data supported the hypotheses that the uptake of MeHg by zooplankton was more efficient than inorganic Hg, and that uptake of MeHg is proportional to supply regardless of lake acidification.

### 6.3 Freshwater Fish

Essentially all mercury found in piscivorous freshwater fish tissue is MeHg (approximately 99%). Early studies that reported less than 95% MeHg may have been limited by their analytical methodology or problems with homogeneity of subsamples (Bloom 1992). While inorganic mercury is always more abundant in natural ecosystems, it is absorbed by fish much less efficiently than MeHg, and if taken up, is eliminated more rapidly than is MeHg (Huckabee et al. 1979; Boudou and Ribeyre 1985; Trudel and Rasmussen 1997). This is because the digestive wall is much more permeable for MeHg than for inorganic mercury (Boudou and Ribeyre 1985), allowing MeHg to be readily transferred to other tissues.

---

<sup>8</sup> Note that these BCFs are based on measurements pre-dating modern collection and analytical techniques.

Biomagnification of MeHg through diet, rather than gill uptake from water, is considered the dominant basis for elevated concentrations in fish (perhaps 90%; Harris and Snodgrass 1993; Rodgers 1994). Recent measurements of mercury speciation in water indicate fish BCFs for MeHg of  $10^6$  to  $10^7$ , and for non-MeHg BCFs are less than  $10^4$  (Bloom 1992). This demonstrates that MeHg biomagnifies in fish by 100 to 1000 times more than other forms of mercury (compared to 10 to 100 times for zooplankton).

Harris and Snodgrass (1993) modelled the uptake of MeHg from water in yellow perch (*Perca flavescens*) and walleye (*Stizostedion vitreum*), as well as uptake through two types of diet. Methylmercury concentrations were modelled with realistic concentrations for uncontaminated water ( $0.05 \text{ ng}\cdot\text{L}^{-1}$ ), for zooplankton ( $0.033 \text{ ng}\cdot\text{g}^{-1} \text{ ww}$ ) and for forage fish ( $0.30 \text{ ng}\cdot\text{g}^{-1} \text{ ww}$ ). Ninety-nine percent of MeHg was accumulated from the diet over several years, taking about one year to reach equilibrium when the diet changed from zooplankton to forage fish. Even when the water concentrations were increased to 'unrealistically high' levels ( $0.3 \text{ ng}\cdot\text{L}^{-1} \text{ MeHg}$ ), and allowing for uncertainty factors, uptake from water was always less than 20% of the total.

Their simulations indicated that excretion was more effective in young fish than from older fish, suggesting that remedial actions may be more effective in reducing mercury in younger fish (may take 10 years to excrete 50% for a 10-year-old walleye; Harris and Snodgrass 1993). The calculations were based on an assumed allometric exponent of -0.6 for Hg elimination. Trudel and Rasmussen (1997) recently calculated the exponent to be closer to -0.2, which would mean that smaller fish excrete MeHg more slowly than the Harris and Snodgrass model predicts. In general, MeHg is excreted very slowly relative to uptake for most fish. Methylmercury is not eliminated rapidly enough to prevent a net accumulation as long as exposure is continued.

Even in the absence of trophic transfer of MeHg, large adult predatory fish would be expected to have higher concentrations of MeHg than their prey because of their longer exposure as a result of age (predatory fish have longer life spans than forage fish) (Huckabee et al. 1979). Fish length, the preferred surrogate for age, is the best single predictor of Hg concentrations in fish (Richardson 1994; G.M. Richardson 1999, O'Connor Associates, pers. com.). Other factors such as differences in food consumption rates, food conversion efficiencies, and fish metabolic rates are also important (Canada-Manitoba Mercury Agreement 1987). The assimilation efficiency of MeHg from food to fish is often estimated at 80% (Norstrom et al. 1976; Harris and Snodgrass 1993; Trudel and Rasmussen 1997).

Recent information indicates that generalisations about the trophic level that a particular species occupies may lead to errors in predictions about potential MeHg concentration. For example, fish may be piscivorous in one lake, but insectivorous or omnivorous in another if no smaller forage fish occupy the lake. This difference in diet may cause the wholly piscivorous fish to have high mercury levels, while the same species in the lake with the shorter food chain may not. Cabana et al. (1994) clearly established that the food chain structure in 81 Ontario lakes had this effect for mercury accumulation by lake trout. Mercury levels in trout from lakes with the longest food chains where pelagic forage fish and the crustacean *Mysis relicta* were present were about 3.6-fold higher than those from lakes with the shortest food chains, where the two trophic components were missing. The importance of food chain length has been demonstrated for other persistent contaminants such as PCBs (Ontario lakes; Rasmussen et al. 1990) and toxaphene (Arctic lakes; Kidd et al. 1993). The relatively inexpensive procedure of measuring stable nitrogen isotopes may be used for screening the trophic level of fish from lakes prior to undertaking measurements of contaminants that biomagnify (Cabana and Rasmussen 1994).

Reservoirs may have fish with elevated mercury concentrations whether or not they have high

pH or alkalinity<sup>9</sup>. In reservoirs, the flooded organic matter and associated increased bacterial activity enhances MeHg production (Canada-Manitoba Mercury Agreement 1987; Ramlal et al. 1987). Any bioavailable MeHg may then be bioaccumulated. Overviews of the mercury problem in specific reservoirs are found in Bodaly and Hecky (1979), Verta et al. (1986), Jackson (1987), Canada-Manitoba Mercury Agreement (1987), Verdon et al. (1991), Anderson et al. (1995), Rodgers et al. (1995), Verdon and Tremblay (1998), and a model of fish mercury levels as related to physical characteristics of Canadian reservoirs was developed by Johnston et al. (1991). While these reports concern reservoirs in Manitoba, Saskatchewan, Ontario, Québec, Labrador and Scandinavia, reservoirs in Alberta have not been associated with elevated mercury in fish to date (AEC 1989-Gleniffer L, Dickson Dam.; AEC 1993-Oldman River Dam). Screening for mercury in fish from reservoirs in British Columbia is in progress (Watson 1992).

Numerous chemical and physical variables of surface waters determine the potential for fish bioaccumulation of mercury (Table 7). In particular, low pH (<6), low alkalinity (acid-neutralising capacity  $50 \mu\text{eq}\cdot\text{L}^{-1}$  or less), and low calcium (< $5 \text{mg}\cdot\text{L}^{-1}$ ) lakes are associated with elevated mercury concentrations in fish (Grieb et al. 1990; Spry and Wiener 1991). Studies of the effect of pH on uptake by fish show conflicting results. In controlled laboratory studies, Rodgers et al. (1987) found no difference in accumulation of MeHg by walleye and trout among three pH treatments (5, 6, and 7); whereas Drummond et al. (1974) found greater MeHg accumulation by brook trout at pH 6 than 9. In drainage lakes (i.e., those that receive terrestrial runoff), DOC more strongly (positively) influences MeHg concentrations in fish than either pH or alkalinity, possibly through DOC-mediated transport of MeHg from the watershed<sup>10</sup> (Richardson 1994). DOC input into lakes is, in turn, negatively correlated to sulphate deposition in the drainage basin because the solubility of organic carbon declines as pH lowers. Thus, in drainage lakes, acidification may indirectly reduce MeHg levels in fish by limiting DOC-mediated transport of MeHg from the drainage basin (Richardson 1994). In contrast, acidification of seepage lakes may increase MeHg levels in fish because methylation is promoted at low pH (Miskimmin et al. 1992; Richardson 1994).

The efficiency of MeHg uptake across the gills of fish is inversely correlated with the calcium ion content of water (Rodgers and Beamish 1983). The water hardness relationship (calcium) has particular relevance in Alberta, where many lakes and rivers are high in calcium and pH. This may explain why fish in many Alberta reservoirs do not have elevated mercury concentrations even though other water bodies in western Canada do. Uptake would be reduced in these high calcium, high pH water bodies compared with softwater lakes. Water hardness makes little difference to mercury toxicity (Keller and Zam 1991). Selenium (Se) may effectively block the accumulation of MeHg in soft, as well as harder, more alkaline waters (Southworth et al. 1994). Accumulation of MeHg by fish in temperate waters is most rapid during summer when the feeding and metabolic rates of fish are most rapid and when the production of MeHg is greatest (Spry and Wiener 1991).

---

<sup>9</sup> High pH and alkalinity are often associated with low Hg levels in fish.

<sup>10</sup> In seepage lakes that have little terrestrial runoff, DOC is less important than pH and alkalinity, and is negatively correlated with MeHg concentrations in fish presumably by inhibiting biotic methylation (Richardson 1994).

**Table 7. Variables associated with mercury concentrations in fish**

| Taxa                                     | Location                       | Fish [Hg] related to                                | Reference              |
|--|--------------------------------|---|------------------------|
| Yellow Perch (Yearling)                  | Muskoka-Haliburton, ON         | pH(-), $A_{dr.bas.}/A_{lake}(+)$                    | Suns & Hitchin 1990    |
| Perch, Pike, Sucker, Bass                | Michigan (UP)                  | pH(-), alkalinity(-), DOC in seepage lakes only (-) | Grieb et al. 1990      |
| Yellow Perch (2 years old)               | Wisconsin                      | pH(-), profundal sediment [Hg](+)                   | Cope et al. 1990       |
| Walleye                                  | 79 Ontario lakes               | DOC(+), Fe(+)                                       | Wren et al. 1991       |
| Northern Pike                            | 79 Ontario lakes               | DOC(+), Fe(+), pH(-), alkalinity(-)                 | Wren et al. 1991       |
| Sunfish                                  | 16 South Central Ontario Lakes | pH(-)   | Wren & MacCrimmon 1983 |
| Yellow Perch                             | 6 N. Ont. Shield lakes         | Lake size (-); Epilimnetic temperature (+)          | Bodaly et al. 1993     |
| Pike                                     | Sweden                         | pH(-), sed. [Hg](+), prox. To emission sources (+)  | Håkanson et al. 1988   |
| Lake trout, Sucker, Perch, Pike, Walleye | 14 Ontario lakes               | Fish weight(+), atmospheric Hg loading(+)           | Johnson 1987           |
| Yellow Perch (3-5 years)                 | 16 Adirondack lakes            | DOC(+), % wetlands(+), total dissolved Al(+)        | Driscoll et al. 1994   |
| Yellow Perch - MeHg                      | 7 Wisconsin seepage lakes      | BCF:DOC(-)  | Watras et al. 1994     |
| Yellow Perch - THg                       | 7 Wisconsin seepage lakes      | BCF:pH(-)   | Watras et al. 1994     |

#### 6.4 Accumulation by Semi-Aquatic Animals and Birds

Semi-aquatic animals and birds that consume aquatic biota bioaccumulate MeHg from their diet, similar to fish. Piscivorous mammals like mink and otter, and birds such as loons, herons and osprey, are top predators in ecosystems and may bioaccumulate substantial MeHg from their diet. Some birds, like the American wigeon (*Anas americana*), tend to accumulate less MeHg because of a diet consisting mostly of aquatic plants (Vermeer et al. 1973). The mechanism for transfer of inorganic Hg and MeHg from the gut to other tissues is the same as described for fish (section 6.3).

Organisms at higher trophic levels also tend to live longer, have more complex physiology and more organs that MeHg will accumulate in. Researchers usually find a lower proportion of MeHg in liver and kidney than in other tissues. While demethylation has been suggested as a mechanism for low MeHg in these organs (Wren et al. 1986), other possibilities may include tissue redistribution or selective excretion.

Mercury levels in mink and otter are elevated relative to the fish or other aquatic biota they consume. Kucera (1983) found that THg was an order of magnitude higher in mink and otter than in fish from the same river system in Manitoba. Somewhat lower average BAFs of 3.9 and 3.4 for mink and otter, respectively, were reported in a wide-ranging study of fish and mustelid populations in New York state (Foley et al. 1988).

Mercury may be passed from adult females to their offspring. Almost 100% of mercury transferred from adult loons through eggs to chicks was organic with no net loss of MeHg from chick tissue as the embryo developed. Methylmercury levels in eggs and in the brain of newly hatched chicks often exceeded levels in the mother's brain (Barr 1986). The transfer of Hg to eggs has been suggested as a method of Hg excretion (Heinz and Hoffman 1998). DesGranges et al. (1998) studied ospreys (*Pandion haliaetus*) in northern Québec and found that adults had higher total mercury concentrations in feathers than their fledglings and that there was no difference in the number of young that fledged per active nest between those located near reservoirs and those located near natural lakes and rivers.

Selenium has long been considered to be a protective agent against mercury toxicity (Wren 1984), a factor that will be discussed further in chapter 8 (Toxicity). Se may also be one factor that reduces MeHg accumulation according to the unexpected finding of low mercury in the tissues of mink and otter near Sudbury compared to other locations in Ontario. The mammals from the metal-contaminated Sudbury environment contained high levels of selenium (Wren and Stokes 1988). Mercury levels in fish from the Sudbury region were also lower than expected (Scheuhammer and Blancher 1994), shifting the potential selenium effect at least one step down on the food web in this case. High levels of sulphide would also be found in the Sudbury area. The solubility products ( $K_s$ ) of HgSe and HgS are  $10^{-58}$  and  $10^{-52}$ , respectively (WHO 1990), which would indicate a strong binding of free mercury ions and a reduction of availability for methylation and accumulation. The Hg-Se ratio has shown an inconsistent tendency to be higher in freshwater fish than in marine fish, which may be an indication of the higher relative mercury levels in certain freshwater environments (Pelletier 1985).

Feather moult, exposure time and residual tissue levels from one season to the next for migratory piscivorous birds are important determinants of mercury body burdens. Hg concentrations were found to decrease in Bonaparte's gulls (*Lacus philadelphia*) as moult progressed over the season (Braune and Gaskin 1987). Loons may spend a summer on a mercury-contaminated territory, then undergo a complete moult including renewal of primary feathers before migrating to winter breeding grounds (Barr 1986). Growing feathers may concentrate mercury up to 12 and 22 times more than breast muscle in adult and juvenile pintail ducks, respectively (Vermeer and Armstrong 1972). In loons, both back and belly feathers contained five to six times the mercury levels in breast muscle. Back feathers of adult loons collected in Ontario from 1971 to 1974 contained an average of 3.8 to 13.4 mg·kg<sup>-1</sup>, belly feathers contained 10.7 to 14.9 mg·kg<sup>-1</sup>, while breast muscle contained 1.02 to 3.26 mg·kg<sup>-1</sup> (Frank et al. 1983). Thus, individual birds that winter on relatively mercury-free waters, should arrive on the breeding grounds with internal mercury levels lower than at any other time of year.

Methylmercury is preferentially retained in dark muscle rather than white muscle fibre of birds (c.f. Barr 1986). Dark and myoglobin-rich muscle fibre is characteristic of loons. Any residual tissue loads from either the previous season or contaminated winter habitats may be augmented sufficiently by contaminated prey (0.35 to 0.50 µg·g<sup>-1</sup>) in time to interfere with reproductive behaviour. For example, no loons hatched chicks in territories with yellow perch that had mercury levels exceeding 0.36 µg·g<sup>-1</sup>, while only one egg was laid in lakes from which mean Hg levels in perch exceeded 0.4 µg·g<sup>-1</sup> (Barr 1986). These observations were from the

## Canadian Tissue Residue Guidelines: Methylmercury

mercury contaminated area of the English-Wabigoon River system in north-west Ontario. Other toxic effects will be discussed in Chapter 8.

## 7.0 OVERVIEW OF DISTRIBUTION AND LEVELS OF METHYLMERCURY IN THE CANADIAN ENVIRONMENT

This chapter describes methylmercury concentrations in aquatic plants, aquatic invertebrates, fish, amphibians, reptiles, semi-aquatic mammals, marine mammals, aquatic birds and non-biological media. Because of the quantity of measurements, note that the information contained in this chapter and tables are examples only and are not exhaustive. Instead, readers are referred to such databases as are being developed by Environment Canada (Kent et al. 1998), and to those that have been compiled for many years by some of the provinces (e.g., Ontario and Québec). Where MeHg concentrations for Canada are unavailable or scarce, information is supplemented with total mercury (THg) concentrations. In the case of fish, researchers routinely only measure THg because it is believed that essentially all mercury in fish tissue is MeHg. Each section is supported by detailed tables contained in Appendix B, with the number listed next to each sub-heading.

### 7.1 Aquatic Plants (App. B-1)

With a few exceptions, the available data indicate that methylmercury usually represents less than 50% of total mercury in freshwater aquatic plants. One study in Heney Lake, Ontario reported levels of 80 to 90  $\mu\text{g}\cdot\text{kg}^{-1}$  THg<sup>11</sup>, and 50  $\mu\text{g}\cdot\text{kg}^{-1}$  MeHg in two genera of filamentous algae (*Mougeotia* and *Spirogyra*), representing 55% MeHg (Stokes et al. 1983). In another lake from the same study (Swan Lake), MeHg represented less than 50% of THg from the same algal groups.

A recent study completed at the Experimental Lakes Area (near Kenora, ON) found that the percentage of MeHg depended on the type of tissue from *Sphagnum augustifolium* that was measured. The lower stem, upper stem, and capitulum contained 0.3%, 1.4% and 2.7% MeHg, respectively (0.3, 1.6, 2.5  $\mu\text{g}\cdot\text{kg}^{-1}$  dw; Moore et al. 1995). They also found that the type of habitat appeared to affect the levels of MeHg in *Sphagnum* spp. Species from a pool/stream habitat contained about 20  $\mu\text{g}\cdot\text{kg}^{-1}$  (19%) MeHg, while species from hollow/lawn and hummock habitats contained about 1 and 0.5  $\mu\text{g}\cdot\text{kg}^{-1}$ , respectively (2.5% and 1.4% MeHg). The macrophyte and sedge species, *Carex aquatilis*, *C. rostrata* and *Sparganium* spp. contained about 2  $\mu\text{g}\cdot\text{kg}^{-1}$  MeHg, on average, representing 37% of THg (Moore et al. 1995).

Macrophytes from the Ottawa River contained lower percentages of MeHg in their roots than their shoots. The roots of *Elodea canadensis*, *Sagittaria latifolia*, *Sparganium augustifolium*, and *Sparganium eurycarpum* contained 8 to 14% (mean = 10.3%) MeHg compared with 23 to 48% (mean = 31.7%) MeHg in their shoots (Mortimer 1985). The levels of THg were 48 to 550  $\mu\text{g}\cdot\text{kg}^{-1}$ , while the specific MeHg levels were not reported. The highest level of total mercury in aquatic plants found in the literature was 1600  $\mu\text{g}\cdot\text{kg}^{-1}$  (min. = 17  $\mu\text{g}\cdot\text{kg}^{-1}$ ), in *Myriophyllum* spp. collected in 1978 from the St. Lawrence River, near Cornwall (Mortimer 1985). *Myriophyllum* from the same area of the St. Lawrence River in 1994 contained THg concentrations in the range of 63 to 240  $\mu\text{g}\cdot\text{kg}^{-1}$  (Thompson-Roberts et al. 1999).

No data were located on levels of methylmercury in estuarine or marine plants in the Canadian environment.

---

<sup>11</sup> All plant measurements on a dw basis unless noted otherwise.

## 7.2 Aquatic Invertebrates (App. B-2)

Many measurements of THg in aquatic invertebrates in Canadian freshwater environments have been made (App. B-2a), but it was not until recently that researchers began to report MeHg in invertebrates. Generally, the proportion of MeHg in invertebrates is higher than in aquatic plants.

In a wide-ranging study in northern Québec, Tremblay et al. (1996) found from 6 to 100% of THg in benthic invertebrates was MeHg. The two highest measured levels of MeHg were found in corixids (*Sigara* spp.) in the La Grande-2 Reservoir. In the south arm of the reservoir, the MeHg concentrations in the invertebrates averaged  $1.52 \mu\text{g}\cdot\text{g}^{-1}$  dw (representing 91% of THg) and in the north arm levels averaged  $1.37 \mu\text{g}\cdot\text{g}^{-1}$  (representing 76% of THg). Larval dragonflies (*Somatochlora* spp.) in the south arm contained 100% MeHg, with levels averaging  $0.62 \mu\text{g}\cdot\text{g}^{-1}$ , while for the same organism in the north arm, only 75% of THg was MeHg. In a control lake, Duncan L, concentrations in various benthic invertebrates ranged from  $0.014 \mu\text{g}\cdot\text{g}^{-1}$  (mayfly, *Leptophlebia* spp.), to  $0.124 \mu\text{g}\cdot\text{g}^{-1}$  (corixid, *Sigara* spp.), with the proportion of MeHg ranging from 11% to 75%. See Appendix B-2 for specific results of this survey and Tremblay (1998) for further discussion of MeHg levels in invertebrates from lakes and reservoirs in northern Québec.

A study of twenty-four lakes in south-central Ontario lakes reported *Daphnia* species to contain 0.019 to  $0.448 \mu\text{g}\cdot\text{g}^{-1}$  dw MeHg (Westcott and Kalff 1996). They did not measure THg, so the proportion represented by MeHg could not be calculated (K. Westcott 1999, pers. com.).

Malley et al. (1996) found that while the kidney of the mussel, *Pyganodon grandis*<sup>12</sup> contained higher concentrations of MeHg than other tissues, the proportion of MeHg in the kidney was the lowest, at 15%; of  $1.99 \mu\text{g}\cdot\text{g}^{-1}$  dw THg in the kidney,  $0.29 \mu\text{g}\cdot\text{g}^{-1}$  was MeHg. The whole organism contained 44% MeHg, while the mantle, viscera, gill and foot contained 45%, 47%, 58% and 62% MeHg, respectively. This represented a range of  $0.11$  to  $0.25 \mu\text{g}\cdot\text{g}^{-1}$  MeHg (Malley et al. 1996). All mussels were from an unmanipulated source lake (Lake 104) at the Experimental Lakes Area, northern Ontario.

Organisms from the contaminated Wabigoon River/Clay Lake area of north-western Ontario showed among the highest levels of THg in freshwater invertebrates. For example, the crayfish, *Orconectes virilis*, from the east basin of Clay Lake collected in 1979-80 contained  $2.2 \pm 0.66 \mu\text{g}\cdot\text{g}^{-1}$  ww THg, and from the Wabigoon River (inflow to Ball Lake) contained  $1.7 \pm 0.49 \mu\text{g}\cdot\text{g}^{-1}$  ww THg (Parks et al. 1991). These are high levels for invertebrates considering that the consumption guideline for human consumers of fish in Canada is  $0.5 \mu\text{g}\cdot\text{g}^{-1}$  ww<sup>13</sup>. If the levels for the crayfish were converted to a dry weight basis, they would be even higher. For example, if tissues were considered at least 50% water, THg concentrations would be twice as high as reported above.

There are relatively few studies in Canada that look at mercury concentrations in marine invertebrates (App. B-2b). Concentrations of THg in the estuarine mussel, *Mytilus edulis*, were reported in areas of the St. Lawrence Estuary for the year of cessation of chlor-alkali plant activities and one year later (1976-1977). Concentrations at the mouth of Saguenay Fjord declined from  $0.63 \mu\text{g}\cdot\text{g}^{-1}$  dw to  $0.17 \mu\text{g}\cdot\text{g}^{-1}$  dw during this period. Similarly, levels of THg declined in two locations on the south shore of the lower estuary and in the Gaspé Peninsula during the same period (Cossa and Rondeau 1985; Appendix B-2).

<sup>12</sup> Formerly *Anodonta grandis grandis*

<sup>13</sup> To be discussed in more detail in chapter 9.



### 7.3 Fish (App. B-3)

As previously described, mercury in fish from natural environments is comprised almost exclusively of MeHg regardless of the composition of diet sources and exposure to water. As such, the data tabulated includes numerous measurements of THg only, which are assumed to be MeHg concentrations unless specifically reported as measured MeHg. While the data included are not all-inclusive, information on freshwater fish mercury levels from eight provinces (Alberta, British Columbia, Manitoba, New Brunswick, Newfoundland, Nova Scotia, Ontario, Québec) and one territory (Northwest Territories) are included.

In some cases, water bodies were either directly contaminated with mercury by mine tailings or industry, and in other cases, MeHg levels became elevated as a result of hydroelectric reservoir development, or by high geologic levels of mercury. As expected, levels are generally highest in piscivorous species such as lake trout, northern pike, walleye and bull trout. The literature is dominated by reports on freshwater fish in comparison to marine fish. All levels are reported as  $\mu\text{g}\cdot\text{g}^{-1}$  wet weight. Age and/or size class of the fish are tabulated where known; wildlife typically consume fish <35 cm in length (US EPA 1993). Results are discussed in alphabetical order by province or territory.

Monitoring programs in Canada typically measure mercury levels in large, commercial or sport fish to address human health concerns. Larger (i.e., older) fish, regardless of other influences, are expected to contain higher levels of mercury because of their longer exposure period relative to smaller (i.e., younger) fish consumed by wildlife. As such, levels reported below may be higher than those expected for prey species of wildlife. Care must be taken when interpreting the relevance of data for large fish to wildlife. The reader is cautioned against generalising the complex relationship between fish size and mercury concentration. While fish size has been shown to be the single best predictor of mercury body burdens, estimating mercury concentrations in small fish from measured concentrations in large fish depends on species and trophic level as well as on site-specific environmental (e.g., pH, alkalinity, DOC, basin/lake morphology) and physiological (e.g., uptake/elimination rates, assimilation efficiency) factors (see chapter 6).

#### 7.3.1 Freshwater fish (App. B-3a)

In Alberta, high levels of mercury in fish were reported from lakes contaminated with mine tailings. Lake trout (*Salvelinus namaycush*), northern pike (*Esox lucius*), and round whitefish (*Prosopium cylindraceum*) collected in 1977 from Giauque Lake in northern Alberta contained averages of 3.79, 1.75 and 1.22  $\mu\text{g}\cdot\text{g}^{-1}$  ww, respectively (Moore and Sutherland 1980). In another contaminated lake, Thompson Lake, northern pike collected in 1978 had an average of 1.7  $\mu\text{g}\cdot\text{g}^{-1}$  THg in muscle tissue. Regardless of whether a lake was contaminated or not, lake whitefish (*Coregonus clupeaformis*) had low average mercury concentrations of 0.2  $\mu\text{g}\cdot\text{g}^{-1}$ . In a number of control lakes, lake trout and northern pike muscle concentrations ranged from 0.11 to 1.0 and 0.05 to 1.91  $\mu\text{g}\cdot\text{g}^{-1}$ , respectively (Moore and Sutherland 1980).

Ramamoorthy et al. (1985) reported mercury levels in seven fish species from the North Saskatchewan River in the vicinity of Edmonton. Average concentrations ranged from 0.245  $\mu\text{g}\cdot\text{g}^{-1}$  for longnose sucker (*Catostomus catostomus*) to 0.743  $\mu\text{g}\cdot\text{g}^{-1}$  for sauger (*Stizostedion canadense*). Two other species that exceeded the consumption guideline of 0.5  $\mu\text{g}\cdot\text{g}^{-1}$  were goldeye (*Hiodon alosoides*; 0.594  $\mu\text{g}\cdot\text{g}^{-1}$ ) and walleye (0.645  $\mu\text{g}\cdot\text{g}^{-1}$ ).

Ramamoorthy et al. (1985) also reported the proportion of MeHg to range from 77% (longnose sucker) to 95% (walleye)<sup>14</sup>.

In British Columbia, mercury in fish has been quantified in a number of lakes and rivers, as well as Williston Lake (reservoir associated with the Bennett Dam on the Peace River). Studies by BC Hydro on a number of other reservoirs are in progress (Watson 1992). The highest mercury concentrations were found in lake trout from Pinchi Lake, which is located in an area of high geological mercury and a former mercury mine. Mercury in these lake trout ranged from 1.06 to 5.78  $\mu\text{g}\cdot\text{g}^{-1}$  (Watson 1992). Bull trout (*Salvelinus confluentus*) levels were relatively high both in Pinchi Lake (0.49 to 0.75  $\mu\text{g}\cdot\text{g}^{-1}$ ) and in Williston Lake (0.25 to 1.62  $\mu\text{g}\cdot\text{g}^{-1}$ ). Kokanee (*Oncorhynchus nerka kennerlyi*) and rainbow trout (*Oncorhynchus mykiss*) from Pinchi Lake contained moderate to high levels at 0.32 to 0.76  $\mu\text{g}\cdot\text{g}^{-1}$  and 0.20 to 0.40  $\mu\text{g}\cdot\text{g}^{-1}$ , respectively.

In a number of rivers in northern BC, mercury concentrations ranged from 0.02 to 0.78  $\mu\text{g}\cdot\text{g}^{-1}$ . On average, rainbow trout and lake whitefish had the lowest levels, while burbot (*Lota lota*) and bull trout had higher levels (Watson 1992).

In Manitoba, extensive monitoring of mercury in fish has been done in association with reservoirs. Average mercury concentrations associated with the Churchill River diversion and flooding (Notigi Reservoir) including Southern Indian Lake exceeded 0.5  $\mu\text{g}\cdot\text{g}^{-1}$  in all species except lake whitefish (Bodaly et al. 1984<sup>15</sup>). Average mercury levels exceeded 2  $\mu\text{g}\cdot\text{g}^{-1}$  in walleye and northern pike from the Rat Lake portion of the Notigi Reservoir, and exceeded 1  $\mu\text{g}\cdot\text{g}^{-1}$  in most other areas of the Notigi Reservoir. Levels were also elevated (>0.75  $\mu\text{g}\cdot\text{g}^{-1}$ ) in these species in five lakes downstream of the Notigi Reservoir. In Southern Indian Lake, walleye and northern pike had average muscle concentrations of 0.57  $\mu\text{g}\cdot\text{g}^{-1}$  and 0.68  $\mu\text{g}\cdot\text{g}^{-1}$ , respectively, while lake whitefish averaged 0.13  $\mu\text{g}\cdot\text{g}^{-1}$  (Bodaly et al. 1984).

Relatively fewer data are available for freshwater fish in the Atlantic provinces, although the dataset has recently grown because of a concern for loon populations in some areas. In a multi-agency study in Kejimikujik National Park, NS, hundreds of measurements in fish tissue have been made (report publications pending, Mercury Team 1998). Two land-locked salmon (*Salmo salar*)<sup>16</sup> from Sisson Branch Reservoir in the St. John River basin (NB), measuring 38 and 58 cm and weighing less than 1.5 kg, contained 0.84 and 1.5  $\mu\text{g}\cdot\text{g}^{-1}$  THg, respectively (Bailey 1985). Twelve lake whitefish from the Green River Reservoir (NB) averaged 0.12  $\mu\text{g}\cdot\text{g}^{-1}$ . In the Smallwood Reservoir in Labrador during the 1970s, land-locked salmon reportedly contained 1.5 to 2.3  $\mu\text{g}\cdot\text{g}^{-1}$  (Bruce and Spencer 1978, c.f. Bailey 1985). Mercury measured in 1977 and 1978 in northern pike from Labrador lakes ranged from 0.01 to 3.08  $\mu\text{g}\cdot\text{g}^{-1}$  (n=117), in lake trout from 0.12 to 3.37  $\mu\text{g}\cdot\text{g}^{-1}$  (n=107) and in brook trout from 0.01-1.69  $\mu\text{g}\cdot\text{g}^{-1}$  (n=96; Bruce et al. 1979).

In Wreck Cove Reservoir, NS, brook trout weighing from 49-211 g had muscle mercury concentrations of 0.27-0.73  $\mu\text{g}\cdot\text{g}^{-1}$ , averaging  $0.50 \pm 0.16 \mu\text{g}\cdot\text{g}^{-1}$  (ADI Nolan Davis Inc. 1994). The anadromous striped bass (*Morone saxatilis*) in the Annapolis River had elevated Hg in muscle tissue ranging from 0.26 - 1.97  $\mu\text{g}\cdot\text{g}^{-1}$  (mean 0.77, mean weight of 7.4 kg); lower for smaller specimens (mean 1.8 kg) from the Shubenacadie River ranging from 0.16 - 1.44  $\mu\text{g}\cdot\text{g}^{-1}$  (mean 0.40  $\mu\text{g}\cdot\text{g}^{-1}$ ; Ray et al. 1984).

---

<sup>14</sup> Note that recent methods and findings stress that 98-100% of THg in fish muscle is MeHg (Bloom 1992).

<sup>15</sup> Others have reported on mercury in these systems; the summary of Bodaly et al. 1984 is used for brevity here.

<sup>16</sup> Species not reported, Atlantic salmon is professional judgement of species based on location.

In the Northwest Territories, mercury concentrations in a variety of species averaged below  $0.5 \mu\text{g}\cdot\text{g}^{-1}$ , with some exceptions. Lake trout from Kaminak Lake, west of Hudson Bay, contained 0.57 to  $2.0 \mu\text{g}\cdot\text{g}^{-1}$  THg (Shilts and Coker 1995). The Kaminak Lake area has high geological mercury. For unknown reasons, northern pike, lake trout and walleye have elevated mercury (means of 0.74, 1.34,  $1.49 \mu\text{g}\cdot\text{g}^{-1}$ ) in Lac Ste. Thérèse in the Johnny Hoe River system (Stephens 1995). The high levels have persisted over time, with measurements in walleye being made in 1975, 1980, 1992 and 1993. Walleye, but not pike or lake whitefish have average concentrations that exceed  $0.5 \mu\text{g}\cdot\text{g}^{-1}$  in another lake in the system, Tseepantee Lake. Two additional lakes (Keller and Taché) have fish with relatively lower THg concentrations.

In the Hay River, NWT, lake whitefish, walleye and northern pike averaged 0.07, 0.22 and  $0.32 \mu\text{g}\cdot\text{g}^{-1}$ , respectively. Similar levels were found for these species in the Slave River and in Leland Lake (Grey et al. 1995).

More studies have been completed in Ontario for a wide range of water bodies than in most other regions of Canada. The Ontario Ministry of the Environment maintains a very large database of mercury concentrations in fish ("probably the largest in the world", G. Mierle 1999, MOE, pers. com.). The most highly mercury contaminated area in the country is likely the English-Wabigoon River system in north-western Ontario. The source of mercury was a chlorine-alkali plant located in the town of Dryden (Rudd et al. 1983). Very high levels of mercury in Clay Lake burbot and walleye were 21.9 and  $15.7 \mu\text{g}\cdot\text{g}^{-1}$  on average, respectively (Fimreite and Reynolds 1973). In Ball Lake, downstream of Clay lake, walleye, rock bass (*Ambloplites upestris*) and northern pike averaged 4.5, 6.2 and  $7.7 \mu\text{g}\cdot\text{g}^{-1}$ . White suckers (*Catostomus commersoni*) from Clay Lake and the Wabigoon River had mercury levels in excess of 3 and  $4 \mu\text{g}\cdot\text{g}^{-1}$ , respectively. Fish from many other connected lakes and rivers also had elevated mercury concentrations (Fimreite and Reynolds 1973).

Cabana et al. (1994) found a positive correlation between lake trout mercury concentrations and length of the food chain in large number of Ontario lakes. Lake trout from lakes with pelagic forage fish and *Mysis* shrimp had an average of  $0.65 \mu\text{g}\cdot\text{g}^{-1}$  compared with  $0.18 \mu\text{g}\cdot\text{g}^{-1}$  in lake trout from lakes that contained *Mysis* but lacked forage fish. The relationship did not exist for smallmouth bass (*Micropterus dolomieu*), a species that does not exploit the pelagic food chain, for which THg averaged from 0.43 to  $0.53 \mu\text{g}\cdot\text{g}^{-1}$  regardless of the length of the food chain.

Lake trout had high mercury concentrations relative to rainbow smelt (*Osmerus mordax*) in a series of Ontario lakes examined from 1977 to 1981. Lakes Muskoka, Mary and Vernon were rated high mercury because lake trout concentrations averaged 2.96 to  $4.49 \mu\text{g}\cdot\text{g}^{-1}$  (0.34 to 0.49 for smelt). Lakes Koshlong, Rosseau and Tadenac were rated moderate because lake trout concentrations averaged 0.61 to  $0.99 \mu\text{g}\cdot\text{g}^{-1}$  (0.18 to 0.26 for smelt), and Lakes Simcoe, Joseph and Bella were rated low Hg because of lake trout concentrations of 0.39 to  $0.55 \mu\text{g}\cdot\text{g}^{-1}$  (0.08 to 0.14 for smelt; MacCrimmon et al. 1983).

Northern pike from 170 lakes and walleye from 255 lakes from across Ontario indicate that fish of a certain size from seemingly unpolluted systems overall average at or above the guideline level of  $0.5 \mu\text{g}\cdot\text{g}^{-1}$ . Northern pike of approximately 54 cm in length averaged  $0.50 \mu\text{g}\cdot\text{g}^{-1}$  (0.07 to  $1.28 \mu\text{g}\cdot\text{g}^{-1}$ ), and walleye of approximately 41 cm in length averaged  $0.58 \mu\text{g}\cdot\text{g}^{-1}$  (0.09 to  $3.24 \mu\text{g}\cdot\text{g}^{-1}$ ) over a wide variety of lakes (Wren et al. 1991).

Yearling yellow perch collected both from the Ottawa River and a number of lakes in north-western Ontario had consistently low levels of mercury. Perch collected in 1974 and 1975 from the Ottawa River contained  $0.13 \mu\text{g}\cdot\text{g}^{-1}$  (Rodgers and Qadri 1982) and from six remote lakes

from NW Ontario averaged 0.04 to 0.14  $\mu\text{g}\cdot\text{g}^{-1}$  (Bodaly et al. 1993). Similarly, yearling yellow perch collected from 16 lakes in the Muskoka-Haliburton area contained 0.031-0.233  $\mu\text{g}\cdot\text{g}^{-1}$  (Suns and Hitchin 1990).

In Québec, studies have focused on northern hydroelectric projects in the Great Whale, La Grande, and Nottaway-Broadback-Rupert (NBR) project areas. The highest levels were found in northern pike and walleye from the La Grande-2 Reservoir. Fish collected from 1982 to 1988 had average muscle concentrations of 1.31 to 2.99  $\mu\text{g}\cdot\text{g}^{-1}$  (pike), and 1.92 to 2.80 (walleye), with the highest averages occurring nine years after flooding (Verdon et al. 1991). Longnose sucker concentrations in the La Grande-2 reached a high of 0.67  $\mu\text{g}\cdot\text{g}^{-1}$  and lake whitefish a high of 0.57  $\mu\text{g}\cdot\text{g}^{-1}$  five years after flooding. Twenty-nine control lakes in the region indicated that fish accumulated lower amounts of mercury with concentrations in northern pike of 0.25 to 0.90, walleye of 0.32 to 1.26 and longnose sucker of 0.06 to 0.32  $\mu\text{g}\cdot\text{g}^{-1}$  (Verdon et al. 1991). More recent publications summarise fish mercury levels at the La Grande complex from 1978 to 1994 (Hydro-Québec and Groupe-conseil Génivar Inc. 1996, 1997).

In the Great Whale and NBR project areas of northern Québec, average mercury concentrations in piscivorous fish exceeded the 0.5  $\mu\text{g}\cdot\text{g}^{-1}$  level. Northern pike and walleye from the NBR region on the south-east coast of James Bay had levels of  $0.93 \pm 0.51$  and  $0.78 \pm 0.46$   $\mu\text{g}\cdot\text{g}^{-1}$ , respectively. Mercury concentrations in these species were only slightly lower in the Great Whale area on the east coast of Hudson Bay (Langlois et al. 1995). Additional data for these regions may be found in Hydro-Québec and Groupe-conseil Génivar Inc. (1997).

Measurements of mercury in northern pike and walleye from 235 lake and 214 river sites across Québec indicated that average levels exceeded 0.5  $\mu\text{g}\cdot\text{g}^{-1}$  in many cases. Northern pike between 40-50, 55-70, and >70 cm measured 0.4, 0.66, and 1.12  $\mu\text{g}\cdot\text{g}^{-1}$ , respectively. Walleye between 30-40, 40-50, and >50 cm averaged 0.53, 0.78, and 1.26  $\mu\text{g}\cdot\text{g}^{-1}$ , respectively (Laliberté 1996).

### 7.3.2 Marine fish (App. B-3b)

A few data for methylmercury in marine fish in the Canadian environment will be discussed. Atlantic herring (*Clupea harengus harengus*) collected in 1981 from the south-western Bay of Fundy had low levels of mercury ranging from 0.005 to 0.015  $\mu\text{g}\cdot\text{g}^{-1}$ . The fish were one to five years in age, and mercury in muscle and whole body was found to increase with age (Braune 1987a). Greenland cod (*Gadus ogac*, 400 mm) caught in the coastal waters of James Bay between 1987 and 1994 contained from 0.14 to 0.42  $\mu\text{g}\cdot\text{g}^{-1}$  of mercury, while the fourhorn sculpin (*Myoxocephalus quadricornis*, 250 mm) from the same regions contained from 0.1 to 0.55  $\mu\text{g}\cdot\text{g}^{-1}$  (Hydro-Québec and Groupe-conseil Génivar Inc. 1997).

### 7.4 Amphibians (App. B-4)

No data were located on levels of methylmercury in amphibians in the Canadian environment. One recent study examined levels of THg in the mudpuppy, *Necturus maculosus*, from various locations in the St. Lawrence and Ottawa Rivers. Average concentrations ranged from 0.087 to 0.239  $\mu\text{g}\cdot\text{g}^{-1}$  on a wet weight (ww) basis, with a maximum individual concentration of 0.445  $\mu\text{g}\cdot\text{g}^{-1}$  ww (Bonin et al. 1995). The high concentration was an adult female from the Beauharnois area of the St. Lawrence River; whole body with gonads removed. The average whole body concentration (without gonads) was  $0.239 \pm 0.137$ , while the gonads of the same six individual females averaged 0.068  $\mu\text{g}\cdot\text{g}^{-1}$  ww.

## 7.5 Reptiles (App. B-4)

No data were located on levels of methylmercury in reptiles in the Canadian environment. Snapping turtle eggs (*Chelydra serpentina*) collected in 1989 and 1990 from the St. Lawrence River and tributaries contained relatively low concentrations of THg (0.09 to 0.15  $\mu\text{g}\cdot\text{g}^{-1}$  ww THg; Bonin et al. 1995).

## 7.6 Mammals

### 7.6.1 Semi-aquatic Mammals (App. B-5)

No data were located on levels of methylmercury in semi-aquatic mammals in the Canadian environment. Several studies that report total mercury concentrations will be described briefly.

In a number of locations in Manitoba, mink (*Mustela vison*) and otter (*Lutra canadensis*) accumulated higher THg concentrations in liver tissue than in kidney or brain tissue. Average liver concentrations ranged from 0.82 to 3.93  $\mu\text{g}\cdot\text{g}^{-1}$  ww for mink, and from 1.75 to 3.67 for otter. Average kidney concentrations ranged from 0.67 to 2.98  $\mu\text{g}\cdot\text{g}^{-1}$  THg for mink, and from 1.07 to 2.37  $\mu\text{g}\cdot\text{g}^{-1}$  for otter. The maximum THg concentration in brain tissue was 0.97  $\mu\text{g}\cdot\text{g}^{-1}$  for mink, and 0.92  $\mu\text{g}\cdot\text{g}^{-1}$  for otter (Kucera 1983). The highest levels tended to be from animals in the vicinity of the Winnipeg River.

In Ontario, MeHg percentages for various tissues in mink and otter from the English River, Muskoka and Sudbury areas were reported. Mink livers and kidneys contained 53% and 43% MeHg, respectively, based on THg concentrations ranging up to 7.5 and 5.54  $\mu\text{g}\cdot\text{g}^{-1}$ , respectively. Otter livers, kidneys and muscle contained 34%, 19%, and 90% MeHg, respectively, based on THg concentrations ranging up to 17.4, 12.6, and 4.26  $\mu\text{g}\cdot\text{g}^{-1}$ , respectively (the highest concentrations were from the English River; Wren et al. 1986). Details of THg concentrations in all tissues are reported in Appendix B-5.

In an Ontario study of mink and otter, THg in liver, kidney and brain tissue showed similar trends as the Manitoba study of decreasing concentration with type of tissue. For mink, the highest average liver tissue concentration was  $2.55 \pm 1.96$  (range of 0.56 to 6.9)  $\mu\text{g}\cdot\text{g}^{-1}$  from the contaminated English River district of north-western Ontario. For otter, the highest average liver tissue concentration was  $3.47 \pm 4.1$  (range of 0.71 to 17.4)  $\mu\text{g}\cdot\text{g}^{-1}$  from the same area (Wren et al. 1986). Muscle concentrations in the province-wide study ranged from 0.38 to 1.62  $\mu\text{g}\cdot\text{g}^{-1}$  for mink, and 0.30 to 1.04 for otter.

In northern Québec, mink from the Great Whale area had the highest average muscle tissue concentrations of  $2.40 \pm 2.24$   $\mu\text{g}\cdot\text{g}^{-1}$  THg (Langlois et al. 1995). Mink from natural environments of northern Québec had liver, kidney and brain tissue concentrations of 4.85, 2.45 and 0.93  $\mu\text{g}\cdot\text{g}^{-1}$  THg (Bélanger and Larivière 1997).

Other semi-aquatic mammals that have been monitored for THg include beaver (*Castor canadensis*) and racoon (*Procyon lotor*). Liver, kidney and muscle tissue from beaver collected near Parry Sound, ON, contained 0.03  $\mu\text{g}\cdot\text{g}^{-1}$  THg. This low level reflects the non-fish diet of the beaver. Liver, kidney and muscle tissue from racoons collected in the same area contained 4.5, 1.1 and 0.3  $\mu\text{g}\cdot\text{g}^{-1}$  THg (Wren 1984).

### 7.6.2 Marine Mammals (App. B-6)

Few studies report on MeHg in marine mammals in the Canadian environment. Some of the highest THg levels found were measurements of marine mammal livers. The following summarises some of the elevated THg and MeHg levels (fully detailed in Appendix B-6) in marine mammal livers in many regions of Canada:

| Species  | THg<br>$\mu\text{g}\cdot\text{g}^{-1}$ | MeHg<br>$\mu\text{g}\cdot\text{g}^{-1}$ | Reference            |
|--|--|---|----------------------|
| Beluga whale (St. Lawrence R.)<br><i>(Delphinapterus leucas)</i>         | 33                                     | NR                                      | Wagemann et al. 1996 |
| Ringed seal (Western Arctic)<br><i>(Phoca hispida)</i>                   | 33                                     | NR                                      | Wagemann 1995        |
| Harbour seal <sup>17</sup> (Great Whale Area)<br><i>(Phoca vitulina)</i> | 29                                     | 4.6                                     | Langlois et al. 1995 |
| Pilot Whale (Atlantic Coast)<br><i>(Globicephala malaena)</i>            | 23                                     | NR                                      | Wagemann et al. 1995 |
| Beluga whale (Great Whale Area)  | 20                                     | 7.3                                     | Langlois et al. 1995 |
| Narwhal (Eastern Arctic)<br><i>(Monodon monoceros)</i>                   | 11                                     | NR                                      | Wagemann et al. 1996 |
| Harp seal (Atlantic coast)<br><i>(Phoca groenlandica)</i>                | 10                                     | NR                                      | Wagemann et al. 1995 |

Livers of ringed seal (*Phoca hispida*) are consumed by indigenous people. Wagemann (1994) determined that the concentration of MeHg in ringed seal liver relative to THg was only 15% to 30%. He later argued that the MeHg percentage in ringed seal liver was much lower (only 2 to 3%) based on more accurate calculation methods (Wagemann et al. 1997; R. Wagemann 1999, DFO, pers. com.). The liver of ringed seals listed above were collected from the western Arctic between 1987 and 1993. Based on Wagemann et al. (1997) the concentration of MeHg in the above liver tissue averaged about  $1.0 \mu\text{g}\cdot\text{g}^{-1}$ . Ringed seal livers from the eastern Arctic contained  $28 \mu\text{g}\cdot\text{g}^{-1}$  THg (3% =  $0.8 \mu\text{g}\cdot\text{g}^{-1}$  MeHg; Wagemann et al. 1995; R. Wagemann 1999, DFO, pers. com.), and from the Great Whale area,  $1.1 \mu\text{g}\cdot\text{g}^{-1}$  MeHg (Langlois et al. 1995).

<sup>17</sup> Land-locked, described as freshwater seals.

There are spatial and temporal trends to mercury accumulations in Arctic marine mammals. In general, tissues of ringed seals and belugas from the western Arctic have higher mercury concentrations than those from the eastern Arctic. While this was attributed to different natural backgrounds of mercury in the two regions, it is important to note that 1) mercury in the liver of belugas increased in the western and eastern Arctic over 10 to 12 years; 2) mercury in the liver of ringed seals in the western Arctic and narwhal in the eastern Arctic showed similar increases; 3) belugas collected in the mid 1990s accumulated twice as much mercury as those collected 10 to 12 years ago; and 4) ringed seals collected in the mid-1990s accumulated three times more mercury than those collected 15 to 20 years ago (Wagemann et al. 1996).

The recent results of Wagemann et al. (1997) indicate that mercury in the muscle tissue of narwhal (*Monodon monoceros*), ringed seal and belugas (*Delphinapterus leucas*) is essentially all MeHg. On the other hand, at least one report indicated that only 34% of THg in ringed seal muscle tissue was MeHg (Langlois et al. 1995). The residues were not excessive in either case, with levels of  $0.6 \mu\text{g}\cdot\text{g}^{-1}$  MeHg (Wagemann et al. 1997) in Arctic seals, and  $0.11 \mu\text{g}\cdot\text{g}^{-1}$  in seals from the Great Whale area (Hudson Bay; Langlois et al. 1995).

Belugas in the St. Lawrence River aged from young-of-the-year to 30 years contained an average of  $33.6 \mu\text{g}\cdot\text{g}^{-1}$  THg in liver tissue (maximum of  $202 \mu\text{g}\cdot\text{g}^{-1}$ ; Wagemann et al. 1996). This was the highest recorded average mercury concentration. Methylmercury levels were not reported. In muscle, belugas from the Arctic, St. Lawrence River, and Great Whale areas contained in excess of  $2 \mu\text{g}\cdot\text{g}^{-1}$  (Langlois et al. 1995; Wagemann et al. 1997). Since much of the mercury in marine mammal muscle tissue is MeHg, belugas from many areas of Canada can be considered to have excessive MeHg in muscle tissue.

## 7.7 Birds (App. B-7)

Numerous studies report THg in various bird tissues, however, relatively few report methylmercury concentrations. While some analyses indicate the majority of mercury in tissues except liver and kidney is MeHg, there is not enough evidence to conclude, as with fish, that all mercury in birds is MeHg. Fish-eating birds commonly have a high proportion of MeHg in muscle and brain tissue. Eggs contain essentially all MeHg, which is transferred from females prior to laying.

An early study on single specimens of immature birds in the vicinity of Clay Lake, north-western Ontario (mercury contaminated) found that 69% to 99% of THg in breast tissue was MeHg. The species examined were blue-winged teal (*Anas discors*), common goldeneye (*Bucephala clangula*), common merganser (*Mergus merganser*), hooded merganser (*Lophodytes cucullatus*), and mallard (*Anas platyrhynchos*). The hooded merganser had the highest MeHg concentration with  $17.8 \mu\text{g}\cdot\text{g}^{-1}$  (99%), while the mallard had the lowest MeHg concentration with  $0.11 \mu\text{g}\cdot\text{g}^{-1}$  (69%). The teal, goldeneye and common merganser had 7.4, 12.3, and  $14.9 \mu\text{g}\cdot\text{g}^{-1}$ , respectively (Vermeer et al. 1973).

Common loons in the Algonquin Park and Parry Sound areas collected from 1971 to 1974 had higher levels of mercury in those individuals that were emaciated. Muscle tissue contained  $2.7$  to  $5.4 \mu\text{g}\cdot\text{g}^{-1}$  in emaciated loons compared to  $0.8$  to  $1.1 \mu\text{g}\cdot\text{g}^{-1}$  in healthy loons (averages; Frank et al. 1983). Mercury concentrations were similarly approximately three times higher in the brain tissue of emaciated loons compared to healthy loons (Appendix B-7).

Another study undertaken in the 1970s involved an extensive examination of the common loon (*Gavia immer*) in the Wabigoon-English River district of north-western Ontario. This is the same

vicinity as Clay Lake described above. Loon eggs contained 97% to 100% MeHg with the highest concentration of  $1.34 \pm 0.53 \mu\text{g}\cdot\text{g}^{-1}$  in a series of six lakes downstream of Dryden on the Wabigoon River system (Barr 1986). Loon chick muscle, brain and liver contained  $0.3 \mu\text{g}\cdot\text{g}^{-1}$  (82-85%),  $0.4 \mu\text{g}\cdot\text{g}^{-1}$  (99%), and  $0.8 \mu\text{g}\cdot\text{g}^{-1}$  (87-92%) MeHg, respectively. Adult loon muscle, brain and liver contained 0.9 to  $2.3 \mu\text{g}\cdot\text{g}^{-1}$  (50-86%), 0.3 to  $0.8 \mu\text{g}\cdot\text{g}^{-1}$  (52-92%), and 0.2 to 1.5 (4-27%) MeHg, respectively (Barr 1986). The pattern seems to be that the young have mainly MeHg in their tissues, and the livers of adults do not favour MeHg accumulation over other forms of mercury (similar to marine mammals). Studies from other less contaminated areas are needed to corroborate this and other conclusions.

In a recent study in Ontario, both adult and chick loon blood mercury concentrations were positively correlated with fish mercury concentrations ( $r = 0.71$  and  $0.51$ , respectively; Scheuhammer et al. 1998). Three of twenty-four lakes had 20 to 50 g perch and sunfish with Hg concentrations exceeding  $0.3 \mu\text{g}\cdot\text{g}^{-1}$ , the level reported to cause reproductive impairment in loons (Barr 1986). As reported by Evers et al. (1998) in their Wisconsin study, Scheuhammer et al. (1998) found that adults had blood mercury concentrations more than ten times higher than their chicks, and males had higher body weights and THg concentrations than females. The highest blood mercury concentrations ( $>2.5 \mu\text{g}\cdot\text{mL}^{-1}$ ) were found on lakes with pH below 6.5. Adult feathers contained 7.65 to  $21 \mu\text{g}\cdot\text{g}^{-1}$  THg, and chick feathers contained  $1.37$  to  $3.35 \mu\text{g}\cdot\text{g}^{-1}$  THg.

Total mercury was examined in a number of other bird species. Bald eagle (*Haliaeetus leucocephalus*) eggs collected from 1990 to 1992 from the lower mainland of British Columbia and Vancouver Island were found to contain from  $0.096$  to  $0.38 \mu\text{g}\cdot\text{g}^{-1}$  THg (Elliott et al. 1996). Nine species of marine birds collected from 1978 to 1984 from the Quoddy region (Bay of Fundy) contained  $0.037$  to  $7.05 \mu\text{g}\cdot\text{g}^{-1}$  THg (Braune 1987b). The highest concentrations were found in the kidney and liver of double-crested cormorants (*Phalacrocorax auritus*). The brain and muscle of this species had levels of  $<1.0 \mu\text{g}\cdot\text{g}^{-1}$  THg, as did all tissues of most other species (Braune 1987b).

Relatively high overall THg levels were reported from a number of birds from the Great Whale area of northern Québec. Common mergansers contained an average of  $17.5 \mu\text{g}\cdot\text{g}^{-1}$  THg in liver tissue, and  $1.3 \mu\text{g}\cdot\text{g}^{-1}$  in muscle tissue. Herring gulls (*Larus argentatus*) contained  $2.9 \mu\text{g}\cdot\text{g}^{-1}$  in liver tissue, and  $1.0 \mu\text{g}\cdot\text{g}^{-1}$  in muscle tissue. Common mergansers from the Nottaway-Broadback-Rupert area (SE coast of James Bay) had elevated tissue concentrations similar to mergansers from Great Whale (Langlois et al. 1995).

Total mercury levels were significantly higher in feathers of adult and nestling ospreys from reservoirs than natural environments in northern Québec. Adult osprey feathers from reservoirs averaged  $51.8 \text{mg}\cdot\text{kg}^{-1}$  dw, and from natural lakes and rivers averaged  $16.5 \text{mg}\cdot\text{kg}^{-1}$  dw (DesGranges et al. 1998). Nestling feathers contained 37.4 and  $7.0 \text{mg}\cdot\text{kg}^{-1}$  dw, respectively, from these locations. Osprey eggs contained  $0.2 \pm 0.1 \text{mg THg}\cdot\text{kg}^{-1}$  wet weight regardless of their location which was explained by the fact that eggs were laid prior to ice-out (therefore prior to high mercury exposure) on the water bodies under study in northern Québec.

A number of researchers have examined aquatic bird eggs in the vicinity of the Great Lakes. Concentrations in herring gull (*Larus argentatus*) eggs from several areas in the Great Lakes ranged from  $0.13$  to  $0.88 \mu\text{g}\cdot\text{g}^{-1}$  THg (Koster et al. 1996). The highest levels reported were for common tern (*Sterna hirundo*) eggs from Lake Ontario taken in 1971, of  $1.1 \mu\text{g}\cdot\text{g}^{-1}$  (Gilbertson 1974), and from Lake St. Clair in 1973, of  $1.1 \mu\text{g}\cdot\text{g}^{-1}$  (Stendell et al. 1976). Tree swallow (*Tachycineta bicolor*) eggs from Akwesasne (north-east of Cornwall), Long Point Tip



(Lake Ontario), Mud Creek (Lake Erie) and Wye Marsh (Georgian Bay) measured in 1991 had very low concentrations of 0.043 to 0.076  $\mu\text{g}\cdot\text{g}^{-1}$  (Bishop et al. 1995).

### 7.8 Non-biological media (App. B-8)

Most of the available information about MeHg in non-biological media in Canada has been published relatively recently because analytical techniques that permit detection of low levels were developed only in the late 1980s (Bloom 1989). Very few laboratories in the country are yet able to make MeHg measurements in water samples. Methylmercury measurements from Ontario and/or Québec have been reported for surface water, rain, snow, seston, sediment, peat porewater and flooded soils.

Information on THg in sediments in the Canadian environment is detailed in the report "Canadian Sediment Quality Guidelines for Mercury" (Environment Canada 1997), while the natural distribution of geological mercury as it relates to sediments in Canada is mapped and described in Friske and Coker (1995). For soil THg information, refer to Environment Canada (1996). Total mercury concentrations in non-biological media are not tabulated in the present report, except where they are supplemental to MeHg concentrations.

Precipitation in a remote area of north-western Ontario contained measurable concentrations of MeHg. Samples of rainwater taken in the early 1990s at the Experimental Lakes Area ranged from 0.017 to 0.049  $\text{ng}\cdot\text{L}^{-1}$  MeHg, or less than 1% of THg. In snow samples, MeHg concentrations ranged from 0.008 to 0.104  $\text{ng}\cdot\text{L}^{-1}$ , representing 0.8% to 2.9% of THg (St. Louis et al. 1995).

Water samples from a variety of surface waters at the Experimental Lakes Area had average MeHg concentrations ranging from 0.03 to 1.38  $\text{ng}\cdot\text{L}^{-1}$ . The higher concentration was from a pond after experimental flooding as part of an experimental reservoir project (Kelly et al. 1995). The lower concentration (0.03  $\text{ng}\cdot\text{L}^{-1}$ ) was for water flowing from an upland area that lacked wetlands, while water from watersheds dominated by wetlands had average levels of  $0.626 \pm 0.126$   $\text{ng}\cdot\text{L}^{-1}$  MeHg (St. Louis et al. 1994). Porewater from peat in wetlands contained 0.6  $\text{ng}\cdot\text{L}^{-1}$  MeHg, which was 10% of THg (Moore et al. 1995).

Water samples taken in the late 1970s from the mercury-contaminated Wabigoon/English River area near Dryden reportedly contained 0.22 to 1.3  $\text{ng}\cdot\text{L}^{-1}$  MeHg (Parks et al. 1989). While these values seem quite possible considering the high levels of mercury found in biota in the area, rigorous clean techniques for sampling and modern analytical methods were not developed at that time.

In the Ottawa and Gatineau Rivers, as much as 36% of THg in water samples was MeHg. Samples from the Ottawa River contained 2.3  $\text{ng}\cdot\text{L}^{-1}$  MeHg (26%), and from the Gatineau River, 4.1  $\text{ng}\cdot\text{L}^{-1}$  MeHg (36%). In the same study, water from Black Lake contained a lower concentration of MeHg (1.3  $\text{ng}\cdot\text{L}^{-1}$ ), but the proportion of MeHg remained high, at 37% (Schintu et al. 1989).

In the Great Whale and Nottaway-Broadback-Rupert areas of northern Québec, water/seston samples taken in the summer of 1989-1991 contained 0.45 and 0.94  $\text{ng}\cdot\text{L}^{-1}$  MeHg, respectively. Samples taken in the fall/winter from these areas contained 0.02 and 0.46  $\text{ng}\cdot\text{L}^{-1}$  MeHg, respectively. The MeHg levels were 9 to 31% of THg (Langlois et al. 1995). Methylmercury levels in water samples from natural lakes of northern Québec from 1994 to 1996 averaged 0.049 ( $\pm 0.004$  SE)  $\text{ng}\cdot\text{L}^{-1}$  ranging from 0.018 to 0.115  $\text{ng}\cdot\text{L}^{-1}$  (N=30; Lucotte et al. 1998).

## Canadian Tissue Residue Guidelines: Methylmercury

Methylmercury levels were approximately 3% of THg.

In the flooded systems of the Robert Bourassa and Laforge-1 Reservoirs of northern Québec, MeHg levels in water samples averaged  $0.28 \text{ ng}\cdot\text{L}^{-1}$  ranging from  $0.03$  to  $0.85 \text{ ng}\cdot\text{L}^{-1}$  (N=87; 1994-1996; Thérien and Morrison 1998). Over the same period, MeHg in water from neighbouring natural lakes was lower averaging  $0.05 \text{ ng}\cdot\text{L}^{-1}$  and ranging from  $0.02$  to  $0.11 \text{ ng}\cdot\text{L}^{-1}$  (N=30). The percent of THg that was MeHg averaged 12% in the reservoirs and 3% in the lakes.

Flooded soils in the vicinity of the La Grande-2 Reservoir of northern Québec contained up to  $0.043 \text{ }\mu\text{g}\cdot\text{g}^{-1}$  dw MeHg depending on the depth the soil sample was taken. Soils at 4-11 cm in depth had the highest concentration of  $0.043 \text{ }\mu\text{g}\cdot\text{g}^{-1}$  MeHg (30%), while the deeper soils from 11-19 cm contained  $0.0019 \text{ }\mu\text{g}\cdot\text{g}^{-1}$  MeHg (6%), the lowest amount. The surface soils from 0-4 cm contained an intermediate amount at  $0.0085 \text{ }\mu\text{g}\cdot\text{g}^{-1}$  MeHg (6.6%; Mucci et al. 1995).

Surface sediments from a number of northern Québec lakes contained from  $0.0001$  to  $0.0006 \text{ }\mu\text{g}\cdot\text{g}^{-1}$  dw MeHg (Mucci et al. 1995).

## **8.0 TOXICITY OF METHYLMERCURY TO PISCIVOROUS ORGANISMS**

While this chapter is intended to describe the toxicity of MeHg to wildlife consumers of fish, because the mechanism of toxicity is similar to other mammals, and because of the abundance of assays using laboratory animals, numerous examples using laboratory animals are also described. Avian wildlife species exposed to excessive MeHg usually incur reproductive and behavioural deficits, while the mammalian piscivores most often suffer neurological effects.

The toxicity of a substance to individual organisms depends on the level of exposure. Assuming that piscivorous birds and mammals consume fish from the same lake, the estimates of daily consumption rates, the trophic level of the fish consumed, and the body weight of the animal all contribute significantly to methylmercury exposure when expressed on a per kg of body weight basis. According to the CCME (1998), the daily food consumption of the otter is approximately 10% of body weight and that of mink is 24%. Assuming that 100% of the otter's diet is from aquatic food sources; piscivorous fish are assumed to comprise roughly 20% of this, while the other 80% consists of lower trophic level fish. On the other hand, mink are assumed to eat exclusively lower trophic fish. As a result of this fact and the high percent of daily bodyweight consumed as fish, otters will have a higher MeHg contact rate than mink (US EPA 1997b). The same reasoning and calculations may be used for piscivorous birds, which with few exceptions (e.g., herons), eat only lower trophic level fish.

The protective effect of selenium against methylmercury toxicity to vertebrates has been known for many years (Ganter et al. 1972). Although efforts to determine the exact mechanism continue, selenium has been shown to bind mercury after demethylation of methylmercury in the liver. The compounds formed in this manner probably include both mercury-selenoproteins and HgSe (Cavalli and Cardellicchio 1995; Palmisano et al. 1995). Because selenium restores the activities of certain liver enzymes, and glutathione status involved in antioxidative defence mechanisms, there may be a biologically derived protection against MeHg toxicity (Hoffman and Heinz 1998).

The consistent finding of lower percent MeHg in bird and mammal livers may be because many vertebrate species possess a capability to detoxify and sequester mercury originating as methylmercury in the diet. The extent to which this capability is developed appears to be related to feeding habits and is most well developed in fish-eating marine mammals and the carnivorous polar bear (Dietz et al. 1990). A correlation between selenium and mercury have also been reported for several seabirds, although the Se-Hg ratio may be higher than the usual 1:1 (Elliott et al. 1992). The capacity and adaptability of this system to detoxify methylmercury is largely unknown. Variable detoxification among individuals of a single species (pilot whales) has been demonstrated; lactating females demonstrated a significantly diminished detoxifying capability (Caurant et al. 1996).

### **8.1 Mode of Toxicity**

Methylmercury is a strong neurotoxicant for most vertebrates because they lack external barriers and substantial internal detoxifying systems. The central nervous system is the most sensitive tissue to MeHg exposure. Methylmercury is more toxic than other forms of mercury as it can pass through the blood-brain barrier and nuclear membranes to react directly with cellular components (Sloss 1995). As described above, the toxic effects of MeHg have been known to decrease with concurrent selenium exposure. The potential effect of selenium in reducing mercury toxicity has been reviewed by Cuvin-Aralar and Furness (1991).

With respect to MeHg uptake by animals, about 95% of an ingested dose of MeHg is absorbed into the bloodstream from the gastrointestinal tract, whereas inorganic Hg is less efficiently absorbed (7 to 15%; Wolfe et al. 1998). Methylmercury distributes to all parts of the body, readily crosses placental barriers, may enter the foetal brain, and is readily accumulated in growing animal fur, bird feathers or human hair. In humans, distribution to tissues may be complete in about two days, except for the brain, which takes about three days (Suzuki et al. 1991). The accumulation of mercury in the brain is greater in mammals than in fish (Zillioux et al. 1993).

Studies on the mechanism of MeHg passage across the blood-brain barrier have indicated that MeHg enters the capillary endothelial cell as a complex with the amino acid, cysteine. Attached to the thiol group of cysteine, MeHg enters the cell on the large neutral amino acid carrier. Given the presence of the MeHg-cysteine complex in plasma, it is thought that this may be the most common entry pathway into all mammalian cells (Suzuki et al. 1991). There even is some evidence that MeHg can cross the cell membranes of the blood-brain barrier by forming a compound that structurally mimics amino acids (Clarkson 1994). Researchers are not sure why brain cells are selectively damaged by MeHg.

Behavioural, physiological, histopathological, and biochemical demonstrations of toxicological effects of MeHg have been found for a variety of animals (Suzuki et al. 1991). Disruption of protein synthesis is one of the earliest biochemical effects (Clarkson 1994). An early symptom of MeHg poisoning in humans is paraesthesia (numbness and tingling) in the hands, feet and around the mouth (Sloss 1995), ataxia (loss of coordination), and gait or limb reflex dysfunction (Moore and Ramamoorthy 1984). Genotoxic (De Flora et al. 1994), developmental and reproductive effects (WHO 1989) have also been reported. Because MeHg can cross placental barriers, teratogenic and embryotoxic effects have been found. A list of the clinical signs and symptoms demonstrated by victims of the "Minimata Bay disease" is found in Table 8.

**Table 8. Frequency of clinical signs and symptoms of the Minimata Bay disease where humans consumed fish and shellfish containing 10-35 µg·g<sup>-1</sup> mercury for an extended period.**

| Symptom or Sign               | Frequency of Occurrence (%) |
|-------------------------------|-----------------------------|
| Constriction of visual fields | 100                         |
| Sensory disturbance           | 100                         |
| Loss of coordination          | 94                          |
| Impairment of speech          | 88                          |
| Impairment of hearing         | 85                          |
| Impairment of gait            | 82                          |
| Tremor                        | 76                          |
| Mental disturbance            | 71                          |
| Exaggerated tendon reflexes   | 38                          |
| Hypersalivation               | 24                          |
| Excessive sweating            | 24                          |
| Muscular rigidity             | 21                          |
| Chorea <sup>a</sup>           | 15                          |
| Pathologic reflexes           | 12                          |
| Athetosis <sup>b</sup>        | 9                           |
| Contractures <sup>c</sup>     | 9                           |

Source: Chang 1979

Notes:

<sup>a</sup> Uncontrollable and irregular muscle movements, especially of the arms, legs, and face.

<sup>b</sup> Inability to maintain one's position of fingers and toes.

<sup>c</sup> An abnormal, often permanent shortening, as of muscle or scar tissue that results in distortion or deformity, especially of a joint of the body.

## 8.2 Toxicity to Mammals

The few existing toxicity studies on wildlife species [mink (*Mustela vison*) and otter (*Lutra canadensis*)] are supplemented with assays utilizing laboratory animals. Most of the studies on wildlife species give MeHg concentrations in food without reporting quantities fed per unit body weight. Estimates of methylmercury consumption on a body weight (bw) per day basis are made preferably by using food intake (FI) and body weight data provided in the original study. Otherwise average daily consumption for an average sized animal species may be used (Table 9). For captive female mink, the average FI:bw ratio is 0.16 kg food·kg<sup>-1</sup> bw per day (Bleavins and Aulerich 1981). The lowest-observed-adverse-effect-level (LOAEL) refers to lowest MeHg concentration tested at which ecologically relevant adverse effects are observed. The no-observable-adverse effect-level (NOAEL) refers to the highest concentration tested at which ecologically relevant adverse effects are not observed. Any tissue concentrations measured at the termination of experiments are reported in Appendix C-1.

### 8.2.1 Acute and Chronic Toxicity (Appendix C-1)

Wren et al. (1987a) found that a diet containing 1 mg·kg<sup>-1</sup> MeHg was lethal to female and male mink within 11 weeks. As such, mink were fed the treatment diet every other day for the

remainder of the 6 month study (~15 weeks); no further mortalities or clinical effects occurred (C. Wren 1999, pers. com.). The researchers believed that these animals might have been additionally stressed by cold weather, causing higher sensitivity than might otherwise have been found. This conclusion is supported by experiments with two strains of mice that were more sensitive to MeHg toxicity at 8°C than at 22 or 38°C (Nomiya et al. 1980). However, cold weather stress is realistic for the Canadian environment, making this study relevant. Higher mortality occurred among female than male mink presumably due to higher rates of food consumption per body weight for females (0.18 kg food·kg<sup>-1</sup> bw per day) compared with males (0.10 kg food·kg<sup>-1</sup> bw per day) for males (Wren et al. 1987a). The authors reported daily intakes of 180 and 100 µg MeHg·kg<sup>-1</sup> bw per day for females and males, respectively. The reduced exposure (1 mg·kg<sup>-1</sup> MeHg every other day) equates daily intake rates of 90 and 50 µg MeHg·kg<sup>-1</sup> bw per day for females and males, respectively. The only sublethal effect reported once the exposure was reduced was that kits were born with a high mercury burden that decreased by five weeks of age. This was evidence of the importance of placental transfer but not transfer of MeHg via mother's milk (Wren et al. 1987a,b).

According to a recent study, fish from James Bay hydroelectric reservoirs may contain levels of MeHg that are toxic to mink. Three groups of female mink were fed daily with natural fish diets containing approximately 0.1, 0.5, and 0.9 mg MeHg·kg<sup>-1</sup> (16, 80, 144 µg·kg<sup>-1</sup> bw per day). The fish were both piscivorous and non-piscivorous species from the northern Québec reservoirs. The study had been originally designed to run for a two year period, however, over 50% of the mink in the 0.9 mg·kg<sup>-1</sup> group died with signs of neurotoxicity between days 80 and 101 (11 to 14 weeks) of the study (Chamberland et al. 1996). This is the same length of exposure that lethal effects were noted for mink fed a similar dosage by Wren et al. (1987a) described above. No clinical signs, neurological damage or reproductive effects were found for the lower dosages (0.1, 0.5 mg MeHg·kg<sup>-1</sup>; Laperle et al. 1998). Brown-coloured urine was observed after day 59 in some mink fed the intermediate and highest dosage, although the toxicological significance of this finding was unknown. The total urine protein and individual protein data corroborated the finding of Wobeser et al. (1976b; discussed below) of little or no renal damage at all dosage levels (Chamberland et al. 1996). The authors reported no confounding effects of other contaminants in the fish diet.

In a similar study during the 1970s, female mink were fed rations that were a mix of ranch mink feed and fish (freshwater drum from Lake Winnipeg), at concentrations of <0.1 mg·kg<sup>-1</sup> (control), 0.25 mg·kg<sup>-1</sup>, and 0.34 mg·kg<sup>-1</sup> (<16, 40, 54 µg·kg<sup>-1</sup> bw per day). After 145 days (21 weeks), MeHg was accumulated in the liver and kidney, but no clinical signs of MeHg toxicity or histopathological lesions were observed in any of the mink (Wobeser et al. 1976a).

Wobeser et al. (1976b) reported that female mink fed a diet containing 1.1 mg·kg<sup>-1</sup> (176 µg·kg<sup>-1</sup> bw per day) MeHg survived, although histopathological abnormalities including pale, yellow livers, lesions in the central nervous system, and axonal degeneration were evident. In addition, two of five mink showed reduced movement toward the end of the three-month (12 week) study, the researchers believed that more effects would have occurred had the exposure period been longer. Anorexia, ataxia and death resulted at the higher dosage of 1.8 mg MeHg·kg<sup>-1</sup> (288 µg·kg<sup>-1</sup> bw per day).

Female and male mink fed a very high concentration of 5 mg MeHg·kg<sup>-1</sup> survived an average of one month (Aulerich et al. 1974). Using body weight and food consumption data provided in the original paper, this dietary dose is converted to 700 and 500 µg MeHg·kg<sup>-1</sup> bw per day for female and male mink, respectively. The mink exhibited typical acute MeHg poisoning symptoms of ataxia, anorexia and limb paralysis prior to death. A diet containing 10 mg·kg<sup>-1</sup>

inorganic Hg (mercuric chloride) for five months produced no effects. As these were the only concentrations administered for the two forms of mercury, LC<sub>50</sub> and LOAEL values were not established.

The single acute toxicity study on otters, although somewhat limited in scope, suggested that otters are as sensitive as mink to MeHg poisoning in laboratory studies. An undescribed diet containing 2 mg·kg<sup>-1</sup> (90 µg·kg<sup>-1</sup> bw per day<sup>18</sup>) over 159 to 213 days (5.3 to 7.1 months) was lethal to two of three otters studied. In addition, at this dose, anorexia occurred in one, and ataxia in two of these otters (O'Connor and Nielsen 1981). The surviving otter displayed both toxic symptoms prior to its euthanasia at the end of the study. A diet of 4 mg·kg<sup>-1</sup> MeHg (170 µg·kg<sup>-1</sup> bw per day) caused anorexic and ataxic symptoms in all three otters over 101 to 123 days, after which the otters were euthanized. A diet containing 8 mg·kg<sup>-1</sup> MeHg (370 µg·kg<sup>-1</sup> bw per day) was lethal to one of three otters within 59 days, with toxic symptoms in all three otters prior to their euthanasia. Because of a high minimum treatment level, a chronic LOAEL and NOAEL (no-observed-adverse-effect-level) could not be determined. In the wild, however, otters may be more susceptible to MeHg poisoning than mink because approximately 20% of the otter diet may consist of higher trophic level fish and, therefore, higher MeHg concentrations (US EPA 1997a; Table 9).

No studies were found on sublethal effects to otters, although at least one study documented sublethal concentrations of mercury in captured otters based on the laboratory studies with mink and otter described above (Halbrook et al. 1994).

#### LABORATORY ANIMALS

A diet containing 0.25 mg·kg<sup>-1</sup> MeHg was highly toxic to domestic cats (*Felis catus*) when administered 5 days per week over 90 days. The cats first showed convulsions at day 68, with a mean survival time of 78 days (Eaton et al. 1980). Methylmercury was less toxic to cats when fed as contaminated tuna fish, where 0.50 mg·kg<sup>-1</sup> fed for up to 11 months caused biochemical changes (Chang et al. 1974), and 0.55 mg·kg<sup>-1</sup> plus 0.29 mg·kg<sup>-1</sup> Se for 188 days induced behavioural changes (Haupt et al. 1988). The latter two studies only indicated the MeHg concentration in food without describing the intake based on body weight.

Domestic cats fed a range of dosages of MeHg daily for up to 2 years showed neurological impairment and minor CNS pathology at the higher dosages. The cats were fed 0.06 mg kg<sup>-1</sup> bw diet per day. The diet was a mix of contaminated and uncontaminated northern pike (Charbonneau et al. 1976). Doses were 0.05 (control), 0.14, 0.33, 0.76, 1.2, and 2.9 mg THg·kg<sup>-1</sup> (3, 8.4, 19.8, 45.6, 73.8 and 176 µg THg·kg<sup>-1</sup> bw per day). Periodical monitoring of the diet indicated that MeHg accounted for 95% of THg. Clinical signs of MeHg toxicity (ataxia, loss of balance and motor incoordination) occurred in groups receiving 176 µg·kg<sup>-1</sup> bw per day after 14 weeks of treatment, and in groups receiving 74 µg·kg<sup>-1</sup> bw per day after 40 weeks of treatment. Subtle neurological deficits (impaired "hopping reaction" and hypalgesia<sup>19</sup>) after 60 weeks of treatment occurred but did not intensify in groups receiving 46 µg·kg<sup>-1</sup> bw per day (0.76 mg·kg<sup>-1</sup>). After two years, there were no overt clinical signs of toxicity at the two lowest dosages and the control (Charbonneau et al. 1976). These findings suggest that 20 µg·kg<sup>-1</sup> bw per day is the approximate NOAEL and 46 µg·kg<sup>-1</sup> bw per day is the LOAEL for chronic dietary exposure to methylmercury in domestic cats.

---

<sup>18</sup> Daily intake rates were calculated by the original authors using measured body weights and food consumption data.

<sup>19</sup> Diminished sensitivity to pain.

Crab-eating macaques (*Macaca fascicularis*) exhibited insensitivity to touch and loss of tactile response (Rice 1989) when dosed with MeHg. Five monkeys were given  $50 \mu\text{g}\cdot\text{kg}^{-1}$  bw per day from birth to seven years. Clinical and neurologic examinations were carried out during the exposure period and for an additional six years. As an indicator of the latent effects of MeHg, objective neurological examinations performed at the end of the observation period revealed that the treated monkeys were clumsier and slower to react when placed in exercise cages than control monkeys (Rice 1989). Dosages of up to  $50 \mu\text{g}\cdot\text{kg}^{-1}$  bw per day fed in apple juice to macaques over 150 days caused no clinical symptoms and no significant difference in cholinesterase activity (Petruccioli and Turillazzi 1991).

Groups of Wistar rats (50 per sex per dose) were administered nominal daily doses of 2, 10, 25, 50 and  $250 \mu\text{g}\cdot\text{kg}^{-1}$  bw per day of MeHg for 26 months (Munro et al. 1980). Female rats that received  $25 \mu\text{g}\cdot\text{kg}^{-1}$  bw per day had reduced body weight gains and showed only minimal clinical signs of neurotoxicity; however, male rats that received this dose showed overt clinical signs of neurotoxicity, had decreased hemoglobin and hematocrit values, reduced weight gains, and showed increased mortality. Histopathological examination of rats of both sexes receiving  $25 \mu\text{g}\cdot\text{kg}^{-1}$  bw per day revealed demyelination<sup>20</sup> of dorsal nerve roots and peripheral nerves. Males had severe renal (kidney) damage, while females had minimal renal damage. This study reported a NOAEL of  $5 \mu\text{g}\cdot\text{kg}^{-1}$  bw per day and a LOAEL of  $25 \mu\text{g}\cdot\text{kg}^{-1}$  bw per day.

A two year feeding study of MeHg chloride was conducted on B6C3F1 mice (60 mice per sex per group) at doses of 0, 0.4, 2, and  $10 \text{mg}\cdot\text{kg}^{-1}$  (0, 40, 170, and  $830 \mu\text{g}\cdot\text{kg}^{-1}$  bw per day) to determine chronic toxicity and possible carcinogenic effects (Mitsumori et al. 1990). The mice were examined clinically during the study and neurotoxic signs characterised by posterior paralysis were observed in 33 males after 59 weeks and three females after 80 weeks in the  $830 \mu\text{g}\cdot\text{kg}^{-1}$  bw per day group. A marked increase in mortality and a significant decrease in body weight gain were also observed in the  $830 \mu\text{g}\cdot\text{kg}^{-1}$  bw per day male dose group, beginning at 60 weeks. Post mortem examination revealed toxic encephalopathy<sup>21</sup> consisting of neuronal necrosis of the brain and toxic peripheral sensory neuropathy<sup>22</sup> in both sexes of the high dose group. An increased incidence of chronic nephropathy<sup>23</sup> was observed in the 170 and  $830 \mu\text{g}\cdot\text{kg}^{-1}$  bw per day males. Based upon this study a NOAEL of  $40 \mu\text{g}\cdot\text{kg}^{-1}$  bw per day and a LOAEL of  $170 \mu\text{g}\cdot\text{kg}^{-1}$  bw per day was determined.

Mitsumori et al. (1983, 1984) administered diets containing 0, 0.4, 2 or  $10 \text{mg}\cdot\text{kg}^{-1}$  of methylmercuric chloride (0, 0.011, 0.05, and  $0.28 \text{mg}\cdot\text{kg}^{-1}$  bw per day in males; 0, 0.014, 0.064, and  $0.34 \text{mg}\cdot\text{kg}^{-1}$  bw per day in females) to 56 per sex per group Sprague-Dawley rats for up to 130 weeks. At week 104, survival was approximately 55, 45, 75, and 10% in control, low-, mid- and high-dose males, respectively, and 70, 75, 75 and 30% in control, low-, mid- and high-dose females, respectively. All males in the high-dose group had died by week 119. Body weight gain was significantly decreased in high-dose males and females after 44 weeks (approximately 10-20%).

Groups of 25 female and 25 male rats were administered methylmercuric chloride at dietary levels of 0, 0.1, 0.5 and  $2.5 \text{mg}\cdot\text{kg}^{-1}$  (0, 0.004, 0.02, and  $0.1 \text{mg}\cdot\text{kg}^{-1}$  bw per day) for two years. No significant effects were observed on growth or food intake except for a 6% decrease in body

---

<sup>20</sup> To destroy or remove the myelin sheath of (a nerve fiber), as through disease.

<sup>21</sup> Any of various diseases of the brain.

<sup>22</sup> A disease or an abnormality of the nervous system.

<sup>23</sup> A disease or an abnormality of the kidney (may be referred to as renal damage).



weight gain at 60 weeks in high-dose females. Survival was 72, 68, 48 and 48% in control, low-, mid- and high-dose males, respectively, and 76, 60, 64 and 56% in control, low-, mid- and high-dose females, respectively (statistical significance not reported). Increases in relative kidney weights were observed in both males and females at the highest dose (Verschuuren et al. 1976).

Chronic exposure to MeHg at levels too low to cause clinical signs of toxicity may cause deficiencies in the immune system of animals. The long-term bioaccumulation of MeHg may cause animals to be more susceptible to diseases that they might otherwise resist (Wolfe et al. 1998).

### **8.2.2 Reproductive and Developmental Toxicity (Appendix C-2)**

Reproductive effects of MeHg in mammals include developmental effects that produce behavioural deficits in offspring, impaired fertility, and foetal death. Swimming ability, operant learning, avoidance, maze learning, and development of reflexes may be affected at low dosages. Effects that occur at successively higher dosages include changes in spontaneous activity, visual function, vocalisation, and convulsions (Chang and Annau 1984; Shimai and Satoh 1985; Eccles and Annau 1987). These effects occur because MeHg selectively targets the foetal brain. Females may also transfer MeHg to their young in milk, although this was previously dismissed as being unimportant in mink (Wren 1987). Foetal red blood cells may contain 30% more MeHg than maternal blood cells (c.f. Wolfe et al. 1998).

Gunderson et al. (1986) administered daily doses of 50-70  $\mu\text{g}\cdot\text{kg}^{-1}$  bw per day of MeHg to eleven crab-eating macaques throughout pregnancy which resulted in blood levels of 1080-1330  $\mu\text{g}\cdot\text{L}^{-1}$  in mothers and 1410-1840  $\mu\text{g}\cdot\text{L}^{-1}$  in offspring. When tested 35 days after birth the infants exhibited significant visual recognition deficits compared to control macaque offspring.

In another study, groups of seven or eight female crab-eating macaques were given 0, 50 or 90  $\mu\text{g}\cdot\text{kg}^{-1}$  bw per day of MeHg through four menstrual cycles (Burbacher et al. 1984). They were mated with untreated males and clinical observations were made for an additional four months. Two of seven 90  $\mu\text{g}\cdot\text{kg}^{-1}$  bw per day treated females aborted and three did not conceive during the four-month mating period; the other two females delivered live infants. Two of seven females of the 50  $\mu\text{g}\cdot\text{kg}^{-1}$  bw per day dose group aborted; the remaining females delivered live infants. All eight females of the control group conceived and six delivered live infants. These reproductive results were not statistically significant. Reproductive failure within dose groups could be predicted by blood Hg levels. The dams did not show clinical signs of MeHg poisoning during the breeding period or gestation but dosed with 90  $\mu\text{g}\cdot\text{kg}^{-1}$  bw per day for one year, four of seven showed adverse neurological signs.

Bornhausen et al. (1980) reported a decrease in operant behaviour performance in four-month-old Wistar rats whose dams had received from 5 to 50  $\mu\text{g}\cdot\text{kg}^{-1}$  bw per day of MeHg on days 6 through 9 of gestation. A statistically significant effect ( $P<0.05$ ) was observed in offspring whose dams had received 10 and 50  $\mu\text{g}\cdot\text{kg}^{-1}$  bw per day during gestation. A LOAEL that could be consumed by pregnant female rats causing decreases in offspring operant behaviour of 10  $\mu\text{g}\cdot\text{kg}^{-1}$  bw was suggested.

### **8.2.3 Carcinogenic and Mutagenic Effects (Appendix C-3)**

Very few studies implicate MeHg as a carcinogen. Based on inadequate data in humans and

limited evidence of carcinogenicity in animals, the US EPA has classified MeHg as a *possible* human carcinogen (US EPA 1995b). Male mice exposed to methylmercuric chloride in the diet had an increased incidence of renal adenomas<sup>24</sup>, adenocarcinomas<sup>25</sup>, and carcinomas. Several negative cancer bioassays were also reported. Although genotoxicity test data suggest that MeHg is capable of producing chromosomal and nuclear damage, there are also abundant negative genotoxicity data (De Flora et al. 1994).

The results are contradictory as to whether MeHg is carcinogenic to rodents. Interpretation of two of the positive studies on mice was complicated by observation of tumours only at doses that exceeded the maximum tolerated dose (MTD)<sup>26</sup>. A fourth dietary study in mice and four separate dietary studies in rats failed to indicate carcinogenicity associated with MeHg exposure.

Methylmercuric chloride was administered in the diet of ICR mice at levels of 0, 0.4, 2 or 10 mg·kg<sup>-1</sup> (0, 0.0375, 0.188, and 0.906 mg·kg<sup>-1</sup> bw per day in males and 0, 0.0318, 0.114, and 0.785 mg·kg<sup>-1</sup> bw per day in females). Sixty mice of each sex were fed the given dose level for 104 weeks (Hirano et al. 1986). Six mice per sex in each group were sacrificed at 26, 52 and 78 weeks. Complete histopathological examinations were performed on all animals found dead, killed *in extremis* or killed by design. Mortality, group mean body weights and food consumption were comparable to controls. The first renal tumour was observed at 58 weeks in a high-dose (0.906 mg·kg<sup>-1</sup> bw) male, and the incidence of renal epithelial tumours (adenomas or adenocarcinomas) was significantly increased in high-dose males (1/32, 0/25, 0/29, and 13/26 in the control, low-, mid- and high-dose males, respectively). Focal hyperplasia of the tubular epithelium was reported to be increased in high-dose males (13/59; other incidences not reported). Epithelial degeneration of the renal proximal tubules and degeneration or fibrosis of the sciatic nerve was reported in high-dose females.

Mitsumori et al. (1981) administered 0, 15 or 30 mg·kg<sup>-1</sup> of methylmercuric chloride in the diet (0, 1.6 and 3.1 mg·kg<sup>-1</sup> bw per day) to 60 ICR mice per sex per group for 78 weeks. Interim sacrifices of up to 6 per sex per group were conducted at weeks 26 and 52. Survival was decreased in a dose-related manner; at week 78 survival was 40, 10 and 0% in control, low- and high-dose males, respectively, and 55, 30 and 0%, in control, low- and high-dose females, respectively (statistical analyses not performed). The majority of high-dose mice (85% males and 98% females) died by week 26 of the study. Examination of the kidneys of mice that died or were sacrificed after 53 weeks showed a significant increase in renal tumours in low-dose males (13/16 versus 1/37 in controls). Renal tumours were found in males but not in females of any group. Evidence of neurotoxicity and renal pathology were observed in the treated mice at both dose levels.

Mitsumori et al. (1983, 1984) administered diets containing 0, 0.4, 2 or 10 mg·kg<sup>-1</sup> of methylmercuric chloride (0, 0.011, 0.05, and 0.28 mg·kg<sup>-1</sup> bw per day in males; 0, 0.014, 0.064, and 0.34 mg·kg<sup>-1</sup> bw per day in females) to 56 per sex per group of Sprague-Dawley rats for up to 130 weeks. No increase in tumour incidence was observed in either males or females. Intermediate dose (0.05 mg·kg<sup>-1</sup> bw per day) males exhibited significantly increased degeneration of the proximal tubular epithelium and hyperplasia of the parathyroid.

---

<sup>24</sup> A benign epithelial tumour having a glandular origin and structure.

<sup>25</sup> A malignant tumour originating in glandular tissue.

<sup>26</sup> The dose that an animal species can tolerate for a major portion of its lifetime without significant impairment or toxic effects other than carcinogenicity (US EPA website: [www.epa.gov](http://www.epa.gov)).

No increase in tumour incidence or decrease in tumour latency was observed in another study using rats of an unspecified species (Verschuuren et al. 1976); groups of 25 female and 25 male rats were administered methylmercuric chloride at dietary levels of 0, 0.1, 0.5 and 2.5 mg·kg<sup>-1</sup> (0, 0.004, 0.02, and 0.1 mg·kg<sup>-1</sup> bw per day) for two years. No effects on the nature or incidence of pathological lesions were observed, and tumours were reported to have been observed with comparable incidence and latency among all of the groups.

No increase in tumour incidence was observed in a multiple-generation reproduction study using Sprague-Dawley rats (Newberne et al. 1972). Groups of rats (30 per sex) were given semi-synthetic diets supplemented with either casein or a fish protein concentrate to yield dietary levels of 0.2 mg·kg<sup>-1</sup> MeHg (0.008 mg·kg<sup>-1</sup> bw per day). Another group of controls received untreated rat chow. Rats that received diets containing MeHg during the two-year study had body weights and haematology comparable to controls. Detailed histopathological analyses revealed no lesions of the brain, liver, or kidney that were attributable to MeHg exposure. Mortality data were not presented.

As reviewed in WHO (1990), MeHg is not a potent mutagen but appears to be capable of causing chromosome damage and nuclear perturbations under some circumstances. For the purposes of this report, MeHg will be considered non-mutagenic at levels found in Canadian environments, however, a few laboratory studies will be described.

MeHg produced chromosomal aberrations and aneuploidy<sup>27</sup> in human peripheral lymphocytes (Betti et al. 1992), sister chromatid exchange in human lymphocytes (Morimoto et al. 1982), and DNA damage in human nerve and lung cells as well as Chinese hamster V-79 cells and rat glioblastoma cells (Costa et al. 1991).

Bone marrow cells of cats treated with MeHg in a study by Charbonneau et al. (1976) were examined for mutagenic effects by Miller et al. (1979). The MeHg treatment caused an increased number of nuclear abnormalities and an inhibition of DNA repair capacity. Methylmercury induced a weak mutagenic response in Chinese hamster V-79 cells (Fiskesjo 1979). Methylmercury also induced histone protein perturbations and influenced factors regulating nucleolus-organising activity (WHO 1990). Moreover, MeHg has been reported to interfere with gene expression in cultures of glioma cells (WHO 1990).

Negative mutagenic findings have also been reported. Methylmercury acetate was reported to be negative in a mouse micronucleus assay (Heddle and Bruce 1977). Matsumoto and Spindle (1982) reported no significant increase in sister chromatid exchange in developing mouse embryos. They did report, however, that the developing mouse embryos were highly sensitive to *in vitro* treatment with MeHg.

### 8.3 Toxicity to Avian Species

As it does in mammals, MeHg readily penetrates the blood-brain barrier in birds and may cause brain lesions, spinal cord degeneration, and CNS dysfunctions. It is a potent neurotoxicant and embryo toxicant. Also similar to mammals, intestinal absorption of MeHg in birds is nearly 100%, while that for inorganic Hg is limited to a few percent (Scheuhammer 1987). The health concerns for the avian wildlife species are usually reproductive and behavioural deficits; mammalian quadrupeds most often suffer ataxia, anorexia and hindlimb paralysis.

---

<sup>27</sup> Having a chromosome number that is not a multiple of the haploid number for the species.

Most of the studies on wildlife species give MeHg concentrations in food without reporting quantities fed per unit body weight. Estimates of methylmercury consumption on a body weight (bw) per day basis are made preferably by using food intake (FI) and body weight data provided in the original study. Otherwise average daily consumption for an average sized animal species may be used (Table 9). The lowest-observed-adverse-effect-level (LOAEL) refers to lowest MeHg concentration tested at which ecologically relevant adverse effects are observed. The no-observable-adverse effect-level (NOAEL) refers to the highest concentration tested at which ecologically relevant adverse effects are not observed. Any tissue concentrations measured at the termination of experiments are reported in Appendix C-4.

### **8.3.1 Acute and Chronic Toxicity**

MeHg poisoning in birds may cause loss of appetite, leading to weight loss, progressive weakness in wings and legs, and difficulty flying, walking or standing (Scheuhammer 1987; Wolfe et al. 1998). Kidney disease and kidney lesions are associated with elevated dietary mercury (Nicholson and Osborn 1984; Snelgrove-Hobson et al. 1988; Rao et al. 1989). Brain residues are most diagnostic for acute mercury poisoning in birds (Wolfe et al. 1998), while blood concentrations have been related to chronic effects (Evers et al. 1998). Total mercury concentrations in liver and kidney tissues of piscivorous birds are insufficient criteria for making toxicologically relevant judgements of possible mercury toxicity because such a small proportion is MeHg, and inorganic Hg may be excreted from these tissues (Scheuhammer et al. 1998).

A recent study indicated contrasting protective effects of selenium on MeHg toxicity to adults compared with young mallards. Selenomethionine protected against MeHg poisoning in adult males, but it worsened the effects of MeHg on the hatching, survival, and growth of offspring, and produced more teratogenic effects (Heinz and Hoffman 1998). Except for one control that died of unknown causes, no adults that were fed a diet containing  $10 \text{ mg}\cdot\text{kg}^{-1}$  MeHg plus  $10 \text{ mg}\cdot\text{kg}^{-1}$  selenomethionine exhibited any overt signs of toxicity. The physiological protection of Se against MeHg poisoning was found to include restoration of plasma and liver GSH-peroxidase activity, partial restoration of liver and brain G-6-PDH activity, and lower hepatic oxidized glutathione (GSSG) concentration (Hoffman and Heinz 1998).

Mallards fed a diet containing  $10 \text{ mg}\cdot\text{kg}^{-1}$  MeHg chloride showed overt signs of toxicity starting at about seven weeks. Males had weakened legs, and by day 73, one of 12 males died and eight others could barely walk (Heinz and Hoffman 1998). The researchers hypothesised that adult females did not suffer from MeHg poisoning during the study because they excreted the MeHg during egg laying, reducing their mercury body burdens (Heinz and Hoffman 1998).

### **8.3.2 Reproductive and Developmental Toxicity**

The avian reproductive system is very sensitive to MeHg toxicity. Overall reproductive success in birds can decrease by 35 to 50% due to dietary MeHg exposures that would be insufficient to cause obvious signs of intoxication in adults (Wolfe et al. 1998).

In a field study of effects to loons (*Gavia immer*) over three years in the contaminated English-Wabigoon district of north-western Ontario, Barr (1986) reported differences in loon reproductive behaviour and success between contaminated and uncontaminated lakes. On Ball Lake, where mercury in perch (mean length 16.1 cm) averaged  $0.36 \text{ mg}\cdot\text{kg}^{-1}$ , no nests were initiated even though loons were present on territories. Based on an adult bird size of 4 kg consuming 800 g of fish per day, this is equivalent to consuming  $72 \text{ }\mu\text{g}\cdot\text{kg}^{-1}\text{bw}\cdot\text{d}^{-1}$  (US EPA 1997b). In other areas of the contaminated section, Barr (1986) suggested that mercury levels in loon eggs may

have been sufficiently high to reduce hatching success because they exceeded levels found to cause embryo mortality in other species. Reductions in egg laying, and in nest site and territorial fidelity were associated with mean mercury concentrations ranging from 0.3 to 0.4 mg·kg<sup>-1</sup> (60 to 80 µg·kg<sup>-1</sup> bw·d<sup>-1</sup>) in perch and crayfish, and from 2 to 3 mg·kg<sup>-1</sup> in adult loon brains and loon eggs. Loons established few territories, laid only one egg per pair, and raised no progeny in waters where mean contamination of small prey species exceeded 0.4 mg·kg<sup>-1</sup> (Barr 1986).

Some male loons from mercury-contaminated Ball Lake were emaciated, which is a symptom of heavy metal poisoning (Borg et al. 1970). Because males consume more than females and eat larger fish, and do not excrete contaminants through egg laying, male loons were considered more susceptible to MeHg toxicity. Barr (1986) concluded that as mercury contamination increases, adult loons were more likely to fail to produce eggs, and that contaminated prey species caused aberrant behaviour in adult loons that resulted in reduced reproductive success. Lakes where no effects to loon behaviour and reproduction were observed contained perch with average mercury concentrations ranging from 0.08 to 0.17 mg·kg<sup>-1</sup> (Barr 1986). The calculated NOAEL for 0.17 mg·kg<sup>-1</sup> is 34 µg·kg<sup>-1</sup> bw·d<sup>-1</sup> based on the daily consumption of 800 g of food by a 4 kg loon (CCME 1998; US EPA 1997b).

The number of adult loons returning to their breeding territories in Wisconsin was found not to be influenced by differences in mercury exposure. Annual return rates of 85 to 90% were recorded for three years, indicating a high rate of adult survival despite blood mercury concentrations of up to 4.2 µg·mL<sup>-1</sup>, and feather concentrations up to 21 µg·g<sup>-1</sup> ww (Evers et al. 1998). Fewer chicks hatched or survived to eight weeks of age on lakes where elevated mercury (>0.3 µg·mL<sup>-1</sup>) was measured in chick blood. While measurements were made as THg, it is assumed that MeHg was ingested because the diet of these loons was fish (fish mercury concentrations were not reported).

Great blue herons (*Ardea herodias*) and double-crested cormorants appear to be more resistant to MeHg toxicity than loons. The reproductive success rates of herons and cormorants at a mercury-contaminated site in California were comparable to success rates from sites that were not contaminated (Wolfe and Norman 1998). The nesting colonies were on Clear Lake, which formerly had open-pit cinnabar mining and ore processing activities on its shoreline. Mercury waste continues to leach into the lake. The mercury concentration in food (minnows) recovered in and around nests, and the gastrointestinal tracts of heron young averaged 0.56 mg·kg<sup>-1</sup> (range of 0.31-0.87 mg·kg<sup>-1</sup>), and the mercury concentration in heron young brain tissue averaged 0.35 mg·kg<sup>-1</sup>. Young cormorants had brain tissue concentrations of 0.63 (0.54 - 0.72 mg·kg<sup>-1</sup>). The dietary intake for heron nestlings was calculated to be 73 to 210 µg MeHg·kg<sup>-1</sup> bw·d<sup>-1</sup> (using a food consumption rate of 0.245 kg·d<sup>-1</sup> for a chick weighing 1.0 kg), with adults having an average rate of 122 µg MeHg·kg<sup>-1</sup> bw·d<sup>-1</sup> (CCME 1998). The number of eggs produced and the growth rates of young herons were no different than from colonies in uncontaminated sites from Washington and Canada (Wolfe and Norman 1998).

Feeding experiments with adult black ducks (*Anas rubripes*) caused a number of reproductive and developmental effects. Breeding pairs were fed 3 mg·kg<sup>-1</sup> MeHg free-choice over two breeding seasons. Clutch size, egg production, number of eggs incubated, hatchability, and survival of ducklings were lower during both years in hens fed MeHg. Whole embryos that failed to hatch contained means of 9.6 and 6.1 mg·kg<sup>-1</sup> mercury during the first and second years, respectively. Brains of dead ducklings were associated with lesions and contained from 3 to 7 mg·kg<sup>-1</sup> mercury (Finley and Stendell 1978). By the second year of study, some adults displayed some behavioural effects characteristic of MeHg poisoning, but otherwise were

healthy. Amounts of MeHg consumed was not reported either on a daily or total experimental basis.

Experiments that compared reproduction and behaviour over two reproductive seasons in mallard ducks fed diets contaminated with methylmercury indicated that effects were less severe in the second season than the first. Adult hens were fed through their first reproductive season diets containing nominal concentrations of 0 (control), 0.5 or 3.0 mg·kg<sup>-1</sup> dw of MeHg; the measured concentrations were <0.05, 0.55, and 3.4 mg·kg<sup>-1</sup> dw<sup>28</sup>, respectively. Egg production ceased after 43, 31, and 21 weeks exposure to 0, 0.5, and 3.0 mg·kg<sup>-1</sup> dw treatment diets, respectively. The only observed effect at the 0.5 mg·kg<sup>-1</sup> dw treatment level within the first 21 weeks of exposure was that the eggs from treated females were smaller than controls (Heinz 1974). Those fed the more highly contaminated diet (3.0 mg·kg<sup>-1</sup> dw) laid fewer eggs, had reduced hatching success and many of the young died within one week of hatching (results not quantified). By the second reproductive season, adult hens had been fed treatment diets continuously for 14 months. As in the first reproductive season, percent survival (89%) of young from hens fed 3.0 mg·kg<sup>-1</sup> dw (measured as 2.88 mg·kg<sup>-1</sup> dw<sup>29</sup>) was also lower than the survival of controls (98%), but the effect was less severe in the second season (Heinz 1976a). Ducklings from the treatment group that died had brain lesions characteristic of MeHg toxicity while those that survived were hyper-responsive in avoidance tests (Heinz 1975; 1976a). In contrast to the first reproductive season, egg production and hatching success for hens fed 3.0 mg·kg<sup>-1</sup> dw were not significantly different from controls in the second reproductive season (Heinz 1976a). There were no consistent differences between the control and 0.5 mg·kg<sup>-1</sup> dw (measured as 0.53 mg·kg<sup>-1</sup> dw<sup>30</sup>) treatments with regards to proportion of eggs laid outside the nest, eggshell thickness, hatching success, or duckling survival to one week during the second reproductive season. With respect to behavioural responses, effects appeared also less severe in the second season than in the first. During the first season, offspring from both treatment groups were hyper-responsive in avoidance tests while during the second season only offspring from the 3.0 mg·kg<sup>-1</sup> dw dose were hyper-responsive (Heinz 1975). Heinz (1976a) hypothesised that the combination of improved reproductive success and fewer behavioural effects for ducks in the 3.0 mg·kg<sup>-1</sup> dw treatment group during the second reproductive season was due in part to a reduction in mercury levels in eggs and in part to a greater ability of adults and young to deal with methylmercury poisoning.

The adult mallards and their second season young in the above study served also as the first and second generations, respectively, of a three generation study (Heinz et al. 1976a;b, 1979). Beginning at 9 days of age, offspring were fed the same nominal dose as their mothers (0 or 0.5 mg·kg<sup>-1</sup> dw; measured as 0.47 mg·kg<sup>-1</sup> dw<sup>31</sup>). At adulthood, second generation treated mallards did not differ from controls in body weight or egg laying rates even though they consumed significantly more food. Treated hens displayed aberrant nesting behaviour and their offspring displayed reduced growth and survival, but not impaired behaviour (Heinz 1976b; Heinz 1979). These results were reversed in the third generation; third generation adults did not consume significantly more food and their offspring had growth rates comparable to controls.

---

<sup>28</sup> For the first reproductive season, Heinz (1974) reports measured concentrations for the nominal dose 0.5 mg·kg<sup>-1</sup> dw of 0.49, 0.53, and 0.64 mg·kg<sup>-1</sup> dw; the mean of these three values is 0.55 mg·kg<sup>-1</sup> dw. Measured concentrations for 3.0 mg·kg<sup>-1</sup> dw were 3.32, 3.32, and 3.55 mg·kg<sup>-1</sup> dw; the mean of these three values is 3.4 mg·kg<sup>-1</sup> dw

<sup>29</sup> For the second reproductive season, the nominal dose of 3.0 mg·kg<sup>-1</sup> dw was measured as 2.22, 2.83, and 3.59 mg·kg<sup>-1</sup> dw; the mean is 2.88 mg·kg<sup>-1</sup> dw (Heinz 1976a).

<sup>30</sup> For the second reproductive season, the nominal dose of 0.5 mg·kg<sup>-1</sup> dw was measured as 0.52, 0.53, and 0.54 mg·kg<sup>-1</sup> dw; the mean is 0.53 mg·kg<sup>-1</sup> dw (Heinz 1976a). These concentrations apply also to the first generation of the three generation study.

<sup>31</sup> For the second and third generations, Heinz (1976b; 1979) reports the means of the measured concentrations as 0.47 and 0.43 mg·kg<sup>-1</sup> dw, respectively, for the nominal dose of 0.5 mg·kg<sup>-1</sup> dw. Results from second and third generation studies were not reported for a 3.0 mg·kg<sup>-1</sup> dw treatment level as in the earlier publications.

Third generation offspring, however, displayed impaired behaviour; fewer ducklings took more time to approach a tape-recorded maternal call (Heinz 1979).

The effect of MeHg on mallard development was studied by applying quantities on eggs externally. Eggs were treated on day 3 of development with microlitre applications of MeHg in a carrier. Dose-related effects on survival, growth and abnormal development were recorded. Almost half of the mercury applied entered the eggs past the shell membranes within several days of treatment. Malformations occurred at  $1 \mu\text{g}\cdot\text{egg}^{-1}$  ( $18 \mu\text{g}\cdot\text{kg}^{-1}$ )<sup>32</sup>; decreased embryonic growth at  $9 \mu\text{g}\cdot\text{egg}^{-1}$  ( $164 \mu\text{g}\cdot\text{kg}^{-1}$ ). With the application of  $27 \mu\text{g}\cdot\text{egg}^{-1}$  ( $490 \mu\text{g}\cdot\text{kg}^{-1}$ ), only 63% of the embryos were alive by 18 days of development (Hoffman and Moore 1979). Residue analyses were made that could be compared with egg concentrations in field studies.

#### **8.4 Toxicity to Piscivorous Fish**

Fish appear to be tolerant to large body burdens of MeHg. Fingerling rainbow trout that were fed a diet containing 4, 8, 16 and  $24 \text{ mg}\cdot\text{kg}^{-1}$  methylmercuric chloride survived and exhibited few signs of toxicity even at the highest dosages (Wobeser 1975). Body concentrations were consistently higher than the food concentrations, with over  $30 \text{ mg}\cdot\text{kg}^{-1}$  measured in a fish fed the  $24 \text{ mg}\cdot\text{kg}^{-1}$  dosage for 15 weeks. In the final five weeks of the experiment, fish fed the two highest dosages showed significantly lower weight gains than the lower dosage groups. The lack of damage to the nervous system was thought to be because the MeHg did not reach toxic levels.

Realistic levels of MeHg in the diet caused sublethal effects to juvenile walleye. Friedmann et al. (1996) fed groups of walleye either 0, 0.1 or  $1.0 \mu\text{g}\cdot\text{g}^{-1}$  MeHg that had been injected into catfish or fathead minnow tissue. After six months, the higher dosage significantly impaired fish growth (length, weight) as well as gonadal development in the male but not the female fish. Pathohistological analysis indicated that testes of fish fed the high ( $1.0 \mu\text{g}\cdot\text{g}^{-1}$ ) MeHg diet showed significant alterations with multifocal cell atrophy compared to controls. Some testicular atrophy was observed to a lesser degree in fish fed the low ( $0.1 \mu\text{g}\cdot\text{g}^{-1}$ ) MeHg diet. Ovaries from both control and MeHg fed fish did not contain any predominant lesions. Adult walleye in the wild are more likely to consume lower trophic level fish, which realistically, may contain the lower concentration of MeHg tested here.

---

<sup>32</sup> Based on an average mallard egg weight of 55 g (Wolfe et al. 1998).

**Table 9. Consumption parameters for mink, otter, kingfisher, loon, osprey, and eagle**

| Species   | Body Weight (kg)            | Ingestion Rate or Food Intake (kg·d <sup>-1</sup> ) | Drinking Rate L·d <sup>-1</sup> | Trophic Level of Food Source | % diet at each trophic level | Reference   |
|---|-----------------------------|---|---------------------------------|------------------------------|------------------------------|---|
| Mink<br>Wild<br>Wild female<br>Captive male<br>Captive female | 0.80<br>0.6<br>1.82<br>0.87 | 0.178<br>0.143<br>0.22<br>0.14                      | 0.081                           | 3                            | 90                           | US EPA 1997b<br>CCME1998<br>Blevins and Aulerich 1981 |
| Otter   | 7.40<br>8.0                 | 1.220<br>0.8  | 0.600                           | 3,4                          | 80,20                        | US EPA 1997b<br>CCME 1998                             |
| Belted Kingfisher   | 0.15                        | 0.075   | 0.017                           | 3                            | 100                          | US EPA 1997b  |
| Loon  | 4.0<br>4.134                | 0.8<br>0.73   | 0.14                            | 3                            | 100                          | US EPA 1997b<br>CCME 1998                             |
| Osprey  | 1.50                        | 0.300   | 0.077                           | 3                            | 100                          | US EPA 1997b  |
| Eagle   | 4.60<br>4.5                 | 0.50<br>0.5   | 0.160                           | 3,4,5 <sup>a</sup>           | 74,18,8                      | US EPA 1997b<br>CCME 1998                             |

Notes:

<sup>a</sup> Level 5 indicates the eagle may consume other piscivorous birds.



## 9.0 EXISTING TISSUE RESIDUE GUIDELINES

### 9.1 Human Health

There are two basic methods of setting human consumption guidelines. In the past, the levels were usually based on a concentration in tissue to be consumed. This is still a practical way for jurisdictions to know how much mercury is in fish or shellfish, but the levels are not necessarily related to the amount of mercury consumed by an individual. Increasingly, guidelines are based on a daily or weekly dosage (i.e.,  $\mu\text{g MeHg}\cdot\text{kg}^{-1}\text{bw}\cdot\text{d}^{-1}$ ) considered to be safe for consumption.

The Canadian and the WHO guidelines for food safety, based on technical reports from annual meetings of the joint FAO/WHO Expert Committee on Food Additives, recommend that the total concentration of mercury in the edible portion of commercial fish must not exceed  $0.5\ \mu\text{g}\cdot\text{g}^{-1}$  on a wet weight basis (WHO 1972, 1976, 1990, HWC 1990; OECD 1994). Swordfish and shark are exempted, and consumption of these species should not exceed one meal per week. Many other countries have adopted the FAO/WHO tissue guideline for commercial fish consumption of  $0.5\ \mu\text{g}\cdot\text{g}^{-1}$ , and relatively few countries have set the higher limit recommended by the US EPA, of  $1.0\ \mu\text{g}\cdot\text{g}^{-1}$  (see Table 10).

Canada has qualified the concentration-based value on the frequency of consumption, which effectively considers the dosage of mercury consumed. Women of childbearing age and children should consume only fish at or below  $0.5\ \mu\text{g}\cdot\text{g}^{-1}$ . From  $0.5$  to  $1.0\ \mu\text{g}\cdot\text{g}^{-1}$ , the average adult eating fish temporarily over a short period may consume up to ten meals per week (2300 g). If consumed throughout the year, fish containing  $0.5$  to  $1.0\ \mu\text{g}\cdot\text{g}^{-1}$  MeHg should be consumed only once per week (200 g). Above  $1.0\ \mu\text{g}\cdot\text{g}^{-1}$ , the average adult eating fish temporarily over a short period should consume no more than seven meals per week (1500 g), and if eaten throughout the year, fish measuring above  $1.0\ \mu\text{g}\cdot\text{g}^{-1}$  should be consumed no more than once per week (portion of 140 g). In Canada, fish measuring over  $1.5\ \mu\text{g MeHg}\cdot\text{g}^{-1}$  are not recommended for human consumption at any time (Appendix D; WHO 1972; 1976; HWC 1990).

The WHO evaluated relationships between response and body burden (hair or blood mercury concentrations) and determined that no adverse effects (NOAEL) were detected with long-term daily methylmercury intakes of  $3\text{--}7\ \mu\text{g}\cdot\text{kg}^{-1}\text{bw}$  (hair mercury concentrations of approximately  $50\text{--}125\ \mu\text{g}\cdot\text{g}^{-1}$ ). They recommend that adults consume no more than  $0.23\ \text{mg MeHg}$  per week, or  $0.47\ \mu\text{g}\cdot\text{kg}^{-1}\text{bw}\cdot\text{d}^{-1}$  (Table 10). They postulate that pregnant women (i.e., fetus) may suffer effects at lower methylmercury exposure levels than non-pregnant adults, suggesting a greater risk for pregnant women (WHO 1990).

The U.S. Food and Drug Administration (FDA) advises persons other than pregnant women and women of child-bearing age to limit their consumption of fish species with methylmercury levels of  $1\ \mu\text{g}\cdot\text{g}^{-1}$  to approximately 200 g per week (one serving). For fish with levels averaging  $0.5\ \mu\text{g}\cdot\text{g}^{-1}$ , regular consumption should be limited to about 500 g per week (two servings).

The FDA action level is based on consideration of the tolerable daily intake (TDI) for MeHg, as well as information on seafood consumption and associated exposure to MeHg. The TDI is the amount of MeHg that can be consumed daily over a long period of time with a reasonable certainty of no harm to adults. The U.S. FDA (and WHO) established a TDI based on a weekly tolerance of  $0.3\ \text{mg}$  of THg per 70 kg person, of which no more than  $0.23\ \text{mg}$  should be present as MeHg. These amounts are equivalent to  $5$  and  $3.3\ \mu\text{g}\cdot\text{kg}^{-1}\text{bw}$ , respectively. Using the values for MeHg, this tolerable level would correspond to approximately  $230\ \mu\text{g}$  per week for a 70 kg person or  $33\ \mu\text{g}$  per person per day, which is the same level of intake recommended for

long term, frequent consumption of fish in Canada. The TDI was calculated from data developed in part by Swedish studies of Japanese individuals poisoned in the episode of Niigata which resulted from the consumption of contaminated fish and shellfish and the consideration of other studies of fish-eating populations (US EPA 1997a).

The US EPA has designated a reference dose (RfD) for humans of  $0.1 \mu\text{g MeHg}\cdot\text{kg}^{-1}\text{bw}\cdot\text{d}^{-1}$ . Another agency, the ATSDR (Agency for Toxic Substances and Disease Registry) formerly recommended a minimum risk level (MRL) of  $0.1 \mu\text{g}\cdot\text{kg}^{-1}\text{bw}\cdot\text{d}^{-1}$ , but they have increased this value to  $0.3 \mu\text{g}\cdot\text{kg}^{-1}\text{bw}\cdot\text{d}^{-1}$  based on the results of new studies of fish and marine mammal consuming populations in the Seychelles Islands (Indian Ocean) and Faroe Islands (North Atlantic Ocean) (US EPA 1997c; ATSDR 1999)<sup>33</sup>. Some are concerned that such a relaxation of the ATSDR standard could jeopardise efforts to curb mercury emissions (Renner 1998). Note that the ATSDR value is a standard (enforceable by law), whereas the EPA RfD is a guideline only.

Thus far, jurisdictions have emphasised reducing the consumption of contaminated fish rather than reducing mercury inputs that cause fish levels to be elevated. This has implications to aboriginal populations that derive important nutritional benefits through consumption of fish. According to the Cree Board of Health and Social Services of James Bay, in most instances limiting fish consumption to prevent mercury contamination carries risks that may be higher than the mercury itself (C. Dumont 1999, pers. com.). They note that fish consumption helps to prevent diseases such as heart disease, diabetes, mellitus and arthritis.

## 9.2 Wildlife

The US EPA has set water quality guidelines that are associated with tissue residue guidelines (reference doses, RfD) to protect mammal and bird consumers of fish. A wildlife criterion (WC) value is defined as the concentration of mercury in water which, if not exceeded, protects avian and mammalian wildlife populations from adverse effects resulting from ingestion of surface waters and from ingestion of aquatic life taken from these surface waters. Thus, the WC is the highest aqueous concentration (dissolved and colloid bound) of mercury that causes no significant reduction in growth, reproduction, viability or usefulness (in a commercial or recreational sense) of a population of animals exposed over multiple generations (US EPA 1997b).

In the U.S., two separate criteria values, expressed as filtered water concentrations, were derived for THg and MeHg. The total Hg criteria was based on a model proposed by the GLWQI (Great Lakes Water Quality Initiative; US EPA 1995c,d), and the MeHg values (WC) were developed as a modification of the GLWQI model by US EPA in their "Mercury Study Report to Congress" (US EPA 1997b,c). The models are based on potential bioaccumulation (BAF<sup>34</sup>) from water and from diet of the mercury species. Because the use of THg rather than MeHg was a source of criticism of the initially proposed WC (and BAF) values, US EPA developed new BAF results for MeHg. The GLWQI was developed based on a NOAEL for piscivorous wildlife of 1.1 ppm, compared with a NOAEL in the report to Congress of 0.33 ppm (both from Wobeser's research, 1975, 1976). The GLWQI value employed an uncertainty factor (UF) of ten, compared with an UF of three in the report to Congress.

The GLWQI for THg was intended to protect wildlife in the Great Lakes region with a value of

---

<sup>33</sup> April 19, 1999 update: <http://www.atsdr.cdc.gov>

<sup>34</sup> BAF = bioaccumulation factor

## Canadian Tissue Residue Guidelines: Methylmercury

1300  $\text{pg}\cdot\text{L}^{-1}$  proposed (US EPA 1997b). The US EPA report to Congress recommended a WC for protection of piscivorous wildlife species of  $50 \text{ pg MeHg}\cdot\text{L}^{-1}$ , which was the lower value of the calculated average avian wildlife criterion of  $61 \text{ pg}\cdot\text{L}^{-1}$  and average piscivorous mammalian wildlife criterion of  $50 \text{ pg}\cdot\text{L}^{-1}$ . This value corresponds to a *total* mercury concentration in the water column of  $641 \text{ pg}\cdot\text{L}^{-1}$ <sup>35</sup> and MeHg concentrations in fish of 0.077 ppm (forage fish = trophic level 3) and 0.346 ppm (piscivorous fish = trophic level 4). The WC, like the human RfD, is predicted to expose wildlife to a safe dose over a lifetime. Recommended water concentrations of  $50 \text{ pg MeHg}\cdot\text{L}^{-1}$  and  $641 \text{ pg THg}\cdot\text{L}^{-1}$  are intended to protect avian wildlife from ingesting more than the safe daily dose i.e., the RfD of  $21 \mu\text{g}\cdot\text{kg}^{-1} \text{ bw}\cdot\text{d}^{-1}$ , and to protect semi-aquatic mammals from ingesting more than  $18 \mu\text{g}\cdot\text{kg}^{-1} \text{ bw}\cdot\text{d}^{-1}$  (US EPA 1997a,b).

The US EPA criteria for wildlife differs fundamentally from Canadian guidelines in that they are based on water concentrations rather than on tissue concentrations that might be consumed by piscivorous wildlife. In other words, the US EPA values originate several steps below the trophic level that the wildlife are feeding on. Given the complexity of mercury behaviour in the environment, including the fact that there may be no relationship between mercury in the water column (especially total Hg), and mercury found in fish, a criterion based on BAF values calculated from water concentrations is less likely to accurately predict MeHg in animal tissues. The report to Congress emphasises that there is a high degree of uncertainty with the BAF calculations. However, at least one study was supportive of the GLWQI (THg) value on a contaminated lake in California (Wolfe and Norman 1998). Also, US EPA's report to Congress did consider a daily dose in the diet (RfD) that should result in no adverse effects.

---

<sup>35</sup> THg water concentration assumes a U.S. national average of 7.8% MeHg.

**Table 10. Worldwide survey of mercury tissue residue guidelines (concentrations or daily limits) for mercury in fish and fisheries products.**

| Country or Agency | Human Consumption Guideline  | Wildlife TRG  | Reference   |
|-------------------|--|---|---|
| Australia         | 0.5 $\mu\text{g}\cdot\text{g}^{-1}$ in fish, crustaceans, molluscs   | --  | OECD 1994   |
| Brazil            | 0.5 $\mu\text{g}\cdot\text{g}^{-1}$ - legal limit for hazardous metals   | --  | US EPA 1989, c.f. Haines et al. 1994              |
| Canada            | 0.5 $\mu\text{g}\cdot\text{g}^{-1}$ in fish <sup>a</sup><br>0.2 $\mu\text{g}\cdot\text{g}^{-1}$ for subsistence fish consumers <sup>b</sup>  | 0.033 $\mu\text{g}\cdot\text{g}^{-1}$ (this report) | HWC 1990; WHO 1972, 1976                          |
| Denmark           | 0.5 $\mu\text{g}\cdot\text{g}^{-1}$ - legal limit for hazardous metals   | --  | US EPA 1989, c.f. Haines et al. 1994              |
| Equador           | 1 $\mu\text{g}\cdot\text{g}^{-1}$ - legal limit for hazardous metals   | --  | US EPA 1989, c.f. Haines et al. 1994              |
| FAO/WHO           | 0.5 $\mu\text{g}\cdot\text{g}^{-1}$ in fish<br>0.23 mg MeHg per week (70 kg person)<br>0.47 $\mu\text{g MeHg}\cdot\text{kg}^{-1}\text{ bw}\cdot\text{d}^{-1}$  | --  | WHO 1972<br>WHO 1990 & Codex Alimentarius 1998    |
| Finland           | 0.5-1.0 $\mu\text{g}\cdot\text{g}^{-1}$ , once per week<br>>1.0 $\mu\text{g}\cdot\text{g}^{-1}$ , no consumption   | --  | OECD 1994   |
| France            | 0.5-0.7 $\mu\text{g}\cdot\text{g}^{-1}$ - legal limit for hazardous metals   | --  | US EPA 1989, c.f. Haines et al. 1994              |
| Germany           | 1 $\mu\text{g}\cdot\text{g}^{-1}$ - legal limit for hazardous metals   | --  | US EPA 1989, c.f. Haines et al. 1994              |
| Great Britain     | 0.2 mg MeHg per week (FAO/WHO)   | --  | OECD 1994   |
| Greece            | 0.7 $\mu\text{g}\cdot\text{g}^{-1}$ - legal limit for hazardous metals   | --  | US EPA 1989, c.f. Haines et al. 1994              |
| Hong Kong         | 0.5 $\mu\text{g}\cdot\text{g}^{-1}$ - legal limit for hazardous metals   | --  | US EPA 1989, c.f. Haines et al. 1994              |
| India             | 0.5 $\mu\text{g}\cdot\text{g}^{-1}$ - legal limit for hazardous metals   | --  | US EPA 1989, c.f. Haines et al. 1994              |
| Israel            | 0.5 $\mu\text{g}\cdot\text{g}^{-1}$ - legal limit for hazardous metals   | --  | US EPA 1989, c.f. Haines et al. 1994              |
| Italy             | 0.7 $\mu\text{g}\cdot\text{g}^{-1}$ - legal limit for hazardous metals   | --  | US EPA 1989, c.f. Haines et al. 1994              |
| Japan             | 0.3-0.4 $\mu\text{g}\cdot\text{g}^{-1}$ - legal limit for hazardous metals   | --  | US EPA 1989, c.f. Haines et al. 1994              |
| Korea             | 0.5 $\mu\text{g}\cdot\text{g}^{-1}$ - legal limit for hazardous metals   | --  | US EPA 1989, c.f. Haines et al. 1994              |
| Netherlands       | 1 $\mu\text{g}\cdot\text{g}^{-1}$ limit for hazardous metals in fish<br>0.5 $\mu\text{g}\cdot\text{kg}^{-1}\text{ bw}\cdot\text{d}^{-1}$ adults<br>0.1 $\mu\text{g}\cdot\text{kg}^{-1}\text{ bw}\cdot\text{d}^{-1}$ pregnant/lactating women | --  | US EPA 1989, c.f. Haines et al. 1994<br>OECD 1994 |
| New Zealand       | 0.5 $\mu\text{g}\cdot\text{g}^{-1}$ - legal limit for hazardous metals   | --  | US EPA 1989, c.f. Haines et al. 1994              |
| Philippines       | 0.5 $\mu\text{g}\cdot\text{g}^{-1}$ - legal limit for hazardous metals   | --  | US EPA 1989, c.f. Haines et al. 1994              |
| Spain             | 0.5 $\mu\text{g}\cdot\text{g}^{-1}$ - legal limit for hazardous metals   | --  | US EPA 1989, c.f. Haines et al. 1994              |
| Sweden            | 1 $\mu\text{g}\cdot\text{g}^{-1}$ - legal limit for hazardous metals   | --  | US EPA 1989, c.f. Haines et al. 1994              |

## Canadian Tissue Residue Guidelines: Methylmercury

| Country or Agency                       | Human Consumption Guideline  | Wildlife TRG  | Reference   |
|---|--|---|---|
| Switzerland                             | 0.5 $\mu\text{g}\cdot\text{g}^{-1}$ - legal limit for hazardous metals   | --  | US EPA 1989, c.f. Haines et al. 1994              |
| Thailand                                | 0.5 $\mu\text{g}\cdot\text{g}^{-1}$ - legal limit for hazardous metals   | --  | US EPA 1989, c.f. Haines et al. 1994              |
| US ATSDR <sup>c</sup>                   | 0.5 $\mu\text{g}\cdot\text{kg}^{-1}$ bw·d <sup>-1</sup> , Nov. 1997 increased from 0.1 $\mu\text{g}\cdot\text{kg}^{-1}$ bw·d <sup>-1</sup> | --  | US EPA 1997c                                      |
| US EPA                                  | 0.1 $\mu\text{g}\cdot\text{kg}^{-1}$ bw·d <sup>-1</sup>  | 21 $\mu\text{g}\cdot\text{kg}^{-1}$ bw per d birds<br>18 $\mu\text{g}\cdot\text{kg}^{-1}$ bw per d mammals <sup>d</sup> | US EPA 1997c                                      |
| US Food & Drug Admin.                   | 1.0 $\mu\text{g}\cdot\text{g}^{-1}$  | --  | US EPA 1997c                                      |
| Venezuela                               | 0.1-0.5 $\mu\text{g}\cdot\text{g}^{-1}$ - legal limit for hazardous metals   | --  | US EPA 1989, c.f. Haines et al. 1994              |
| World Health Organization - see FAO/WHO |  |   |   |
| Zambia                                  | 0.2-0.3 $\mu\text{g}\cdot\text{g}^{-1}$ - legal limit for hazardous metals   | --  | US EPA 1989, c.f. Haines et al. 1994 <sup>e</sup> |

### Notes:

- <sup>a</sup> WHO/Canada- Shark, tuna, and swordfish exempt should be eaten no more than once per week.
- <sup>b</sup> Subsistence level issued by Health Canada's Medical Services Branch in 1979, and reiterated in 1984 (N.P. McEwen 1999, Health Canada, pers. com.; App. D)
- <sup>c</sup> Agency for Toxic Substances and Disease Registry
- <sup>d</sup> US EPA wildlife TRG (RfD) are in review, and are based on wildlife criteria (WC) which are water concentrations of 50 pg MeHg·L<sup>-1</sup> and 641 pg total.
- <sup>e</sup> Haines et al. 1994 comprises the supporting documentation for MacDonald 1994.

## 10.0 CANADIAN TISSUE RESIDUE GUIDELINES FOR METHYLMERCURY

### 10.1 Protocol

TRGs are concentrations of environmental contaminants (that have a potential to biomagnify in the tissues of aquatic organisms; e.g., fish) recommended to protect wildlife that consume aquatic biota in freshwater, estuarine, or marine ecosystems. The protocol for the derivation of Canadian tissue residue guidelines (TRGs) for the protection of wildlife that consume aquatic biota is fully described in "Protocol for the derivation of Canadian tissue residue guidelines for the protection for wildlife that consume aquatic biota" (CCME 1998). While the protocol promotes a standard method of derivation, it is intended as a flexible procedural guide that should not replace best scientific judgement when developing guidelines. Note that the current protocol is intended as a working document for the testing and application of the methodology (CCME 1998).

Briefly, TRG derivation follows the evaluation of toxicological data from a range of avian and mammalian species. The goal is to protect the most sensitive species and life stages of wildlife. Certain minimum toxicological data set requirements must be met (see 3.3.1 - 3.4.3 in CCME 1998). The steps in derivation include:

1. Selecting high quality chronic studies from which LOAEL and/or NOAEL values are available; values may be adjusted to daily intake rates in units of  $\text{mg}\cdot\text{kg}^{-1}$  bw per day if not originally presented as such
2. Selecting an Uncertainty Factor (UF) based on the reality of accurate extrapolation from differences in gender, life stage, other species, lab-to-field, and other factors;
3. Calculation of Tolerable Daily Intakes (TDI) in  $\text{mg}\cdot\text{kg}^{-1}$  bw per day as the geometric mean of the selected LOAEL and NOAEL<sup>36</sup> divided by the UF. TDIs are calculated separately for mammalian and avian wildlife species.
4. Calculation of Reference Concentrations (RCs) for a wide range of wildlife species, where  $\text{RC} = \text{TDI} \times (\text{Body Weight}/\text{Food Intake})$ ;  $\text{bw}=\text{kg}$ ;  $\text{FI}=\text{kg}\cdot\text{d}^{-1}$ ).

As the lowest TDI will not necessarily result in the lowest acceptable dietary concentration, the avian and mammalian TDIs are used in conjunction with the body weights (bw) and daily food intake rates (FI) of Canadian wildlife species to calculate RCs. The lowest RC is carried forward as the TRG. For the purposes of deriving national TRG values, the mammalian and avian RCs must be as inclusive as possible to accommodate all species and regions in Canada. Therefore, the derivation is based on the highest mammalian and avian FI:bw ratios known for Canadian wildlife, namely 0.24 for mink and 0.94 for Wilson's storm petrel (CCME 1998).

## 10.2 Guideline Derivation and Rationale

As detailed in chapter 8, there are a number of studies from both laboratory and wildlife species on MeHg toxicity. According to the CCME protocol, wildlife species are preferred and as such, high quality studies on mallards and minks were considered in calculating the TRG.

### 10.2.1 Avian Species

Three generations of mallard ducks fed a contaminated diet indicated that successive generations are susceptible to MeHg effects although there was no evidence that the effects became progressively more severe from one generation to the next (Heinz 1979). A dietary concentration of  $0.5 \text{ mg}\cdot\text{kg}^{-1}$  dw of methylmercury is the LOAEL because it resulted in reduced growth and survival in offspring of second generation hens and in ecologically-relevant behavioural effects in offspring of third-generation hens (Heinz 1976b, 1979). In addition, hens fed  $0.5 \text{ mg}\cdot\text{kg}^{-1}$  dw laid a greater percentage of eggs outside their nest boxes than control birds although the effect was statistically significant only in the second generation and when results for all three generations were combined (Heinz 1979). The nominal dietary concentration of  $0.5 \text{ mg}\cdot\text{kg}^{-1}$  dw is equivalent to an intake rate of  $75 \mu\text{g}\cdot\text{kg}^{-1}$  bw per day assuming a measured dietary concentration of  $0.48 \text{ mg}\cdot\text{kg}^{-1}$  dw<sup>37</sup> and a food consumption rate of 0.156 kg of dry food per kg of body weight per day<sup>38</sup> (Heinz 1979). The NOAEL of  $13 \mu\text{g}\cdot\text{kg}^{-1}$  bw per day was calculated by dividing the LOAEL by 5.6 (CCME 1993). This calculated NOAEL of  $13 \mu\text{g}\cdot\text{kg}^{-1}$  bw per day is higher than the intake rate ( $6.4 \mu\text{g}\cdot\text{kg}^{-1}$  bw per day) calculated for those

---

<sup>36</sup> NOAEL may be estimated if necessary as  $\text{LOAEL} \div 5.6$  (CCME 1993)

<sup>37</sup> The measured dietary concentration is  $0.48 \text{ mg}\cdot\text{kg}^{-1}$  dw, when the means for each generation (0.53, 0.47, and  $0.43 \text{ mg}\cdot\text{kg}^{-1}$  dw) are averaged (Heinz 1979). Individual measured concentrations could not be used because they were not reported for the second and third generations (Heinz 1976b; 1979).

<sup>38</sup> When results were combined, second and third generation mallards fed the  $0.5 \text{ mg}\cdot\text{kg}^{-1}$  dw treatment diet consumed 0.156 kg of dry food per kg of body weight per day; food consumption of the first generation was not recorded (Heinz 1979).

birds fed control feed ( $0.05 \text{ mg}\cdot\text{kg}^{-1} \text{ dw}$ ;  $0.128 \text{ kg}$  of dry food per  $\text{kg}$  of body weight per day).

The avian TDI of  $31 \text{ }\mu\text{g}\cdot\text{kg}^{-1} \text{ bw}$  per day (Table 11) was calculated as the geometric mean of the LOAEL ( $75 \text{ }\mu\text{g}\cdot\text{kg}^{-1} \text{ bw}$  per day) and NOAEL ( $13 \text{ }\mu\text{g}\cdot\text{kg}^{-1} \text{ bw}$  per day) without the application of an uncertainty factor (UF). Not applying an UF is justified in this case because the experiments were conducted on a Canadian wildlife species (mallard), over three generations (a true chronic study), and at MeHg dosages that would commonly be found in wild fish. Dividing the TDI by the highest FI:bw ratio for wild birds (0.94 for Wilson's storm petrel) results in an avian RC of  $33 \mu\text{g}\cdot\text{kg}^{-1}$  diet on a wet weight basis (Tables 11 and 12).

For perspective, the avian TDI of  $31 \text{ }\mu\text{g}\cdot\text{kg}^{-1} \text{ bw}$  per day is comparable to the reference dose (RfD) recommended by the US EPA ( $21 \text{ }\mu\text{g}\cdot\text{kg}^{-1} \text{ bw}$  per day) to protect avian wildlife (US EPA 1997a), although the derivation procedures of the two values are not the same. The avian TDI is about 2.4 times lower than the daily ingested dose that resulted in the LOAEL for mallards discussed above. In contrast, the avian TDI is 66 times higher than the daily intake recommended by the WHO for humans ( $0.47 \text{ }\mu\text{g}\cdot\text{MeHg}\cdot\text{kg}^{-1} \text{ bw}$  per day; WHO 1990).

Weaknesses in the critical study used to calculate the avian RC include the fact that the mercury dosage was added to dry food (rather than using contaminated fish), and was fed to a species (mallard) that is not principally a piscivore. Moreover, although a dietary concentration of  $0.5 \text{ mg}\cdot\text{kg}^{-1} \text{ dw}$  of methylmercury produced adverse effects in all three generations, effects were not consistent from one generation to the next. In addition, because effects were found at the lowest test dosage, the NOAEL had to be calculated from the LOAEL. Nevertheless, the long-term chronic toxicity experiment completed by Heinz was the most comprehensive MeHg study available.

### 10.2.2 Mammalian Species

A study on domestic cats indicated that cats are sensitive to MeHg exposure (Charbonneau et al. 1976). As wild cats rarely consume aquatic biota, and because good quality studies on mink were available, the cat and Wistar rat assays (Munro et al. 1980) were not used for the derivation, but were referred to for comparison purposes only.

The most sensitive results from wildlife-based studies included reduced fertility and neurotoxic effects in female mink followed by death ( $>50\%$ ) at  $0.9 \text{ mg}\cdot\text{kg}^{-1}$  ( $144 \text{ }\mu\text{g}\cdot\text{kg}^{-1} \text{ bw}$  per day) (Chamberland et al. 1996). No ecologically relevant effects including clinical signs, neurological damage or impaired reproduction were observed in female mink at  $0.5 \text{ mg}\cdot\text{kg}^{-1}$  ( $80 \text{ }\mu\text{g}\cdot\text{kg}^{-1} \text{ bw}$  per day) and as such, this exposure was used as the NOAEL (Chamberland et al. 1996; Laperle et al. 1998).

Strengths of the critical study include the use of contaminated fish from James Bay reservoirs, a critical wildlife species (mink), and three treatment levels (Chamberland et al. 1996). Neurological effects considered ecologically relevant include ataxia, staggering, splaying or paralysis in the hind limbs, and anorexia. These effects and others were reported in many of the studies that described MeHg neurotoxicity. Such effects would render an animal unable to properly feed, escape from predators, reproduce, and would increase its susceptibility to other stressors.

The TDI for mammals of  $22 \text{ }\mu\text{g}\cdot\text{kg}^{-1} \text{ bw}$  per day was calculated as the geometric mean of the LOAEL ( $144 \text{ }\mu\text{g}\cdot\text{kg}^{-1} \text{ bw}$  per day) and NOAEL ( $80 \text{ }\mu\text{g}\cdot\text{kg}^{-1} \text{ bw}$  per day) divided by an uncertainty factor of 5. An UF of 5 was chosen because the endpoint of the LOAEL was death, the study

## Canadian Tissue Residue Guidelines: Methylmercury

lasted just over three months (subchronic), and to permit extrapolation to other wildlife species. A mammalian RC of  $92 \mu\text{g}\cdot\text{kg}^{-1}$  diet on a wet weight basis was obtained by dividing the TDI by the highest FI:bw ratio for wild mammals (0.24 for female mink; Tables 8 and 9).

The mammalian TDI is comparable to the US EPA's reference dose (RfD) for semi-aquatic mammals of  $18 \mu\text{g}\cdot\text{kg}^{-1}$  bw per day and is approximately equal to the chronic NOAEL ( $20 \mu\text{g}\cdot\text{kg}^{-1}$  bw per day) estimated from a two-year feeding study with domestic cats (Charbonneau et al. 1976). The mammalian TDI is 47 times higher than the intake recommended by the WHO for humans ( $0.47 \mu\text{g}\cdot\text{kg}^{-1}$  bw per day; WHO 1990).



### 10.3 Methylmercury Guideline

For the protection of all life stages of all birds and mammals that consume aquatic organisms, the lowest RC,  $33 \mu\text{g}\cdot\text{kg}^{-1}$  for Wilson's storm petrel, is recommended as the TRG. This guideline refers to concentration of methylmercury on a wet weight basis in the aquatic organisms (i.e., fish and other aquatic biota) that is not expected to adversely affect the wildlife that consume them. To protect all wildlife at a site, TRGs should be applied to the highest known aquatic trophic level.

For the purposes of deriving a national TRG value, the avian and mammalian RCs must be as inclusive as possible to accommodate all species and regions in Canada and are therefore based on the highest mammalian and avian FI:bw ratios known for Canadian wildlife, namely 0.24 for mink and 0.94 for Wilson's storm petrel (CCME 1998). We recognise that in areas where, for example, Wilson's storm petrel is not found, the use of the lowest RC (highest FI:BW ratio) may not be appropriate; in which case, RCs for resident wildlife may be applied. For this reason, a list of suggested RCs for a suite of mammalian and avian receptors is provided (see Table 12).

The recommended MeHg TRG of  $33 \mu\text{g}\cdot\text{kg}^{-1}$  diet, ww is lower than most MeHg concentrations that have been measured in sport fish from Canadian freshwaters (see chapter 7; Appendix B-3). Concern about human consumption advisories has led to the testing of commercially important species. Fish typically consumed by wildlife are much smaller and are from lower trophic levels, and could have proportionately less Hg than their larger commercial sport fish counterparts. As such, the reader is cautioned about making inferences regarding risks to wildlife from mercury levels in large (>35 cm) fish. The wildlife TRG, including any calculated site-specific values, must be relatively low due to the high proportion of fish in the diet of the species in question.

**Table 11. TRG derivation and calculations.**

STEP 1: CALCULATION OF TOLERABLE DAILY INTAKE

|               |   |  |
|---------------|---|--|
| $TDI_{AVIAN}$ | = | $(LOAEL \times NOAEL)^{0.5} / \text{Uncertainty factor}$   |
| Where:        |   |  |
| LOAEL         | = | $0.48 \text{ mg}\cdot\text{kg}^{-1} \text{ food} \times 0.156 \text{ kg food}\cdot\text{kg}^{-1} \text{ bw}$ |
|               | = | $75 \text{ }\mu\text{g}\cdot\text{kg}^{-1} \text{ bw per day (Heinz 1979)}$                                  |
| and,          |   |  |
| NOAEL         | = | $LOAEL/5.6 \text{ (CCME 1993)}$  |
|               | = | $13 \text{ }\mu\text{g}\cdot\text{kg}^{-1} \text{ bw per day}$   |
| Therefore,    |   |  |
| $TDI_{AVIAN}$ | = | $(75 \times 13)^{0.5} / 1$   |
|               | = | $31 \text{ }\mu\text{g}\cdot\text{kg}^{-1} \text{ bw per day.}$  |

|                   |   |  |
|-------------------|---|--|
| $TDI_{MAMMALIAN}$ | = | $(LOAEL \times NOAEL)^{0.5} / \text{Uncertainty factor}$   |
| Where:            |   |  |
| LOAEL             | = | $0.9 \text{ mg}\cdot\text{kg}^{-1} \text{ food} \times 0.16 \text{ kg food}\cdot\text{kg}^{-1} \text{ bw}$ |
|                   | = | $144 \text{ }\mu\text{g}\cdot\text{kg}^{-1} \text{ bw per day (Chamberland et al. 1996)}$                  |
| and,              |   |  |
| NOAEL             | = | $0.5 \text{ mg}\cdot\text{kg}^{-1} \text{ food} \times 0.16 \text{ kg food}\cdot\text{kg}^{-1} \text{ bw}$ |
|                   | = | $80 \text{ }\mu\text{g}\cdot\text{kg}^{-1} \text{ bw per day.}$  |
| Therefore,        |   |  |
| $TDI_{MAMMALIAN}$ | = | $(144 \times 80)^{0.5} / 5$  |
|                   | = | $22 \text{ }\mu\text{g}\cdot\text{kg}^{-1} \text{ bw per day.}$  |

STEP 2: CALCULATION OF REFERENCE CONCENTRATIONS

|               |   |  |
|---------------|---|--|
| $RC_{AVIAN}$  | = | $TDI_{AVIAN} / (FI:BW_{MAX})$                  |
| Where:        |   |  |
| $TDI_{AVIAN}$ | = | $31 \mu\text{g}\cdot\text{kg}^{-1}$ bw per day |
| and,          |   |  |
| $FI:BW_{MAX}$ | = | 0.94 (i.e., FI:BW for Wilson's storm petrel).  |
| Therefore,    |   |  |
| $RC_{AVIAN}$  | = | $31 / 0.94$                                    |
|               | = | $33 \mu\text{g}\cdot\text{kg}^{-1}$ diet, ww.  |

|                   |   |  |
|-------------------|---|--|
| $RC_{MAMMALIAN}$  | = | $TDI_{MAMMAL} / (FI:BW_{MAX})$                 |
| Where:            |   |  |
| $TDI_{MAMMALIAN}$ | = | $22 \mu\text{g}\cdot\text{kg}^{-1}$ bw per day |
| and,              |   |  |
| $FI:BW_{MAX}$     | = | 0.24 (i.e., FI:BW for mink).                   |
| Therefore,        |   |  |
| $RC_{MAMMALIAN}$  | = | $22 / 0.24$                                    |
|                   | = | $92 \mu\text{g}\cdot\text{kg}^{-1}$ diet, ww   |

STEP 3:

|   |
|---|
| $TRG = \text{LOWEST } RC = RC_{AVIAN} = 33 \mu\text{g}\cdot\text{kg}^{-1}$ diet, ww |
|---|

Canadian Tissue Residue Guidelines: Methylmercury

**Table 12. Reference concentrations (RCs;  $\mu\text{g}\cdot\text{kg}^{-1}$  diet, ww)<sup>a,b</sup> for Canadian wildlife species derived from the lowest avian (31  $\mu\text{g}\cdot\text{kg}^{-1}$  bw per day) or mammalian (22  $\mu\text{g}\cdot\text{kg}^{-1}$  bw per day) tolerable daily intakes (TDIs) for methylmercury and food intake (FI) to body weight (BW) ratios<sup>c</sup>.**

| Species  | FI:BW Ratio | RC  | Species   | FI:BW Ratio | RC        |
|--|-------------|-----|---|-------------|-----------|
| <b>Birds</b>                                       |             |     | <b>Birds</b>  |             |           |
| <u>Anseriformes</u>                                |             |     | <u>Ciconiiformes</u>                                      |             |           |
| Buffle head ( <i>Bucephala albeola</i> )           |             |     | Great blue heron ( <i>Ardea herodias</i> )                |             |           |
| Male   | 0.36        | 86  | Male  | 0.21        | 148       |
| Female   | 0.42        | 74  | Female  | 0.22        | 141       |
| Common goldeneye ( <i>Bucephala clangula</i> )     |             |     | Green-backed heron ( <i>Butorides striatus</i> )          | 0.24        | 129       |
| Male   | 0.29        | 107 |   |             |           |
| Female   | 0.31        | 100 | <u>Procellariiformes</u>                                  |             |           |
| Mallard ( <i>Anas platyrhynchos</i> )              | 0.23        | 135 | <b>Wilson's storm-petrel (<i>Oceanites oceanicus</i>)</b> | <b>0.94</b> | <b>33</b> |
| Oldsquaw ( <i>Clangula hyemalis</i> )              |             |     | Fork-tailed storm-petrel ( <i>Oceanodroma furcata</i> )   | 0.73        | 42        |
| Male   | 0.29        | 107 | Northern fulmar ( <i>Fulmarus glacialis</i> )             |             |           |
| Female   | 0.31        | 100 | Male  | 0.34        | 91        |
| Wood duck ( <i>Aix sponsa</i> )                    |             |     | Female  | 0.38        | 82        |
| Male   | 0.34        | 91  |   |             |           |
| Female   | 0.35        | 89  | <b>Mammals</b>  |             |           |
| American wigeon ( <i>Anas americana</i> )          |             |     | <u>Mustelidae</u>   |             |           |
| Male   | 0.32        | 97  | Sea otter ( <i>Enhydra lutris</i> )                       |             |           |
| Female   | 0.32        | 97  | Male  | 0.18        | 122       |
| Lesser Scaup ( <i>Aythya affinis</i> )             |             |     | Female  | 0.2         | 110       |
| Male   | 0.31        | 100 | American mink ( <i>Mustela vison</i> )                    |             |           |
| Female   | 0.32        | 97  | Female  | 0.24        | 92        |
| Common merganser ( <i>Mergus merganser</i> )       |             |     | River Otter ( <i>Lutra canadensis</i> )                   | 0.1         | 220       |
| Male   | 0.24        | 129 |   |             |           |
| Female   | 0.27        | 115 | <u>Pinnipedia</u>   |             |           |
| Red-breasted merganser ( <i>Mergus serrator</i> )  |             |     | Harbour Seal ( <i>Phoca vitulina</i> )                    |             |           |
| Male   | 0.21        | 148 | Male  | 0.16        | 138       |
|  |             |     | Female  | 0.17        | 129       |
| <u>Falconiformes</u>                               |             |     | Northern fur seal ( <i>Callorhinus ursinus</i> )          |             |           |
| Bald eagle ( <i>Haliaeetus leucocephalus</i> )     | 0.11        | 282 | Male  | 0.13        | 169       |
| Osprey ( <i>Pandion haliaetus</i> )                | 0.2         | 155 | Female  | 0.18        | 122       |
|  |             |     | Northern elephant seal ( <i>Mirounga angustirostris</i> ) |             |           |
| <u>Coraciiformes</u>                               |             |     | Male  | 0.08        | 275       |
| Belted kingfisher ( <i>Ceryle alcyon</i> )         | 0.5         | 62  | Female  | 0.1         | 220       |
|  |             |     | Northern seal lion ( <i>Eumetopias jubata</i> )           |             |           |
| <u>Gaviiformes</u>                                 |             |     | Male  | 0.1         | 220       |
| Common loon ( <i>Gavia immer</i> )                 | 0.18        | 172 | Female  | 0.12        | 183       |
|  |             |     | Walrus ( <i>Odobenus rosmarus</i> ), eastern Arctic race  |             |           |
| <u>Charadriiformes</u>                             |             |     | Male  | 0.11        | 200       |
| Common tern ( <i>Sterna hirundo</i> )              | 0.61        | 51  | Female  | 0.11        | 200       |
| Herring gull ( <i>Larus argentatus</i> )           |             |     | Walrus ( <i>Odobenus rosmarus</i> ), Pacific Ocean race   |             |           |
| Male   | 0.28        | 111 | Male  | 0.1         | 220       |
| Female   | 0.29        | 107 | Female  | 0.1         | 220       |
| Ring-billed gull ( <i>Larus delawarensis</i> )     |             |     |   |             |           |
| Male   | 0.17        | 182 | <u>Ursidae</u>  |             |           |
| Black-legged kittiwake ( <i>Rissa tridactyla</i> ) |             |     | Polar Bear ( <i>Ursus maritimus</i> )                     |             |           |
| Male   | 0.38        | 82  | Male  | 0.12        | 183       |
| Razorbill ( <i>Alca torda</i> )                    | 0.32        | 97  |   |             |           |
| Common murre ( <i>Uria aalge</i> )                 |             |     |   |             |           |
| Male   | 0.39        | 79  |   |             |           |
| Female   | 0.3         | 103 |   |             |           |
| Thick-billed murre ( <i>Uria lomvia</i> )          | 0.3         | 103 |   |             |           |
| Black guillemot ( <i>Cepphus grylle</i> )          | 0.4         | 78  |   |             |           |
| Atlantic puffin ( <i>Fratercula arctica</i> )      | 0.39        | 79  |   |             |           |
| Tufted puffin ( <i>Fratercula cirrhata</i> )       | 0.32        | 97  |   |             |           |

Notes:

<sup>a</sup> Reference Concentration refers to the concentration of methylmercury in aquatic biota that when consumed by wildlife is not expected to result in adverse effects.

<sup>b</sup> US EPA ranked Hg susceptibility as: kingfisher > river otter > loon = osprey = mink > bald eagle.

<sup>c</sup> FI: BW values from CCME (1998)

## 11.0 DATA GAPS

Research on methylmercury has lagged behind research on inorganic mercury because of former limitations in sample collection and analytical methodologies. As a result, a number of data gaps exist. Knowledge of specific environmental variables that influence the mercury cycle, biological accumulation, and toxicity are also contradictory. Other data gaps relate to essential toxicity and physiological studies on critical wildlife species. More information is required in the following areas:

- The influence of abundance and range of DOC types on MeHg bioavailability and accumulation by aquatic organisms;
- What levels of dietary selenium are protective, and how much of a role do internal demethylating pathways play in protecting piscivorous birds and mammals;
- Dietary absorption efficiency of MeHg from natural sources for the key piscivorous wildlife species;
- Studies that analyse for both total and MeHg so that differences between aquatic systems can be definitively linked to differences in MeHg levels. Analysing the two mercury species together will contribute to an understanding of existing data, much of which has been reported as total mercury;
- Data on MeHg levels in wildlife are needed to identify populations that are potentially at risk. Feathers and fur hold promise in this regard due to the potential for “non-invasive” determination of mercury residues;
- Whenever possible, tissue samples should be analysed for selenium. Sampling efforts with wildlife for Se and MeHg should be accompanied by analyses of likely food items;
- Chronic toxicity studies are needed for representative mammalian and avian species where a NOAEL is available in addition to a LOAEL;
- Chronic studies, if possible, should be completed with piscivorous bird species such as loons, eagles, or osprey;
- Chronic toxicity studies with river otter have never been completed;
- More specific knowledge about the diets of critical wildlife is required. Fish sampling efforts are frequently focused on species that are relevant to human consumers but that may be of little significance to wildlife. **There is an additional need to collect information for small fish (<35 cm), macroinvertebrates, amphibians and reptiles.** Seasonal and spatial effects on predation should be explored and methods developed to describe this information adequately. Additional life history data is needed to fully characterise the nature and extent of exposure to MeHg. Complicating factors must be considered, including migratory behaviours and sex-specific differences in distribution and resource allocation; and
- As the above data gaps are filled, it may be possible to extrapolate from effects of MeHg on individuals (as described in this report) to populations, communities and ecosystems.

## **12.0 GUIDELINE IMPLEMENTATION**

For substances that have a strong potential to biomagnify through the food chain, proper implementation of the TRGs requires consideration of several factors including: 1) the identification of the wildlife species requiring protection, 2) food intake: body weight ratios and diet preferences of those wildlife species, and 3) size class and trophic level of prey (diet) items on which the species of concern feeds. Implementation of TRG for MeHg by risk managers may be particularly challenging given the toxicity of methylmercury, its widespread occurrence due to both natural and anthropogenic sources, and possible adverse effects in wild populations (Wren et al. 1986; Wobeser et al. 1976a,b; G. Mierle 1999, MOE, pers. com.). The apparent stringency of this guideline is due in part to the small size and high food intake rate of the avian species on which it is based, the Wilson's storm petrel.

To protect all wildlife species, the TRG applies broadly to the highest known aquatic trophic level (CCME 1998). Environment Canada and the CCME cautiously encourage jurisdictions to modify the guidelines into site-specific objectives where necessary. This process may include taking into account species which are protected by the national guideline but which are not present, or are not reasonably expected to be present, in the sites under consideration. Guidelines may be modified also if factors affecting toxicity or bioavailability are adequately known. For example, hardness and DOC make metals less toxic to aquatic life under some conditions and guideline modification can take this into account. More specifics on guideline modification are provided in the CCME document entitled "*Scientific and Technical Guidance on Water Quality Guideline Implementation*" (in press). Once a suitable site-specific tissue residue objective is determined, it must be applied to the appropriate dietary items (prey species), in terms of both size and trophic level.

### **12.1 Use of Appropriate Wildlife Species**

The TRG was calculated from a tolerable daily intake (based on the toxicity of mercury to mallard ducks), and the highest food intake: body weight ratio for wildlife species, namely that for Wilson's storm petrel (see chapter 10). Wilson's storm petrel, as a marine bird, is not present in all jurisdictions and, therefore, may not be representative of resident species in freshwater ecosystems. Moreover, large birds, such as bald eagles, with small food intake: body weight ratios may tolerate a more highly contaminated diet than small birds such as terns which eat more food per body weight (i.e., large food intake: body weight ratios), assuming all other factors equal. For this reason a list of food intake: body weight ratios and corresponding reference concentration values are provided for a number of wildlife species (Table 12). Reference concentrations may themselves be modified, for example, by adjusting the food intake: body weight ratio on a site-specific basis for a given wildlife species.

### **12.2 Application of TRG to Appropriate Size Class of Dietary Items**

Different wildlife species consume different size classes of fish species. For example, mink typically consume fish 15 to 20 cm in length while heron consume fish 20 to 25 cm in length. Generally, wildlife do not consume fish longer than 35 cm (U.S. EPA 1997c). As a result of bioaccumulation and food chain transfer, larger fish of a given species commonly contain higher methylmercury tissue concentrations than smaller fish; therefore, the TRG or site-specific objective should be applied to the size class of fish (prey) on which the wildlife species of concern feeds. Otherwise, erroneous conclusions may be drawn regarding the potential for adverse effects in wildlife consumers of aquatic biota. Notwithstanding, wildlife species that scavenge for food consume diet items of various sizes that are sometimes in excess of 35 cm.

Ultimately, reference concentrations should be compared to the concentrations in known prey and other diet items, regardless of size class or source.

### **12.3 Trophic Level Considerations**

To develop site-specific tissue residue objectives, the trophic level at which a wildlife species feeds needs to be considered to ensure adequate protection of the resident wildlife consumers of aquatic biota. For example, a site-specific objective designed to protect wildlife that consume trophic level 2 species (e.g., common goldeneye) may not be protective of wildlife that prey on trophic level 3 species (e.g., cormorants). The following sections and Tables 10 and 11 are intended to provide general guidance necessary for the proper implementation of TRGs for bioaccumulative substances. The classification for aquatic trophic levels (Table 13; from US EPA 1995e) provides guidance on the aquatic organisms found at each trophic level, as well as the wildlife species that feed at each aquatic trophic level in freshwater environments. Feeding habits for several consumers of aquatic life are presented in Table 14 to aid guideline users in choosing an appropriate trophic level at which to apply the guideline to protect the wildlife species of concern.

#### **12.3.1 Uncertainties with Establishing Distinct Trophic Levels**

A major difficulty with the trophic levels described above is that many species feed at more than one trophic level and therefore do not fit wholly into discrete trophic levels. While species can be fairly readily identified as plants or herbivores, the carnivorous species are difficult to categorise into distinct trophic levels. For example, lake trout will consume both benthic invertebrates (trophic level 2) and small fish (trophic level 3). Mink will feed on muskrats (which feed mainly on emergent vegetation, therefore trophic level 1), shiners (trophic level 3), and some walleye (trophic level 4). In general, the closer a species is to the top of a food web, the more likely it is to feed on prey from more than one trophic level (US EPA 1995e). To overcome the uncertainty in assigning the diet of a wildlife species to a specific trophic level, the highest trophic level at which a species feeds was used for its classification.

Uncertainty also exists when attempting to estimate the food chain or web that supports a given wildlife species in a specific location. First, a food chain analysis implies a certain consistency in feeding patterns over time. The diets of animals, however, vary both with season because of changes in nutritional needs and availability of prey, and with organism age and size as larger animals can take larger prey. Thus, analysis of a particular food chain or web for an organism over a short period of time may not be indicative of the potential for bioaccumulation and biomagnification over longer time periods.

Finally, ecosystem types that would otherwise be similar (e.g., oligotrophic lakes of a certain size) may support substantially different food webs and numbers of trophic levels to the top predators depending on the history of the ecosystem, species introductions, and species loss. It has been shown that much of the large between-lake differences in contaminant levels among fish species results from differences in the length of the food chains (Rasmussen et al. 1990). This underlines the importance of understanding local food webs if the TRGs are being applied on a site-specific basis.

#### **12.3.2 Basic Aquatic Trophic Levels**

General trophic level guidance for freshwater ecosystems is provided in Table 13. Each trophic level has a general description of the species that feed within that trophic level and what their

diet resembles. This table is not intended to replace best scientific judgement at a given site.

### **12.3.3 Prey Trophic Levels of Representative Species**

Information on the feeding habits and prey trophic levels of several species in both freshwater and marine ecosystems is presented in Table 14. Again, this table is only meant for guidance and is not intended to replace best scientific judgement at a given site.



**Table 13. Basic freshwater aquatic trophic levels.**

| Trophic Level | Organisms                           | Examples  |
|---------------|-------------------------------------|---|
| <b>1</b>      | <b>Primary Producers</b>            |   |
|               | Phytoplankton                       | Free-floating unicellular algae.  |
|               | Periphyton                          | Algae attached to substrates (e.g., rocks, mud, surfaces of aquatic macrophytes).   |
|               | Emergent macrophytes                | Marsh plants rooted in the littoral zone with leaves and stems exposed (e.g., cattails and bulrushes).  |
|               | Floating aquatic macrophytes        | Higher plants that are free-floating and not rooted to a substrate (e.g., duckweed, pondweed, water hyacinth).  |
|               | Submerged aquatic macrophytes       | Higher plants rooted to a substrate in the littoral zone that are submerged underwater (e.g., <i>Hydrilla</i> ).  |
|               | Detritus                            | Dead and decaying plant and animal particulate matter. It may float or settle to the bottom.  |
| <b>2</b>      | <b>Herbivores and Detritivores</b>  |   |
|               | Planktivores                        | Generally are zooplankton that filter phytoplankton and floating detritus and bacteria from the water column.   |
|               | Order Anostraca                     | Fairy shrimp.   |
|               | Conchostraca                        | Clam shrimp.  |
|               | Cladocera                           | Water fleas (e.g., <i>Daphnia</i> ).  |
|               | Copepoda                            | Copepods.   |
|               | Diptera                             | Aquatic larvae of several species of flies.   |
|               | Class Bivalvia                      | Mussels and clams.  |
|               | Periphyton and Macrophyte Consumers | Herbivorous invertebrates that feed on the periphyton (and associated detritus) on the surfaces of plants and other aquatic substrates, or that feed on macroalgae and higher plants. |
|               | Order Amphipoda                     | Scuds, side-swimmers, or freshwater shrimp.   |
|               | Decapoda                            | Crayfish.   |
|               | Coleoptera                          | Halplidae (crawling water beetles).   |
|               | Diptera                             | Aquatic larvae of flies and midges.   |
|               | Lepidoptera                         | Caterpillars of butterflies and moths   |
|               | Gastropoda                          | Snails.   |
|               | Class Amphibia                      | Tadpoles, larval newts.   |
|               | Bony fishes                         | Several species including lake herring, shad, and carp.   |
|               | Detritivores                        | Scavenge food from substrates or through filter-feeding.  |
|               | Order Copepoda                      | Copepods.   |
|               | Ostracoda                           | Seed shrimp.  |
|               | Amphipoda                           | Pontoporeia, scuds, side-swimmers, or freshwater shrimp.  |
|               | Diptera                             | Aquatic larvae of midges and several species of flies.  |
|               | Decapoda                            | Grass shrimp.   |
| <b>3</b>      | <b>Small Carnivores</b>             |   |
|               | Invertebrate Carnivores             |   |
|               | Order Caldocera                     | <i>Leptodora</i> spp., <i>Polyphemus</i> spp.   |
|               | Mysidae                             | <i>Mysis</i> spp.   |
|               | Coleoptera                          | Dytiscidae (predatory diving beetles).  |
|               | Hemiptera                           | Aquatic bugs such as water bugs, water boatmen, and water striders.   |
|               | Odonata                             | Dragonfly and damselfly nymphs and adults.  |
|               | Ephemeroptera                       | Nymphs of some mayflies.  |

## Canadian Tissue Residue Guidelines: Methylmercury

| Trophic Level | Organisms                   | Examples  |
|---------------|-----------------------------|---|
|               | Invertebrate Omnivores      | Many groups of invertebrates include species that feed on both plant and animal matter or that display a wide variety of dietary habits among species. One of the more ubiquitous of these groups are the Trichoptera (or caddis flies). This group includes many omnivorous species, but some species are exclusively grazers, scrapers, suspension feeders, filter feeders, or carnivores.  |
|               | Small Vertebrate Carnivores | Species that feed on zooplankton and benthic invertebrates.   |
|               | Fish species                | Forage fish species including shiners (up to 5 to 6 cm), mudminnows, darters, killifish, sticklebacks, longnose dace, mummichog, suckers, alewife, smelt, sculpin, chub, crappie, bullheads, sunfish, white perch, yellow perch (until age 4), carp.  |
| <b>4</b>      | <b>Large Carnivores</b>     |   |
|               | Predacious fish species     | The larger carnivores, or predators, can feed on the smaller carnivores, but also tend to feed on the larger herbivorous species. In freshwater aquatic ecosystems, the larger carnivores are exclusively fish. Examples of key species include trout (particularly greater than 30 cm), salmon, walleye, northern squawfish (> 30 cm), burbot (> 50 cm), redbreast and chain pickerel, coho salmon, Arctic char (> 20 cm), channel catfish (> 45 cm), gar, and pike. |

Source: US EPA 1995e

**Table 14. Feeding habits and prey trophic levels of representative freshwater amphibian, reptilian, avian, and mammalian species.**

| Species                   | Prey Trophic Level | Feeding Habits  |
|---------------------------|--------------------|---|
| <b>Amphibians</b>         |                    |   |
| Frogs                     | 2                  | Capture insects that may have an aquatic or terrestrial food chain base.  |
| Salamanders               | 2                  | Adults and nymphs are carnivorous.  |
| Mudpuppies                | 2                  | Feed on worms, crayfish, insects, and small fish.   |
| <b>Reptiles</b>           |                    |   |
| Water snakes              | 3                  | Feed primarily on small fish and frogs.   |
| Common snapping turtle    | 3                  | Feed primarily on aquatic invertebrates, small fish, and amphibians.  |
| Eastern painted turtle    | 2                  | Feed on insects, snails, and bits of lily pad.  |
| <b>Birds</b>              |                    |   |
| Herring gull              | 4                  | Herring gulls frequent a wide variety of coastal areas including freshwater and marine. They are highly opportunistic feeders and feed on a wide variety of foods depending on availability including fish, squid, crustaceans, molluscs, worms, insects, small mammals and birds, duck and gull eggs and chicks, and garbage. Freshwater fish usually considered trophic level 3 (up to 23 cm in length, i.e., small freshwater drum, alewife, smelt) comprise a large proportion of herring gull diets in most populations, although some trophic level 4 fish are also taken. About 75% of their diet is fish. |
| Bald eagle                | 4                  | Bald eagles are generally found in coastal areas, lakes, and rivers. They will eat dead or dying fish over most of their range, but also catch live fish swimming near the surface and often eat mammals and birds. In general, bald eagles can be described as opportunistic feeders, eating whatever food source is most plentiful and easy to capture. Bald eagles can capture and carry relatively large prey (i.e., > 60 cm in length) such as chain pickerel, burbot, and lake trout.   |
| Belted kingfisher         | 3                  | Belted kingfishers are typically found along rivers and streams and along lake and pond edges. They are also common on sea coasts and estuaries. Kingfishers feed predominantly on fish, although they sometimes consume large numbers of crayfish. The largest fish prey taken are generally less than 18 cm, the average length being closer to 5 to 8 cm. Freshwater species known to be captured by kingfishers include brook trout, sculpins, blacknose dace, creek chub, common shiner, darters, brook stickleback, redbelly dace, fathead minnow, and suckers.   |
| Osprey                    | 3                  | Osprey are found near fresh or salt water and are almost completely piscivorous. Although, they have been observed on occasion to take other prey including birds, frogs, and crustaceans. Freshwater fish prey include gizzard shad, yellow perch, and salmonids. Most of the fish captured by osprey are between 10 and 35 cm in length.  |
| Ring-billed gull          | 3                  | Although smaller in body size, ring-billed gulls inhabit areas similar to the herring gull. They are generalised and opportunistic foragers, consuming large quantities of fish as well as significant quantities of terrestrial, aquatic, and aerial invertebrates. A study in the Great Lakes found ring-billed gulls to consume mainly smelt, alewives, and sticklebacks. They also consumed earthworms, ephemeroptera (mayflies), homoptera (cicadas), coleoptera (beetles), and diptera (chironomid midges).   |
| Black-crowned night heron | 3                  | Black-crowned night herons nest in a wide variety of freshwater, brackish, and saltwater habitats. They are considered opportunistic general predators, feeding primarily on fish, amphibians, and insects (beetles, flies, and dragonfly nymphs) in freshwater habitats, and on molluscs, spiders, small mammals (e.g., voles), and birds and eggs. Fish species up to 17 cm in length have been taken and include   |

## Canadian Tissue Residue Guidelines: Methylmercury

| Species                        | Prey Trophic Level | Feeding Habits  |
|--------------------------------|--------------------|---|
|                                |                    | whiting, herring, carp, pickerel, suckers, horn-pouts, black bass, perch, gizzard shad, alewife, and eels.  |
| Common tern                    | 3                  | Common terns usually breed on island or coastal beach habitat. They forage in large flocks over schools of small fish, with fish comprising over 90% of their diet. They feed secondarily on crustaceans and insects. In freshwater, common terns have been found to prey on alewife, smelt, bluntnose minnow, common shiners, emerald shiners, and trout-perch.  |
| Forster's tern                 | 3                  | Forster's tern breeds in freshwater and saltwater marshes and on marshy borders of ponds and lakes. Little information exists on the feeding preferences of the Forster's tern, but it is known that it feeds on insects as it flies over marshes and also feeds on fish. Chicks are fed minnows almost exclusively. They also eat frogs, scavenge dead fish, and occasionally take the eggs of the American coot.  |
| Caspian tern                   | 3                  | Caspian terns breed on flat sand or gravel beaches, shell banks, and occasionally marshes in both marine and freshwater. They dive primarily for fish and occasionally take crustaceans. Fish species eaten include alewife, smelt, yellow perch, pumpkinseed, and rock bass. Fish taken are usually 10-12 cm in length.  |
| Black tern                     | 3                  | Black terns breed in shallow freshwater marshes. During the breeding season, they eat mainly insects, including dragonflies, moths, grasshoppers and crickets, beetles, spiders, water scorpions, mayflies, and caddisflies, and smaller amounts of grubs, larvae, small fish, molluscs, and crayfish.  |
| Double-crested cormorant       | 3                  | Double-crested cormorants frequent coasts, bays, estuaries, marine islands, freshwater lakes and their islands, ponds, rivers, sloughs, and swamps. Their diet consists primarily of schooling fish, but may include some small invertebrates. Double-crested cormorants often take small forage fishes such as sticklebacks, sculpins, and burbot, and also take yellow perch, white sucker, and tulibee. Most of the fish taken range from 12-15 cm in length.  |
| Common and American mergansers | 3                  | The American merganser is recognised by some as a subspecies of the common merganser. Common mergansers generally occur in a variety of open freshwater habitats including cold and warm water rivers, ponds, lakes, and inland bays. In riverine habitats, they have been observed to prefer trout and young salmon, and in larger bodies of water will utilise forage fish in large schools. In lakes and rivers, the mergansers were capturing small salmonids and small to medium sculpins (3 to 12 cm in length).  |
| Red-breasted merganser         | 3                  | The habit of the red-breasted merganser is much more marine than that of the common merganser. In the breeding season, it inhabits inland waters as well as coasts and marine islands. In other seasons it inhabits tidewater and inshore marine areas. In freshwater areas, red-breasted mergansers ate salmon eggs, salmonids, sticklebacks, sculpins, schizopods (crustacean), and caddis larvae.  |
| Great blue heron               | 3                  | Great blue herons are found in a variety of freshwater and marine habitats, including freshwater lakes and rivers, brackish marshes, lagoons, mangroves, and coastal wetlands. They are often seen on tidal flats and sandbars, and occasionally forage in wet meadows, pastures, and other terrestrial habitats. Fish are the preferred prey, but they also eat amphibians, reptiles, crustaceans, insects, birds, and mammals. Most of the fish captured represent trophic level 3 (including horn-pouts, shiners, perch, suckers, black bass, herrings, small pickerel) and are less than 25 cm in length. |
| Loons                          | 3                  | Loons feed primarily on small fish at trophic level 3.  |
| Western grebe                  | 3                  | The western grebe diet includes trophic level 2 and 3 aquatic invertebrates and small fish between 3 and 20 cm in length.   |
| Lesser scaup                   | 2                  | Lesser scaup are found on large lakes and bays during the fall and winter, and are common on smaller bodies of water during the spring.   |

## Canadian Tissue Residue Guidelines: Methylmercury

| Species             | Prey Trophic Level | Feeding Habits   |
|---------------------|--------------------|--|
|                     |                    | Most populations consume primarily aquatic invertebrates year round. Common prey include snails, clams, amphipods, midges, chironomids, and leeches. In spring and summer, egg-laying females and ducklings feed at about trophic level 2.   |
| Common goldeneye    | 2                  | During the summer on inland lakes, common goldeneyes tend to consume aquatic invertebrates (75%) and plant material (25%).   |
| Mallard duck        | 2                  | Mallards prefer natural bottomland wetlands and rivers to reservoirs and farm ponds. They feed primarily on seeds of aquatic plants and cultivated grains, although they also consume aquatic invertebrates, particularly during the breeding season. During the summer, 80-90% of their diet is made up of aquatic invertebrates.   |
| Snow goose          | 1                  | Snow geese are almost entirely herbivorous year-round.   |
| Canada goose        | 1                  | Canada geese are almost exclusively vegetarian year-round. They graze on terrestrial and wetland plants.   |
| American black duck | 1                  | The spring and summer diet in freshwater habitat is 80% aquatic plants.  |
| <b>Mammals</b>      |                    |  |
| River otter         | 4                  | River otters are almost exclusively aquatic and are found in food rich coastal areas, the lower portions of streams and rivers, estuaries, nonpolluted waterways, the lakes and tributaries that feed rivers, and areas showing little human impact. They primarily consume fish, but may also consume crustaceans, aquatic insects (e.g., stonefly nymphs, aquatic beetles), amphibians, insects, birds (e.g., ducks), mammals, and turtles. Otters feed primarily on trophic level 3 fish, including suckers, mudminnows, shiners, darters, and carp. They also capture some perch and level 4 walleye, but very few trout and burbot. The average length of fish captured is about 13 cm. |
| Harbour seal        | 4                  | Some freshwater harbour seals are found in Canada. They were found to be depleting older lake trout populations in Québec, although all of the fish were less than 55 cm in length.  |
| Mink                | 3                  | Mink are found associated with aquatic habitats of all kinds, including water ways such as rivers, streams, lakes, and ditches, as well as swamps, marshes, and backwater areas. Mink are generalist and opportunistic feeders, taking whatever prey is locally abundant. The diet of mink consists primarily of prey linked to aquatic ecosystems, including crayfish, frogs, fish, muskrat, and waterfowl. Terrestrial prey include shrews, mice, and voles. Fish species taken include trout, sculpins, blacknose dace, creek chub, suckers, darters, and redbelly dace measuring less than 18 cm in length.  |
| Raccoon             | 2                  | Raccoons are found near virtually every aquatic habitat, particularly in hardwood swamps, mangroves, floodplain forests, and fresh and salt water marshes. Raccoons consume less fish than do river otters or mink. They consume a high amount of fruit and nuts., as well as aquatic invertebrates. Their diet rarely contains trophic level 3 fish.  |

### 13.0 REFERENCES

Abernathy, A.R., and P.M. Cumbie. 1977. Mercury accumulation by largemouth bass (*Micropterus salmoides*) in recently impounded reservoirs. Bull. Environ. Contam. Toxicol. 17: 595-602.

Aceto, M. A.M. Foglizzo, E. Mentasti, G. Sacchero, and C. Sarzanini. 1995. Mercury speciation in biological samples. Intern. J. Environ. Anal. Chem. 60: 1-3.

ADI Nolan Davis INC. 1994. Limnology and fisheries studies at Wreck Cove Reservoir. Nova Scotia Power Inc. Sydney, NS. No. 24-3117-023-1.

AEC (Alberta Environmental Centre). 1989. A five-year study of mercury in fish from a newly formed reservoir (Gleniffer Lake, Alberta). Alberta Environmental Centre, Vegreville, AB. 24 pp., AECV89-R4.

———. 1993. Oldman River Dam: Mercury in fish-Interim report 1993. Alberta Environmental Centre, Vegreville, AB. 13 pp., AECV94-R9.

AEP (Alberta Environmental Protection). 1992. Mercury and its compounds in the environment: their chemistry, toxicology and human health hazards with special reference to Alberta. Draft Criteria Document. Standards Research and Development Branch. Environmental Assessment Division. Edmonton AB, 8 Chapters.

Allard, M., and P.M. Stokes. 1989. Mercury in crayfish species from thirteen Ontario lakes in relation to water chemistry and smallmouth bass (*Micropterus dolomieu*) mercury. Can. J. Fish. Aquat. Sci. 46: 1040-1046.

Amyot, M.G. Mierle, D.R.S. Lean, and D.J. McQueen. 1994. Sunlight-induced formation of dissolved gaseous mercury in lake waters. Environ. Sci. Technol. 28(13): 2366-2371.

Anderson, A. 1979. Mercury in soils. In: The biogeochemistry of mercury in the environment, J.O. Nriagu, ed. Elsevier/North-Holland Biomedical Press, New York. pp. 79-112.

Anderson, M.R., D.A. Scruton, U.P. Williams, and J.F. Payne. 1995. Mercury in fish in the Smallwood Reservoir, Labrador, twenty-one years after impoundment. Water Air Soil Pollut. 80: 927-930.

Andersson, I., H. Parkman, and A. Jernelov. 1990. The role of sediments as sink or source for environmental contaminants- a case study of mercury and chlorinated organic compounds. Limnologica 20: 347-359.

Armstrong, J.E. 1946. Geology and mineral deposits of northern British Columbia west of the Rocky Mountains. Dept. of Mines and Resources. Geological Survey Bulletin No. 5, 46 pp.

Aulerich, R.J., R.K. Ringer, and S. Iwamoto. 1974. Effects of dietary mercury on mink. Arch. Environ. Contam. Toxicol. 2: 43-51.

Back, R.C., and C.J. Watras. 1995. Mercury in zooplankton of northern Wisconsin lakes: taxonomic and site-specific trends. Water Air Soil Pollut. 80: 931-938.

## Canadian Tissue Residue Guidelines: Methylmercury

Bailey, H.S. 1985. 1983 Toxic Chemical Survey: a survey of nine impoundments in the Saint John River Basin. Environment Canada. Moncton, N.B. pp. 1-40.

Barr, J.F. 1986. Population dynamics of the common loon (*Gavia immer*) associated with mercury-contaminated waters in North-western Ontario. Canadian Wildlife Service. Occasional paper 56. 23 pp.

Bélanger, D., and N. Larivière. 1997. Développement et validation de biomarqueurs d'effets physiopathologiques précoces chez certains piscivores relativement à leur exposition au méthylmercure. Rapport final. Faculté de médecine vétérinaire, Université de Montréal. 105 pp. and Annexes.

Betti, C., T. Davini, and R. Barale. 1992. Genotoxic activity of methyl mercury chloride and dimethyl mercury in human lymphocytes. *Mutat. Res.* 281(4): 255-260.

Bishop, C.A., M.D. Koster, A.A. Chek, D.J.T. Hussell, and K. Jock. 1995. Chlorinated hydrocarbons and mercury in sediments, red-winged blackbirds (*Agelaius phoeniceus*) and tree swallows (*Tachycineta bicolor*) from wetlands in the Great Lakes-St. Lawrence River Basin. *Environ. Toxicol. Chem.* 14: 491-501.

Bleavins, M.R., and R.J. Aulerich. 1981. Feed consumption and food passage in mink (*Mustela vison*) and European ferrets (*Mustela putorius furo*). *Lab. Animal Sci.* 31: 268-269.

Bloom, N.S. 1989. Determination of picogram levels of methylmercury by aqueous phase ethylation, followed by cryogenic gas chromatography with cold vapour atomic fluorescence detection. *Can. J. Fish. Aquat. Sci.* 46: 1131-1140.

———. 1992. On the chemical form of mercury in edible fish and marine invertebrate tissue. *Can. J. Fish. Aquat. Sci.* 49: 1010-1017.

Bloom, N.S., and C.J. Watras. 1989. Observations of methylmercury in precipitation. *Sci. Total Environ.* 87/88: 199-207.

Blum, J.E., and R. Bartha. 1980. Effect of salinity on methylation of mercury. *Bull. Environ. Contam. Toxicol.* 25: 404-408.

Bodaly, R.A., and R.E. Hecky. 1979. Post-impoundment increases in fish mercury levels in the Southern Indian Lake reservoir, Manitoba. *Can. Fish. Mar. Serv. MS Rep.* 1531.

Bodaly, R.A., R.E. Hecky, and R.J.P. Fudge. 1984. Increases in fish mercury levels in lakes flooded by the Churchill River Diversion, Northern Manitoba. *Can. J. Fish. Aquat. Sci.* 41: 682-691.

Bodaly, R.A., J.W.M. Rudd, R.J.P. Fudge, and C.A. Kelly. 1993. Mercury concentrations in fish related to size of remote Canadian Shield lakes. *Can. J. Fish. Aquat. Sci.* 50: 980-987.

Bonin, J., J.-L. Desgranges, C.A. Bishop, J. Rodrigue, A. Gendron, and J.E. Elliott. 1995. Comparative study of contaminants in the mudpuppy (Amphibia) and the common snapping turtle (Reptilia), St. Lawrence River, Canada. *Arch. Environ. Contam.* 28: 184-194.

## Canadian Tissue Residue Guidelines: Methylmercury

Borg, K., K. Erne, E. Hanco, and H. Wanntorp. 1970. Experimental secondary methylmercury poisoning in the Goshawk (*Accipiter g. gentilis* L.). *Environ. Pollut.* 1: 91-104.

Borgmann, U., and D. M. Whittle. 1991. Contaminant concentration trends in Lake Ontario lake trout (*Salvelinus Namaycush*). *J. Great Lakes Res.* 17: 368-381.

Bornhausen, M., H.R. Musch, and H. Greim. 1980. Operant behavior performance changes in rats after prenatal methylmercury exposure. *Toxicol. Appl. Pharmacol.* 56: 305-310.

Boudou, A., and F. Ribeyre. 1985. Experimental study of trophic contamination of *Salmo gairdneri* by two mercury compounds - HgCl<sub>2</sub> and CH<sub>3</sub>HgCl - analysis at the organism and organ levels. *Water Air Soil Pollut.* 26: 137-148.

Boudou, A., M. Delnomdedieu, D. Georgescauld, F. Ribeyre, and E. Saouter. 1991. Fundamental roles of biological barriers in mercury accumulation and transfer in freshwater ecosystems (analysis at organism, organ, cell and molecular levels). *Water Air Soil Pollut.* 56: 807-822.

Braune, B.M. 1987a. Mercury accumulation in relation to size and age of Atlantic herring (*Clupea harengus harengus*) from the Southwestern Bay of Fundy, Canada. *Arch. Environ. Contam. Toxicol.* 16: 311-320.

———. 1987b. Comparison of total mercury levels in relation to diet and molt for nine species of marine birds. *Arch. Environ. Contam. Toxicol.* 16: 217-224.

Braune, B.M., and D.E. Gaskin. 1987. Mercury levels in Bonaparte's gulls (*Larus philadelphia*) during autumn molt in the Quoddy Region, New Brunswick, Canada. *Arch. Environ. Contam. Toxicol.* 16: 539-549.

Bruce, W.J., and K.D. Spencer. 1978. Mercury levels in Labrador fish, 1977-78. Technical Report #111. Department of Fisheries and Oceans. St. John's, NF.

Bruce, W.J., K.D. Spencer, and E. Arsenault. 1979. Mercury content data for Labrador fishes, 1977-78. Fisheries and Marine Service Data Report 142.

Burbacher, T.M., C. Monnett, L.S. Grant, and N.K. Mottet. 1984. Methylmercury exposure and reproductive dysfunction in the nonhuman primate. *Toxicol. Appl. Pharmacol.* 75: 18-24.

Cabana, G., and J.B. Rasmussen. 1994. Modelling food chain structure and contaminant bioaccumulation using stable nitrogen isotopes. *Nature* 372: 255-257.

Cabana, G., A. Tremblay, J. Kalff, and J.B. Rasmussen. 1994. Pelagic food chain structure in Ontario lakes: a determinant of mercury levels in lake trout (*Salvelinus namaycush*). *Can. J. Fish. Aquat. Sci.* 51: 381-389.

Canada-Manitoba Mercury Agreement. 1987. Summary Report. Canada-Manitoba agreement on the study and monitoring of mercury in the Churchill River Diversion. Joint report by Environment Canada and Environment and Workplace Safety and Health of Manitoba. Winnipeg, Manitoba. 77 pp. + Maps.

Caurant, F., M. Navarro, and J.-C. Amiard. 1996. Mercury in pilot whales: possible limits to the



detoxification process. *Sci. Tot. Environ.* 186: 95-104.

Cavalli, S., and N. Cardellicchio. 1995. Direct determination of seleno-amino acids in biological tissues by anion-exchange separation and electrochemical detection. *J. Chromatog.* 706(A): 429-436.

CCME (Canadian Council of Ministers of The Environment). 1993. Protocols for deriving water quality guidelines for the protection of agricultural water uses. Appendix XV. In: *Canadian Water Quality Guidelines*, Canadian Council of Ministers of the Environment (CCREM), ed. CCREM, Ottawa, Ontario.

———. 1998. A protocol for the derivation of Canadian tissue residue guidelines for the protection of wildlife that consume aquatic biota. CCME, Winnipeg, MB and Environment Canada, Science Policy and Environmental Quality Branch, Ottawa, ON.

CEPA (Canadian Environmental Protection Act), S.C. 1999, c.33.

Chamberland, G., D. Bélanger, A. Dallaire, J.S. Blais, L. Vermette, and N. Larivière. 1996. Urinary protein excretion of semidomesticated mink in a chronic methylmercury study. *J. Toxicol. Environ. Health* 47: 285-297.

Chang, L.W. 1979. Pathological effects of mercury poisoning. In: *The biogeochemistry of mercury in the environment*, J.O. Nriagu, ed. Elsevier/North-Holland Biomedical Press. New York. pp. 519-580.

Chang, L.W., and Z. Annau. 1984. Developmental neuropathology and behavioral teratology of methylmercury. In: *Neurobehavioral Teratology*, J. Yanai, ed. Elsevier, Amsterdam. pp. 405-432.

Chang, L.W., S. Yamachghi, and A.W.J. Dudley. 1974. Neurological changes in cats following long-term diet of mercury contaminated tuna. *Acta Neuropathol. (Berl)* 27: 171-176.

Charbonneau, S.M., I.C. Munro, E.A. Nera, and others. 1976. Chronic toxicity of methylmercury in the adult cat. *Toxicology*. 5: 337-349.

Clarkson, T.W. 1994. The toxicology of mercury and its compounds. Chapter VIII.1. In: *Mercury pollution: Integration and synthesis*, C.J. Watras and J.W. Huckabee, eds. Lewis Publishers, USA. pp. 631-641.

Codex Alimentarius. 1998. General Standard for Contaminants in Foods. Food and Safety Information Centre, DLO-State Institute for Quality Control of Agricultural Products. Wageningen, Netherlands.

Cope, W.G., J.G. Weiner, and R.G. Rada. 1990. Mercury accumulation in yellow perch in Wisconsin seepage lakes: relation to lake characteristics. *Environ. Toxicol. Chem.* 9: 931-940.

Coquery, M., and P.M. Welbourn. 1995. The relationship between metal concentration and organic matter in sediments and metal concentration in the aquatic macrophyte *Eriocaulon septangulare*. *Water Res.* 29(9): 2094-2102.

Cossa, D., and J.-G. Rondeau. 1985. Seasonal, geographical and size-induced variability in

mercury content of *Mytilus edulis* in an estuarine environment: a re-assessment of mercury pollution level in the Estuary and Gulf of St. Lawrence. *Mar. Biol.* 88: 43-49.

Cossa, D., M. Coquery, C. Gobeil, and J.M. Martin. 1996. Mercury fluxes at the ocean margins. In: *Regional and global mercury cycles: Sources, fluxes and mass balances*, W. Baeyens, ed. Kluwer Publ. pp. 229.

Costa, M., N.T. Christie, O. Cantoni, J.T. Zelikoff, X.W. Wang, and T.G. Rossman. 1991. DNA damage by mercury compounds: An overview. In: *Advances in mercury toxicity*, T. Suzuki, N. Imura, and T.W. Clarkson, eds. Plenum Press, New York, NY. pp. 255-273.

Cox, M.F., H.W. Holm, H.J. Kania, and R.L. Knight. 1975. Methylmercury and total mercury in selected stream biota. In: *Trace substances in environmental health, IX.*, D.D. Hemphill, ed. Univ. Missouri, Columbia, MO. pp. 151-155.

Craig, P.J., and P.D. Bartlett. 1978. The role of hydrogen sulphide in environmental transport of mercury. *Nature (Lond.)* 275: 635-637.

Crowder, A. 1991. Acidification, metals and macrophytes. *Environ. Pollut.* 71: 171-203.

Cuvin-Aralar, M.L.A., and R.W. Furness. 1991. Mercury and selenium interaction: a review. *Ecotoxicol. Environ. Safety.* 21: 348-364.

Czuba, M., and D.C. Mortimer. 1980. Stability of methyl-mercury and inorganic mercury in aquatic plants (*Elodea densa*). *Can. J. Bot.* 58: 316-320.

Davis, R.B., D.S. Anderson, and F. Berge. 1985. Palaeolimnological evidence that lake acidification is accompanied by loss of organic matter. *Nature (Lond.)* 316: 436-438.

De Flora, S., C. Bennicelli, and M. Bagnasco. 1994. Genotoxicity of Mercury Compounds. A Review. *Mutation Research*, Elsevier Science B.V. 317: pp. 57-79.

Desgranges, J.-L., J. Rodrigue, B. Tardif, and M. Laperle. 1998. Breeding success of ospreys under high seasonal methylmercury exposure, Ch. 6.2. In: *Mercury in natural environments and hydroelectric reservoirs of northern Québec (Canada)*. M. Lucotte, ed. Hydro-Québec, Environment Canada, Montreal. pp. 274-280.

Dietz, R., C.O. Nielsen, M.M. Hansen, and C.T. Hansen. 1990. Organic mercury in Greenland birds and mammals. *Sci. Tot. Environ.* 95: 41-51.

D'Itri, F.M. 1972. *The environmental mercury problem*. CRC Press. Inc., Boca Raton, Fla.

———. 1991. Mercury contamination- what we have learned since minamata. *Environ. Mon. Assess.* 19: 165-182.

Douglas, J. 1991. Mercury in the environment. *EPRI Journal*. December: 4-11.

Driscoll, C.T., C. Yan, C.L. Schofield, R.K. Munson, and J.G. Holsapple. 1994. The mercury cycle and fish in the Adirondack lakes. *Environ. Sci. Technol.* 28(3): 136A-143A.

Drummond, R.A., G.F. Olson, and A.R. Batterman. 1974. Cough response and uptake of mercury by brook trout, *Salvelinus fontinalis*, exposed to mercuric compounds at different

## Canadian Tissue Residue Guidelines: Methylmercury

hydrogen-ion concentrations. *Trans. Amer. Fish. Soc.* 103: 244-249.

Eaton, R.D.P., D.C. Secord, and P. Hewitt. 1980. An experimental assessment of the toxic potential of mercury in ringed-seal liver for adult laboratory cats. *Toxicol. Appl. Pharmacol.* 5: 514-521.

Eccles, C.U., and Z. Annau. 1987. Prenatal exposure to methylmercury. In: *The toxicity of methylmercury*, C.U. Eccles and Z. Annau, eds. Johns Hopkins University Press, Baltimore, MD. pp. 114-130.

Eisler, R. 1987. Mercury hazards to fish, wildlife, and invertebrates: a synoptic review. *Biol. Report 85(1.10)*, Contaminant Hazard Reviews No. 10, U.S. Dept. of Interior, Fish & Wildlife Service. Washington DC. 90 pp.

EPRI (Electric Power Research Institute). 1996. Protocol for estimating historic atmospheric mercury deposition. TR-106768. Palo. Alto, CA. 49 pp.

Elliott, J.E., A.M. Scheuhammer, F.A. Leighton, and P.A. Pearce. 1992. Heavy metal and metallothionein concentrations in Atlantic Canadian seabirds. *Arch. Environ. Contam. Toxicol.* 22: 63-73.

Elliott, J.E., R.J. Norstrom, and G.E.J. Smith. 1996. Patterns, trends, and toxicological significance of chlorinated hydrocarbon and mercury contaminants in bald eagle eggs from the Pacific Coast of Canada, 1990-1994. *Arch. Environ. Contam. Toxicol.* 31: 354-367.

Engstrom, D.R., and E.B. Swain. 1997. Recent declines in atmospheric mercury deposition in the upper midwest. *Environ. Sci. Technol.* 31: 960-967.

Engstrom, D.R., E.B. Swain, T.A. Henning, M.E. Brigham, and P.L. Brezonik. 1994. Atmospheric mercury deposition to lakes and watersheds. A quantitative reconstruction from multiple sediment cores. In: *Environmental Chemistry of Lakes and Reservoirs*, L.A. Baker, ed. *Advances in Chemistry Series 237*. pp. 33-66.

Environment Canada. 1996. Canadian soil quality guidelines for inorganic mercury: Environmental Supporting Document- Final Draft. December 1996. Science Policy and Environmental Quality Branch, Guidelines Division, Ottawa. Unpub.

———. 1997. Canadian sediment quality guidelines for mercury. Supporting document. Environmental Conservation Service, Ecosystem Science Directorate, Science Policy and Environmental Quality Branch, Guidelines and Standards Division. Ottawa 103 pp. + Append.

Environment Canada and US EPA (Environmental Protection Agency). 1997. Canada -- United States strategy for the virtual elimination of persistent toxic substances in the Great Lakes. The Great Lakes binational toxics strategy. Prepared for website publication by P. Pranckevicius. (<http://www.epa.gov/glnpo/p2/bns.html>).

EPMAP (Expert Panel on Mercury Atmospheric Processes). 1994. Mercury atmospheric processes: A synthesis report. Workshop proceedings Sept. 1994. Tampa FL. EPRI Report No. TR-104214. 23 pp.

Evans, R.D. 1995. Mercury levels in otters. Proceedings of the 1995 Canadian mercury network

## Canadian Tissue Residue Guidelines: Methylmercury

workshop, T. Brydges, A. Perez-Fuentetaja, and D.J. McQueen, eds. York University, Toronto. pp. 6-8.

Evers, D.C., J.D. Kaplan, M.W. Meyer, P.S. Reaman, W.E. Braselton, A. Major, N. Burgess, and A.M. Scheuhammer. 1998. Geographic trend in mercury measured in common loon feathers and blood. *Environ. Toxicol. Chem.* 17: 173-183.

Faust, B.C. 1992. The octanol/water distribution coefficients of methylmercuric species: the role of aqueous-phase chemical speciation. *Environ. Toxicol. Chem.* 11: 1373-1376.

Fehr, F., and C.A.R. Dennis. 1975. Investigations into the use and hazard of mercury in Saskatchewan. Prairie Institute of Environmental Health. Regina, Saskatchewan.

Fimreite, N. 1971. Effects of methylmercury on ring-necked pheasants, with special reference to reproduction. Occasional Paper No. 9. Canadian Wildlife Service, Ottawa, ON.

Fimreite, N., and L.M. Reynolds. 1973. Mercury contamination of fish in northwestern Ontario. *J. Wildl. Manage.* 37: 62-68.

Finley, M.T., and R.C. Stendell. 1978. Survival and reproductive success of black ducks fed methyl mercury. *Environ. Pollut.* 16(1): 51-64.

Fisheries Act, R.S. 1985, c. F-14.

Fiskesjo, G. 1979. Two organic mercury compounds tested for mutagenicity in mammalian cells by use of the cell line V 79-4. *Hereditas.* 90: 103-109.

Fitzgerald, W.F. 1986. Cycling of mercury between the atmosphere and oceans. In: The role of air-sea exchange in geochemical cycling, P. Buat-Menard, ed. NATO Advanced Science Institutes Series, Reidel, Dordrecht, pp. 363-408.

Fitzgerald, W.F., and C.J. Watras. 1989. Mercury in the surficial waters of rural Wisconsin lakes. *Sci. Total. Environ.* 87/88: 223-232

Foley, R.E., S.J. Jackling, R.J. Sloan, and M.K. Brown. 1988. Organochlorine and mercury residues in wild mink and otter: comparison with fish. *Environ. Toxicol. Chem.* 7: 363-374.

Foster, T.J. 1987. The genetics and biochemistry of mercury resistance. *CRC. Crit. Rev. Microbiol.* 15: 117-140.

Frank, R., H. Lumsden, J.F. Barr, and H.E. Braun. 1983. Residues of organochlorine and mercury in eggs and tissues of healthy and emaciated common loons (*Gavia immer*), Ontario, Canada, 1968-80. *Arch. Environ. Contam. Toxicol.* 12: 641-654.

Freeman, H.C., and D.A. Horne. 1972. Total mercury and methylmercury content of the American eel. *J. Fish. Res. Bd. Can.* 30: 454-456.

Friedmann, A.S., M.C. Watzin, T. Brinck-Johnsen, and J.C. Leiter. 1996. Low levels of dietary methylmercury inhibit growth and gonadal development in juvenile walleye (*Stizostedion vitreum*). *Aquat. Toxicol.* 35: 265-278.

Friske, P.W.B., and W.B. Coker. 1995. The importance of geological controls on the natural

distribution of mercury in lake and stream sediments across Canada. *Water Air Soil Pollut.* 80: 1047-1051.

Furukawa, K., T. Suzuki, and K. Tonomura. 1969. Decomposition of organic mercurial compounds by mercury-resistant bacteria. *Agric. Biol. Chem.* 33: 128-130.

Furutani, A., and J.W.M. Rudd. 1980. Measurement of mercury methylation in lake water and sediment samples. *Appl. Environ. Microbiol.* 40(4): 770-776.

Gagnon, C., and N.S. Fisher. 1997. Bioavailability of sediment-bound methyl and inorganic mercury to a marine bivalve. *Environ. Sci. Technol.* 31: 993-998.

Ganther, H.E., C. Goudie, M.L. Sunde, M.J. Kipecky, P. Wagner, S.H. Oh, and W.G. Hoekstra. 1972. Selenium relation to decreased toxicity of methyl mercury added to diets containing tuna. *Science* 175: 1122-1124.

Gilbertson, M. 1974. Seasonal changes in organochlorine compounds and mercury in common terns of Hamilton harbour, Ontario. *Bull. Environ. Contam. Toxicol.* 12: 726-732.

Gill, G.A., and K.W. Bruland. 1990. Mercury speciation in surface freshwater systems in California and other areas. *Environ. Sci. Technol.* 24: 1392-1400.

Gill, G.A., and W.F. Fitzgerald. 1985. Mercury sampling of open ocean waters at the picomolar level. *Deep-Sea Res.* 32: 287-297.

———. 1987. Picomolar mercury measurements in seawater and other materials using stannous chloride reduction and two-stage gold amalgamation with gas phase detection. *Mar. Chem.* 20: 227-243.

Gilmour, C.C., and G.S. Riedel. 1995. Measurement of Hg methylation in sediments using high specific-activity hg and ambient incubation. *Water Air Soil Pollut.* 80: 747-756.

Gilmour, C.C., E.A. Henry, and R. Mitchell. 1992. Sulfate stimulation of mercury methylation in freshwater sediments. *Environ. Sci. Technol.* 26: 2281-2286.

Grey, B.J., S.M. Harbicht, and G.R. Stephens. 1995. Mercury in fish from rivers and lakes in southwestern Northwest Territories. Ottawa, Canada. Department of Indian Affairs and Northern Development. pp. 1-61.

Grieb, T.M., C.T. Driscoll, S.P. Gloss, C.L. Schofield, G.L. Bowie, and D.B. Porcella. 1990. Factors affecting mercury accumulation in fish in the Upper Michigan Peninsula. *Environ. Toxicol. Chem.* 9: 919-930.

Gunderson, V.M., K.S. Grant, J.F. Fagan, and N.K. Mottet. 1986. The effect of low-level prenatal methylmercury exposure on visual recognition memory of infant crab-eating macaques. *Child Dev.* 57: 1076-1083.

Haines, M.L., K. Brydges, M.J. Macdonald, S.L. Smith, and D.D. Macdonald. 1994. A review of environmental quality criteria and guidelines for priority substances in the Fraser River basin: Supporting documentation. Report prepared for Environment Canada. Pacific and Yukon Region. North Vancouver, British Columbia.

## Canadian Tissue Residue Guidelines: Methylmercury

Håkanson, L., A. Nilsson, and T. Andersson. 1988. Mercury in fish in Swedish lakes. *Environ. Pollut.* 49: 145-162.

Halbach, S. 1985. The octanol/water distribution of mercury compounds. *Arch. Toxicol.* 57: 139-141.

Halbrook, R.S., J.H. Jenkins, P.B. Bush, and N.D. Seabolt. 1994. Sublethal concentrations of mercury in river otters: monitoring environmental contamination. *Arch. Environ. Contam. Toxicol.* 27: 306-310.

Hammer, U.T., A.T. Merkowsky, and P.M. Huang. 1988. Effects of oxygen concentrations on release of mercury from sediments and accumulation by *Ceratophyllum demersum* and *Anodonta grandis*. *Arch. Environ. Contam. Toxicol.* 17: 257-262.

Harris, R.C., and W.J. Snodgrass. 1993. Bioenergetic simulations of mercury uptake and retention in walleye (*Stizostedion vitreum*) and yellow perch (*Perca flavescens*). *Water Pollut. Res. J. Can.* 28(1): 217-236.

Headon, C.M., R.J. Hall, and G. Mierle. 1996. Dynamics of radiolabelled methylmercury in crayfish (*Orconectes virilis*). *Can. J. Fish. Aquat. Sci.* 53: 2862-2869.

Hecky, R.E., D.J. Ramsey, R.A. Bodaly, and N.E. Strange. 1991. Increased methylmercury contamination in fish in newly formed freshwater reservoirs. In: *Advances in mercury toxicology*, T.F.W. Clarkson, T. Suzuki, and A. Imura, eds. Plenum Press, New York. pp. 33-52.

Heddle, J.R., and W.R. Bruce. 1977. Comparison of the micronucleus and sperm assay for mutagenicity with the carcinogenic activities of 61 different agents. In: *Origins of Human Cancer*, Vol. 4., H.H. Hiatt, J.D. Watson, J.A. Winsten, eds. Cold Spring Harbor Conferences.

Heinz, G. 1974. Effects of low dietary levels of methyl mercury on mallard reproduction. *Bull. Environ. Contam. Toxicol.* 11: 386-392.

———. 1975. Effects of methylmercury on approach and avoidance behavior of mallard ducklings. *Bull. Environ. Contam. Toxicol.* 13: 554-564.

———. 1976a. Methyl mercury: second year feeding effects on mallard reproduction and duckling behavior. *J. Wildl. Manage.* 40: 82-90.

———. 1976b. Methyl mercury: second generation reproductive and behavioral effects on mallards. *J. Wildl. Manage.* 40(4): 710-715.

Heinz, G.H. 1979. Methylmercury: reproductive and behavioural effects on three generations of mallard ducks. *J. Wildl. Manage.* 43: 394-400.

Heinz, G.H., and D.J. Hoffman. 1998. Methylmercury chloride and selenomethionine interactions on health and reproduction in mallards. *Environ. Toxicol. Chem.* 17: 139-145.

Hendzel, M.R., and D.M. Jamieson. 1976. Determination of mercury in fish. *Anal. Chem.* 48: 926-928.

Hildebrand, S.G., R.H. Strand, and J.W. Huckabee. 1980. Mercury accumulation in fish and

invertebrates of the North Fork Holston River, Virginia and Tennessee. *J. Environ. Qual.* 9: 393-400.

Hirano, M., K. Mitsumori, K. Maita, and Y. Shirasu. 1986. Further carcinogenicity study on methylmercury chloride in ICR mice. *Jpn. J. Vet. Sci.* 48(1): 127-135.

Hoffman, D.J., and G.H. Heinz. 1998. Effects of mercury and selenium on glutathione metabolism and oxidative stress in mallard ducks. *Environ. Toxicol. Chem.* 17: 161-166.

Hoffman, D.J., and J.M. Moore. 1979. Teratogenic effects of external egg applications of methyl mercury in the mallard, *Anas platyrhynchos*. *Teratology* 20: 453-462.

Haupt, K.A., L.A. Essick, E.B. Shaw, D.K. Alo, J.E. Gilmartin, W.H. Gutenmann, C.B. Littman, and D.J. Lisk. 1988. A tuna fish diet influences cat behavior. *J. Toxicol. Environ. Health* 24: 161-172.

Huckabee, J.W., J.W. Elwood, and S.G. Hildebrand. 1979. Accumulation of mercury in freshwater biota." In: *The Biogeochemistry of Mercury in the Environment*, J.O. Nriagu, ed. Elsevier/North-Holland Biomedical Press, pp. 276-302.

Hudson, R.J.M., S.A. Gherini, C.J. Watras, and D.B. Porcella. 1994. A mechanistic model of the biogeochemical cycle of mercury in lakes. In: *Mercury pollution: Integration and synthesis*. C.J. Watras and J.W. Huckabee, eds. Lewis Publishers. pp. 473-526.

Hurley, J.P., C.J. Watras, and N.S. Bloom. 1989. Biogeochemistry of mercury in aquatic ecosystems. Aqueous speciation of mercury: seston component. *Semiannual Report to EPRI*. Madison WI. 10 pp.+ Figures.

———. 1991. Mercury cycling in a northern Wisconsin seepage lake: The role of particulate matter in vertical transport. *Water Air Soil Pollut.* 56: 543-551.

Hurley, J.P., D.P. Krabbenhoft, C.L. Babiartz, and A.W. Andren. 1994. Cycling of mercury across the sediment-water interface in seepage lakes. In: *Environmental Chemistry of Lakes and Reservoirs*, L.A. Baker, ed. American Chemical Society, Washington DC. pp. 425-449.

Hurley, J.P., J.M. Benoit, C.L. Babiartz, M.M. Shafer, A.W. Andren, J.R. Sullivan, R. Hammond, and D.A. Webb. 1995. Influences of watershed characteristics on mercury levels in Wisconsin rivers. *Environ. Sci. Technol.* 29: 1867-1875.

HWC (Health and Welfare Canada). 1990. Food chemical contaminants - Assessing health risks. Health Protection Branch. Ottawa, Canada. 2 pp.

Hydro-Québec and Groupe-Conseil Génivar Inc. 1996. Evolution des teneurs en mercure des poissons du complexe La Grande (1978-1994), Annexe 2. Hydro-Québec. 509 pp.

———. 1997. Summary Report - Evolution of fish mercury levels at the La Grande complex, Québec (1978-1994). Hydro-Québec. 143 pp.+ Append./maps.

IJC (International Joint Commission). 1990. Production, usage and atmosphere emissions of 14 priority toxic chemicals. Appendix 2 to the Proceedings of the Workshop on Great Lakes Atmospheric Deposition, 29-31 October 1986.

Jackson, T.A. 1987. Methylation, demethylation, and bio-accumulation of mercury in lakes and reservoirs of northern Manitoba, with particular reference to effects of environmental changes caused by the Churchill-Nelson River diversion. Can. Data Rep. Fish. Aquat. Sci. 628.

———. 1988a. The mercury problem in recently formed reservoirs of northern Manitoba (Canada): effects of impoundment and other factors on the production of methyl mercury by microorganisms in sediments. Can. J. Fish. Aquat. Sci. 45: 97-121.

———. 1988b. Accumulation of mercury by plankton and benthic invertebrates in riverine lakes of northern Manitoba (Canada): Importance of regionally and seasonally varying environmental factors. Can. J. Fish. Aquat. Sci. 45: 1744-1757.

———. 1989. The influence of clay minerals, oxides, and humic matter on the methylation and demethylation of mercury by micro-organisms in freshwater sediments. Appl. Organo. Chem. 3: 1-30.

———. 1993. Effects of environmental factors and primary production on the distribution and methylation of mercury in a chain of highly eutrophic riverine lakes. Water Poll. Res. J. Can. 28: 177-216.

Jana, S. 1988. Accumulation of Hg and Cr by three aquatic species and subsequent changes in several physiological and biochemical plant parameters. Water Air Soil Pollut. 38: 105-109.

Jaques, A.P. 1987. Summary of emissions of antimony, arsenic, cadmium, chromium, copper, lead, manganese, mercury and nickel in Canada. Environment Canada. May 1987.

Jensen, S., and A. Jernelöv. 1969. Biological methylation of mercury in aquatic organisms. Nature 223: 753-754.

Jernelöv, A., and C. Ramel. 1994. Mercury in the environment. Synopsis of the SCOPE meeting on mercury held at the Royal Academy of Science, Stockholm, 28-30 October 1993. Ambio 23: 166.

Johnson, M.G. 1987. Trace element loadings to sediments of fourteen Ontario lakes and correlations with concentrations in fish. Can. J. Fish. Aquat. Sci. 44: 3-13.

Johnston, T.A., R.A. Bodaly, and J.A. Mathias. 1991. Predicting fish mercury levels from physical characteristics of boreal reservoirs. Can. J. Fish. Aquat. Sci. 48: 1468-1475.

Keller, A.E., and S.G. Zam. 1991. The acute toxicity of selected metals to the freshwater mussel, *Anodonta imbecilis*. Environ. Toxicol. Chem. 10: 539-546.

Kelly, C.A., J.W.M. Rudd, V.L. St.Louis, and A. Heyes. 1995. Is total mercury concentration a good predictor of methyl mercury concentration in aquatic systems? Water Air Soil Pollut. 80: 715-724.

Kent, R.A., P.-Y. Caux, R. Allan, R. Post, and J. Parks. 1998. A Canada-wide GIS analysis of methylmercury in fish - exploring relative risks to wildlife and human health. In: W. Pilgrim, N. Burgess, and M-F. Giguère (Eds.). Mercury in Eastern Canada and the Northeast States. Proceedings of the Conference held in Fredericton, New Brunswick, Sept. 21-23, 1998



## Canadian Tissue Residue Guidelines: Methylmercury

- Kidd, K.A., J.E. Eamer, and D.C.G. Muir. 1993. Spatial variability of chlorinated bornanes (toxaphene) in fishes from Yukon lakes. *Chemo.* 27: 1975-1986.
- King, D.G., and I.M. Davies. 1987. Laboratory and field studies of the accumulation of inorganic mercury by the mussel *Mytilus edulis* (L.) *Mar. Pollut. Bull.* 18(1): 40-45
- Koster, M.D., D.P. Ryckman, D.V.C. Weseloh, and J. Struger. 1996. Mercury levels in great lakes herring gull (*Larus argentatus*) eggs, 1972-1992. *Environ. Pollut.* 93: 261-270.
- Krabbenhoft, D.P., and D.A. Rickert. 1995. Mercury contamination of aquatic ecosystems. Fact Sheet FS-216-95, US Department of the Interior, US Geological Survey.
- Krabbenhoft, D.P., J.M. Benoit, C.L. Babiarz, J.P. Hurley, and A.W. Andren. 1995. Mercury cycling in the Allequash Creek watershed, northern Wisconsin. *Water Air Soil Pollut.* 80: 425-433.
- Krenkel, P.A. 1974. Mercury, environmental considerations, Part II. *CRC Crit. Rev. Environ. Control.* 4: 251-339.
- Kristensen, P. 1982. Time-dependent variation of mercury in a stream sediment and the effect upon mercury content in *Gammarus pulex* (L.). *Water Res.* 16: 759-764.
- Kucera, E. 1983. Mink and otter as indicators of mercury in Manitoba waters. *Can. J. Zool.* 61: 2250-2256.
- Laliberté, D. 1996. Teneurs en mercure et en composés organochlorés dans les poissons capturés au lac Aylmer en 1995. Ministère de l'Environnement et de la Faune du Québec. Direction des écosystèmes aquatiques, Québec. 32 pp.
- Landers, D H., C. Gubala, M. Verta, M. Lucotte, K. Johansson, T. Vlasova, and W.L. Lockhart. 1998. Using lake sediment mercury flux ratios to evaluate the regional and continental dimensions of mercury deposition in Arctic and boreal ecosystems. *Atmospheric Environ.* 32 (5): 919-928.
- Langlois, C., R. Langis, and M. Perusse. 1995. Mercury contamination in northern Québec environment and wildlife. *Water Air Soil Pollut.* 80: 1021-1024.
- Laperle, M., J. Sbeghen, and D. Messier. 1998. Assessment of the ecotoxic risk of methylmercury exposure in mink (*Mustela vison*) inhabiting northern Québec, Ch. 6.1. In: *Mercury in natural environments and hydroelectric reservoirs of northern Québec (Canada)*, M. Lucotte, ed. Hydro-Québec, Environment Canada, Montreal. pp. 262-273.
- Lee, Y.H. 1987. Determination of methyl- and ethylmercury in natural waters at sub-nanogram per liter using SCF-adsorbent preconcentration procedure. *Intern. J. Environ. Anal. Chem.* 29: 263-276.
- Lee, Y-H., and H. Hultberg. 1990. Methyl mercury in some Swedish surface waters. *Environ. Toxicol. Chem.* 9: 833-841.
- Lee, Y-H., and C. Iverfeldt. 1991. Measurement of methylmercury and mercury run-off, lake and rain waters. *Water Air Soil Pollut.* 56: 309-321.

Lee, Y-H., H. Hultberg, and I. Andersson. 1985. Catalytic effect of various metal ions on the methylation of mercury in the presence of humic substances. *Water Air Soil Pollut.* 25: 391-400.

Leermakers, M., M. Elskens, S. Panutrakul, F. Monteny, and W. Baeyens. 1993. Geochemistry of mercury in an intertidal flat of the Scheldt Estuary. In: *Marine and Estuarine Gradients; ECSA Symposium*, Ghent, Belgium. 1991, P. Miere and M. Vincz, eds. Institute of Natural Conservation, Ministry of Flemish County. pp. 267-277.

Liang, L., M. Horvat, and N.S. Bloom. 1994. An improved speciation method for mercury by GC/CVAFS after aqueous phase ethylation and room temperature precollection. *Talanta* 41: 371-379.

Lindqvist, B., A. Jernelöv, K. Johansson, and H. Rodhe. 1984. Mercury in the Swedish environment, global and local sources. SNV PM 1816, Swedish Environment Protection Agency, S-171 85. Solna, Sweden.

Lindqvist, O., K. Johansson, M. Aastrup, A. Andersson, L. Bringmark, L. Hovsenius, C. Iverfeldt, M. Meili, and B. Timm. 1991. Mercury in the Swedish environment - recent research on causes, consequences and corrective methods. *Water Air Soil Pollut.* 55: i-261.

Lodenius, M., A. Seppanen, and M. Herranen. 1983. Accumulation of mercury in fish and man from reservoirs in northern Finland. *Water Air Soil Pollut.* 19: 237-246.

Lucotte, M., A. Mucci, C. Hillaire-Marcel, P. Pichet, and A. Grondin. 1995. Anthropogenic mercury enrichment in remote lakes of northern Québec (Canada). *Water Air Soil Pollut.* 80: 467-476.

Lucotte, M., S. Montgomery, B. Caron, and M. Kainz. 1998. Mercury in natural lakes and unperturbed terrestrial ecosystems of northern Québec, Ch. 3.1. In: *Mercury in natural environments and hydroelectric reservoirs of northern Québec (Canada)*, M. Lucotte, ed. Hydro-Québec and Environment Canada, pp. 46-80.

MacCrimmon, H.R., C.D. Wren, and B.L. Gots. 1983. Mercury uptake by lake trout, *Salvelinus namaycush*, relative to age, growth, and diet in Tadénac Lake with comparative data from other precambrian shield lakes. *Can. J. Fish. Aquat. Sci.* 40: 114-120.

MacDonald, D.D. 1994. A review of environmental quality criteria and guidelines for priority substances in the Fraser River basin. Report prepared for Environment Canada. Pacific and Yukon Region. North Vancouver, British Columbia.

Major, M.A., D.H. Rosenblatt, and K.A. Bostian. 1991. The octanol/water partition coefficient of methylmercuric chloride and methylmercuric hydroxide in pure water and salt solutions. *Environ. Toxicol. Chem.* 10: 5-8.

Malley, D.F., A.R. Stewart, and B.D. Hall. 1996. Uptake of methylmercury by the floater mussel, *Pyganodon grandis* (Bivalvia, Unionidae), caged in a flooded wetland. *Environ. Toxicol. Chem.* 15: 928-936.

Mason, R.P., W.F. Fitzgerald, and F.M.M. Morel. 1994. The biogeochemical cycling of elemental mercury: anthropogenic influences. *Geochim. Cosmochim. Acta* 58: 3191-3198.

## Canadian Tissue Residue Guidelines: Methylmercury

Mason, R.P., J.R. Reinfelder, and F.M.M. Morel. 1995. Bioaccumulation of mercury and methylmercury. *Water Air Soil Pollut.* 80: 915-921.

Mathers, R.A., and P.H. Johansen. 1985. The effects of feeding ecology on mercury accumulation in walleye (*Stizostedion vitreum*) and pike (*Esox lucius*) in Lake Simcoe. *Can. J. Zool.* 63: 2006-2012.

Matsumoto, N., and A. Spindle. 1982. Sensitivity of early mouse embryos to methylmercury toxicity. *Toxicol. Appl. Pharmacol.* 64: 108-117.

Maury, R., A. Boudou, F. Ribeyre, and P. Engrand. 1988. Experimental study of mercury transfer between artificially contaminated sediment ( $\text{CH}_3\text{HgCl}$ ) and macrophytes (*Elodea densa*). *Aquat. Toxicol.* 12: 213-228.

Maury-Brachet, R., F. Ribeyre, and A. Boudou. 1990. Actions and interactions of temperature and photoperiod on mercury accumulation by *Elodea densa* from sediment source. *Ecotoxicol. Environ. Safety* 20: 141-155.

McMurtry, M.J., D.L. Wales, W.A. Scheider, G.L. Beggs, and P.E. Dimond. 1989. Relationship of mercury concentrations in lake trout (*Salvelinus namaycush*) and smallmouth bass (*Micropterus dolomieu*) to the physical and chemical characteristics of Ontario lakes. *Can. J. Fish. Aquat. Sci.* 46: 426-434.

Mercury Team. 1998. Mercury in Atlantic Canada - A progress report. Environment Canada. Atlantic Region. 117 pp.

Metcalfe, C.D. 1986. An Introduction to aquatic toxicology. Peterborough, ON: Trent University. 172 pp.

Mierle, G. 1995. Sources of mercury to lakes on the Precambrian shield of Ontario. Proceedings of the 1995 Canadian Mercury Network Workshop. Sept 29-30. pp. 86-89.

Mierle, G., and R. Ingram. 1991. The role of humic substances in the mobilization of mercury from watersheds. *Water Air Soil Pollut.* 56: 349-357.

Miller, C.T., Z. Zawidska, E. Nagy, and S.M. Charbonneau. 1979. Indicators of genetic toxicity in leukocytes and granulocytic precursors after chronic methylmercury ingestion by cats. *Bull. Environ. Contam. Toxicol.* 21: 296-303.

Miller, R.W. 1975. The role of humic acids in the uptake and release of mercury by freshwater sediments. *Verh. Internat. Verein. Limnol.* 19: 2082-2086.

Minerals Yearbook. 1989. Mercury. 1: 705-708. US Bureau of Mines.

Miskimmin, B.M. 1991. Effect of natural levels of dissolved organic carbon (DOC) on methyl mercury formation and sediment-water partitioning. *Bull. Environ. Contam. Toxicol.* 47: 743-750.

Miskimmin, B.M., J. W. M. Rudd, and C.A. Kelly. 1992. Influence of dissolved organic carbon, pH, and microbial respiration rates on mercury methylation and demethylation in lake water. *Can. J. Fish. Aquat. Sci.* 49: 17-22.

## Canadian Tissue Residue Guidelines: Methylmercury

Mitra, S. 1986. Mercury in the ecosystem: Its dispersion and pollution today. Transtech Publications Ltd., Switzerland. ISBN-0-87849-529-0. 327 pp.

Mitsumori, K., K. Maita, T. Saito, S. Tsuda, and Y. Shirasu. 1981. Carcinogenicity of methylmercury chloride in ICR mice: Preliminary note on renal carcinogenesis. *Cancer Lett.* 12: 305-310.

Mitsumori, K., K. Takahashi, O. Matano, S. Goto, and Y. Shirasu. 1983. Chronic toxicity of methylmercury chloride in rats: Clinical study and chemical analysis. *Jpn. J. Vet. Sci.* 45(6): 747-757.

Mitsumori, K., K. Maita, and Y. Shirasu. 1984. Chronic toxicity of methylmercury chloride in rats: Pathological study. *Jpn. J. Vet. Sci.* 46(4): 549-557.

Mitsumori, K., M. Hiarano, H. Ueda, K. Maiata, and Y. Shirasu. 1990. Chronic toxicity and carcinogenicity of methylmercury chloride in B6C3F1 mice. *Fund. Appl. Toxicol.* 14: 179-190.

Mo, S.C., D.S. Choi, and J.W. Robinson. 1989. Uptake of mercury from aqueous solution by duckweed: the effects of pH, copper and humic acid. *J. Environ. Sci. Health* 24A: 135-146.

Moore, J.W., and S. Ramamoorthy 1984. Mercury, Chapter 7 In: *Heavy Metals in Natural Waters*, J.W. Moore and S. Ramamoorthy, eds. Springer-Verlag, New York. pp. 125-160.

Moore, J.W., and D.J. Sutherland. 1980. Mercury concentrations in fish inhabiting two polluted lakes in northern Canada. *Water Res.* 14: 903-907.

Moore, T.R., J.L. Bubier, A. Heyes, and R.J. Flett. 1995. Methyl and total mercury in boreal wetland plants, Experimental Lakes Area, Northwestern Ontario. *J. Environ. Qual.* 24: 845-850.

Morimoto, K., S. Iijima, and A. Koizumi. 1982. Selenite prevents the induction of sister-chromatid exchanges by methyl mercury and mercuric chloride in human whole-blood cultures. *Mutat. Res.* 102: 183-192.

Mortimer, D.C. 1985. Freshwater aquatic macrophytes as heavy metal monitors - the Ottawa River experience. *Environ. Monitor. Assess.* 5: 311-323.

Mucci, A., M. Lucotte, S. Montgomery, Y. Plourde, P. Pichet, and H.V. Tra. 1995. Mercury remobilization from flooded soils in a hydroelectric reservoir of Northern Québec, La Grande-2: results of a soil resuspension experiment. *Can. J. Fish. Aquat. Sci.* 52: 2507-2517.

Munro, I.C., E.A. Nera, S.M. Charbonneau, B. Junkins, and Z. Zawidzka. 1980. Chronic toxicity of methylmercury in the rat. *J. Environ. Pathol. Toxicol.* 3(5-6): 437-447.

Nagase, H., Y. Ose, T. Sato, and T. Ishikawa. 1982. Methylation of mercury by humic substances in an aquatic environment. *Sci. Total Environ.* 24: 133-142.

Nagpal, N.K. 1989. *Ambient Water Quality Criteria for Mercury*. Province of British Columbia. Water Management Branch, Ministry of Environment. Summary Report (11 pp) + Technical Appendix (105 pp.).

Newberne, P.M., O. Glaser, and L. Friedman. 1972. Chronic exposure of rats to methyl mercury

in fish protein. *Nature*. 237: 40-41.

Nicholson, J.K., and D. Osborn. 1984. Kidney lesions in juvenile starlings *Sturnus vulgaris* fed on a mercury-contaminated synthetic diet. *Environ. Pollut. (A)* 33: 195-206.

Nomiyama, K., K. Matsui, H. Nomiyama. 1980. Effects of temperature and other factors on the toxicity of methylmercury in mice. *Toxicol. Appl. Pharmacol.* 56: 392-398.

Norstrom, R.J., A.E. Mckinnon, and A.S.W. Defreitas. 1976. A bioenergetics-based model for pollutant accumulation by fish--simulation of PCB and methylmercury residue levels in Ottawa River yellow perch. *J. Fish. Res. Board Can.* 33: 248-267.

NPRI (National Pollutant Release Inventory). 1997. Summary Report 1995. Minister of Supply and Services Canada. 230 pp. <http://www.npri-inrp.com/queryform.cfm>.

Nriagu, J.O. 1979. *The biogeochemistry of mercury in the environment*. Elsevier/North-Holland Biomedical Press, New York. 696 pp.

———. 1989. A global assessment of natural sources of atmospheric trace metals. *Nature* 338: 47-49.

Nriagu, J.O., and J.M. Pacyna. 1988. Quantitative assessment of worldwide contamination of air, water and soils by trace metals. *Nature* 333: 134-139.

O'Connor, D.J., and S.W. Nielsen. 1981. Environmental survey of methylmercury levels in wild mink (*Mustela vison*) and otter (*Lutra canadensis*) from the northeastern United States and experimental pathology of methylmercurialism in the otter. Proc. Aug 3-11, Worldwide Furbearer Conference, Frostburg, MD. pp. 1728-1745.

OECD (Organisation for Economic Co-Operation and Development). 1994. Risk reduction monograph No. 4: Mercury - Background and national experience with reducing risk. OECD Environment Monograph Series No. 103. Environment Directorate. Paris. 159 pp.

Paine, P.J. 1994. Compliance with chlor-alkali mercury regulations, 1986-1989: Status report. Environment Canada. Environmental Protection Service. Industrial Programs Branch. Ottawa, ON. 43 pp.

Palmisano, F., N. Cardellicchio, and P.G. Zambonin. 1995. Speciation of mercury in dolphin liver: a two-stage mechanism for the demethylation accumulation process and role of selenium. *Mar. Environ. Res.* 40: 109-121.

Parkman, H., and M. Meili. 1993. Mercury in macroinvertebrates from Swedish forest lakes: influence of lake type, habitat, life cycle, and food quality. *Can. J. Fish. Aquat. Sci.* 50: 521-534.

Parks, J.W., A. Lutz, and J.A. Sutton. 1989. Water column methylmercury in the Wabigoon/English River-lake system: factors controlling concentrations, speciation, and net production. *Can. J. Fish. Aquat. Sci.* 46: 2184-2202.

Parks, J.W., J.A. Sutton, J.D. Hollinger, and D.D. Russell. 1988. Uptake of mercury by caged crayfish. *Appl. Organo. Chem.* 2: 181-184.

## Canadian Tissue Residue Guidelines: Methylmercury

Parks, J.W., C. Curry, D. Romani, and D.D. Russell. 1991. Young northern pike, yellow perch and crayfish as bioindicators in a mercury contaminated watercourse. *Environ. Mon. Assess.* 16: 39-73.

Pelletier, E. 1985. Mercury-selenium interactions in aquatic organisms: a review. *Mar. Environ. Res.* 18: 111-132.

Petrucchioli, L., and P. Turillazzi. 1991. Effect of methylmercury on acetylcholinesterase and serum cholinesterase activity in monkeys, *Macaca fascicularis*. *Bull. Environ. Contam. Toxicol.* 46: 769-773.

Pilgrim, W. and EMAN (Ecological Monitoring And Assessment Network). 1998. Mercury activities and measurements in the eastern Canadian provinces, Ch. VIII. In: Northeast states and eastern Canadian provinces - Mercury Study - a framework for action, M. Tatsutani, ed. NESCAUM, NEWMOA, NEIWPC, EMAN, 64 pp.

Ramamoorthy, S. 1982. Mercury analysis in environmental samples. Alberta Environmental Centre, Vegreville, AB. AECV82-M2. 61 pp.

Ramamoorthy, S., and D.J. Kushner. 1975. Heavy metal binding sites in river water. *Nature* 256: 399-401.

Ramamoorthy, S., T.C. Cheng, and D.J. Kushner. 1982. Effect of microbial life stages on the fate of methylmercury in natural waters. *Bull. Environ. Contam. Toxicol.* 29: 167-173.

———. 1983. Mercury speciation in water. *Can. J. Fish. Aquat. Sci.* 40: 85-89.

Ramamoorthy, S., J.W. Moore, and L. George. 1985. Partitioning of mercury in the North Saskatchewan River. *Chemosphere* 14: 1455-1468.

Ramlal, P.S., J.W.M. Rudd, A. Furutani, and L. Xun. 1985. The effect of pH on methyl mercury production and decomposition in lake sediments. *Can. J. Fish. Aquat. Sci.* 42: 685-692.

Ramlal, P.S., J.W.M. Rudd, and R.E. Hecky. 1986. Methods for measuring specific rates of mercury methylation and degradation and their use in determining factors controlling net rates of mercury methylation. *Appl. Environ. Microbiol.* 51: 110-114.

Ramlal, P.S., C. Anema, C. Furutani, R.E. Hecky, and J.W.M. Rudd. 1987. Mercury methylation studies at Southern Indian Lake, Manitoba: 1981-1983. *Can. Tech. Rep. Fish. Aquat. Sci.* 1490.

Ramlal, P.S., C.A. Kelly, J.W.M. Rudd, and A. Furutani. 1993. Sites of methyl mercury production in remote Canadian Shield lakes. *Can. J. Fish. Aquat. Sci.* 50: 972-979.

Rao, P.V., S.A. Jordan, and M.K. Bhatnagar. 1989. Ultrastructure of kidney of ducks exposed to methylmercury, lead and cadmium in combination. *J. Environ. Pathol. Toxicol. Oncol.* 9: 19-44.

Rasmussen, J.B., D.J. Rowan, D.R.S. Lean, and J.H. Carey. 1990. Food chain structure in Ontario lakes determines PCB levels in lake trout (*Salvelinus namaycush*) and other pelagic fish. *Can. J. Fish. Aquat. Sci.* 47: 2030-2038.

Ray, S., B.M. Jessop, J. Coffin, and D.A. Swetnam. 1984. Mercury and polychlorinated

biphenyls in striped bass (*Morone saxatilis*) from two Nova Scotia rivers. *Water Air Soil Pollut.* 21: 15-23.

Renner, R. 1998. Mercury health standard eased by U.S. agency. *Environ. Sci. Technol.* 32(1): 8A-9A.

Rice, D.C. 1989. Delayed neurotoxicity in monkeys exposed developmentally to methylmercury. *Neurotox.* 10: 645-650.

Richardson, G.M. 1994. Physical, chemical and geochemical factors influencing methylmercury accumulation in freshwater fish and humans in Ontario, Canada. Ph.D. Thesis, University of Ottawa ON. 195 pp.

Rieberger, K. 1992. Metal concentrations in fish tissue from uncontaminated BC lakes. *Water Quality Branch, BC Ministry of the Environment, Lands, and Parks.* 97 p.

Rissanen, K., J. Erkama, and J.K. Miettinen. 1970. Experiments on microbiological methylation of mercury (2+) ion by mud and sludge under aerobic and anaerobic conditions. *FAO Technical Conference on Marine Pollution and Its Effects on Living Resources and Fishing.* Nov. 6-25, Rome, Italy.

Robinson, J.B., and O.H. Tuovinen. 1984. Mechanisms of microbial resistance and detoxification of mercury and organomercury compounds: physiological, biochemical, and genetic analyses. *Microbiol. Rev.* 48: 95-124.

Rodgers, D.W. 1994. You are what you eat and a little bit more: bioenergetics-based models of methylmercury accumulation in fish revisited. In: *Mercury pollution: Integration and synthesis*, C.J. Watras and J.W. Huckabee, eds. Lewis Publishers, Ann Arbor, MI. pp. 427-439

Rodgers, D.W., and F.W.H. Beamish. 1983. Water quality modifies uptake of waterborne methylmercury by rainbow trout, *Salmo gairneri*. *Can. J. Fish. Aquat. Sci.* 40: 824-828.

Rodgers, D.W., and S.U. Qadri. 1982. Growth and mercury accumulation in yearling yellow perch, *Perca flavescens*, in the Ottawa River, Ontario. *Environ. Biol. Fish.* 7: 377-383.

Rodgers, D.W., T.A. Watson, J.S. Langan, and T.J. Wheaton. 1987. Effects of pH and feeding regime on methylmercury accumulation within aquatic microcosms. *Environ. Pollut.* 45: 261-274.

Rodgers, D.W., M. Dickman, and X. Han. 1995. Stories from old reservoirs: sediment Hg and Hg methylation in Ontario hydroelectric developments. *Water Air Soil Pollut.* 80: 829-839.

Ronald, K. S.V. Tessaro, J.F. Uthe, H.C. Freeman, and R. Frank. 1977. Methylmercury poisoning in the harp seal (*Pagophilus groenlandicus*). *Sci. Tot. Environ.* 8: 1-11.

RSC (Royal Society of Chemistry). 1994. D.O.S.E., J. Crombie, ed. RSC, Thomas Graham House, The Science Park, Cambridge. 875 pp.

Rudd, J.W.M. 1995. Sources of methylmercury to aquatic ecosystems. *Water Air Soil Pollut.* 80: 697-713.

## Canadian Tissue Residue Guidelines: Methylmercury

Rudd, J.W.M., A. Furutani, and M. Turner. 1980. Mercury methylation by fish intestinal contents. *Appl. Environ. Microbiol.* 40: 777-782.

Rudd, J.W.M., M.A. Turner, A. Furutani, A. Swick, and B.E. Townsend. 1983. The English-Wabigoon River system: I. A synthesis of recent research with a view towards mercury amelioration. *Can. J. Fish. Aquat. Sci.* 40: 2206-2217.

Ryder, J. P. 1974. Organochlorine and mercury residues in gull's eggs from western Ontario. *Can. Field-Nat.* 88: 349-352.

Sang, S., and B. Lourie. 1997. Mercury emission inventory in Ontario. *Environ. Monitor. Assess.* 47: 79-88.

Saouter, E., L. Hare, P.G.C. Campbell, A. Boudou, and F. Ribeyre. 1993. Mercury accumulation in the burrowing mayfly *Hexagenia rigida* (Ephemeroptera) exposed to CH<sub>3</sub>HgCl or HgCl<sub>2</sub> in water and sediment. *Water Res.* 27: 1041-1048.

Särkkä, J. 1979. Mercury and chlorinated hydrocarbons in zoobenthos of Lake Paijanne, Finland. *Arch. Environ. Contam. Toxicol.* 8: 161-173.

Scheuhammer, A. M. 1987. The chronic toxicity of aluminum, cadmium, mercury, and lead in birds: A review. *Environ. Pollut.* 46: 263-295.

Scheuhammer, A. M., and P. J. Blancher. 1994. Potential risk to common loons (*Gavia immer*) from methylmercury exposure in acidified lakes. *Hydrobiol.* 279/280: 445-455.

Scheuhammer, A.M., A.H.K. Wong, and D. Bond. 1998. Mercury and selenium accumulation in common loons (*Gavia immer*) and common mergansers (*Mergus merganser*) from eastern Canada. *Environ. Toxicol. Chem.* 17: 197-201.

Schintu, M., T. Kauri, and A. Kudo. 1989. Inorganic and methyl mercury in inland waters. *Water Res.* 23: 699-704.

Schroeder, W.H., G. Yarwood, and H. Niki. 1991. Transformation processes involving mercury species in the atmosphere -- results from a literature survey. *Water Air Soil Pollut.* 56: 653-666.

Scruton, D.A., E.L. Petticrew, L.J. LeDrew, M.R. Anderson, U.P. Williams, B.A. Bennett, and E.L. Hill. 1994. Methylmercury levels in fish tissue from three reservoir systems in Insular Newfoundland, Canada. In: *Mercury pollution: Integration and synthesis*, C.J. Watras and J.W. Huckabee, eds. Lewis Publishers. Boca Raton. pp. 441-455

Sellers, P., C.A. Kelly, J.W.M. Rudd, and A.R. MacHutchon. 1996. Photodegradation of methylmercury in lakes. *Nature* 380: 694-696.

Shilts, W.W., and W.B. Coker. 1995. Mercury anomalies in lake water and in commercially harvested fish, Kaminak Lake area, District of Keewatin, Canada. *Water Air Soil Pollut.* 80: 881-884.

Shimai, S., and H. Satoh. 1985. Behavioral teratology of methylmercury. *J. Toxicol. Sci.* 10: 199-216.



Sloss, L.L. 1995. Mercury emissions and effects - the role of coal. Perspectives. IEA Coal Research. London, U.K. pp. 1-39.

Smith, A.L., R.H. Green, and A. Lutz. 1975. Uptake of mercury by freshwater clams (family Unionidae). J. Fish. Res. Bd. Can. 32: 1297-1303.

Smith, C.M., and C. Rowan-West 1996. Mercury in Massachusetts: an evaluation of sources, emissions, impacts and controls. Office of Research and Standards, Bureau of Strategic Policy and Technology, Mass. Dept. Environ. Prot. (Available as an Internet manuscript).

Snelgrove-Hobson, S.M., P.V. Rao, and M.K. Bhatnagar. 1988. Ultra-structural alterations in the kidneys of Peking ducks fed methylmercury. Can. J. Vet. Res. 52: 89-98.

Sorenson, J.A., G.E. Glass, K.W. Schmidt, J.K. Huber, and G.R. Rapp Jr. 1990. Airborne mercury deposition and watershed characteristics in relation to mercury concentrations in water, sediments, plankton, and fish of eighty northern Minnesota lakes. Environ. Sci. Technol. 24: 1716-1727.

Southworth, G.R., M.J. Peterson, S.M. Adams, and B.G. Blaylock. 1994. Estimation of appropriate background concentrations for assessing mercury contamination in fish. Bull. Environ. Contam. Toxicol. 53: 211-218.

Spry, D.J., and J.G. Wiener. 1991. Metal bioavailability and toxicity to fish in low-alkalinity lakes: a critical review. Environ. Pollut. 71: 243-304.

St. Louis, V.L., J.W.M. Rudd, C.A. Kelly, K.G. Beaty, N.S. Bloom, and R.J. Flett. 1994. Importance of wetlands as sources of methyl mercury to boreal forest ecosystems. Can. J. Fish Aquat. Sci. 51: 1065-1076.

St. Louis, V.L., J.W.M. Rudd, C.A. Kelly, and L.A. Barrie. 1995. Wet deposition of methyl mercury in northwestern Ontario compared to other geographic locations. Water Air Soil Pollut. 80: 405-414.

Stendell, R.C., H.M. Ohlendorf, E.E. Klaas, and J.B. Elder. 1976. Mercury in eggs of aquatic birds, Lake St. Clair-1973. Pestic. Monit. J. 10: 7-9.

Stephens, G.R. 1995. Mercury concentrations in fish in a remote Canadian Arctic lake. Water Air Soil Pollut. 80: 633-636.

Stokes, P.M., and C.D. Wren. 1987. Bioaccumulation of mercury by aquatic biota in hydroelectric reservoirs: A review and consideration of mechanisms. In: Lead, mercury, cadmium and arsenic in the environment, T.C. Hutchinson and K.M. Meema, eds. John Wiley and Sons Ltd. Toronto, Canada. pp. 255-277.

Stokes, P.M., R.C. Bailey, and G.R. Groulx. 1985. Effects of acidification on metal availability to aquatic biota, with special reference to filamentous algae. Environ. Health Persp. 63: 79-87.

Stokes, P.M., S.I. Dreier, M.O. Farkas, and R.A.N. McLean. 1983. Mercury accumulation by filamentous algae: a promising biological monitoring system for methyl mercury in acid-stressed lakes. Environ. Pollut. (B) 5: 255-271.

## Canadian Tissue Residue Guidelines: Methylmercury

Summers, A.O. 1986. Organization, expression, and evolution of genes for mercury resistance. *Ann. Rev. Microbiol.* 40: 607-634.

Summers, A.O., and S. Silver. 1978. Microbial transformations of metals. *Ann. Rev. Microbiol.* 32: 637-672.

Sundberg, J., and A. Oskarsson. 1992. Placental and lactational transfer of mercury from rats exposed to methylmercury in their diet: speciation of mercury in the offspring. *J. Trace Elem. Exp. Med.* 5: 47-56.

Suns, K., and G. Hitchin. 1990. Interrelationships between mercury levels in yearling yellow perch, fish condition and water quality. *Water Air Soil Pollut.* 50: 255-265.

Suzuki, T., N. Imura, and T.W. Clarkson. 1991. Overview. In: *Advances in Mercury Toxicology*. T. Suzuki, N. Imura, and T.W. Clarkson, eds. Plenum Press, New York. pp. 1-32.

Thérien, N. and K. Morrison. 1998. In vitro release of mercury from flooded organic matter. Ch. 4.1. In: *Mercury in natural environments and hydroelectric reservoirs of northern Québec (Canada)*, M. Lucotte, ed. Hydro-Québec and Environment Canada, pp. 159-184.

Thompson, E.S. 1996. Mercury in Ontario wetlands: Concentrations in water, sediments, and a common aquatic plant in relation to geochemical variables. M.Sc. Thesis. The University of Ottawa, Ottawa, Ontario. pp. 113.

Thompson-Roberts, E.S., F.R. Pick, and G.E.M. Hall. 1999. Total Hg in water, sediment, and four species of aquatic macrophytes in the St. Lawrence River, near Cornwall, Ontario. *J. Great Lakes Res.* 25 (2): 294-304.

Tremblay, A. 1998. Bioaccumulation of mercury and methylmercury in invertebrates from natural boreal lakes, Ch. 3.2. In: *Mercury in natural environments and hydroelectric reservoirs of northern Québec (Canada)*, M. Lucotte, ed. Hydro-Québec, Environment Canada, Montreal. pp.81-103.

Tremblay, A., M. Lucotte, and D. Rowan. 1995. Different factors related to mercury concentration in sediments and zooplankton of 73 Canadian lakes. *Water Air Soil Pollut.* 80: 961-970.

Tremblay, A., M. Lucotte, and I. Rheault. 1996. Methylmercury in a benthic food web of two hydroelectric reservoirs and a natural lake of northern Québec (Canada). *Water Air Soil Pollut.* 91: 255-269.

Trudel, M., and J.B. Rasmussen. 1997. Modeling the elimination of mercury by fish. *Environ. Sci. Technol.* 31: 1715-1722.

US Department of Commerce. 1993. Comments on OECD Co-operative Risk Reduction Activities for Certain Dangerous Chemicals: Mercury (Draft Status Report). Letter of 22 December 1993 to Jim Darr, US EPA.

US DOH and HS (Dept. of Health and Human Services). 1994. Toxicological profile for mercury. Prep. By Clement International Corp. for Public Health Service. TP-93/10. 357 pp.

## Canadian Tissue Residue Guidelines: Methylmercury

US EPA (U.S. Environmental Protection Agency). 1989. Assessing human health risks from chemically contaminated fish and shellfish. A guidance manual. EPA-503/8-89-002. Office of Marine and Estuarine Protection and Office of Water Regulations and Standards. Washington, District of Columbia.

———. 1992. Characterization of products containing mercury in municipal solid waste in the United States, 1970 to 2000. US EPA, Office of Solid Waste, Municipal and Industrial Solid Waste Division, April 1992, #EPA 530-R-92-013.

———. 1993. Wildlife Exposure Factors Handbook. Volume 1 and 2. EPA/R-93/187a-b. Office of Health and Environmental Assessment Office of Research and Development. Washington, D.C. December 1993.

———. 1995a. Method 1669 - Method for sampling ambient water for determination of metals at EPA ambient criteria levels. Office of Water, Office of Science and Technology, Engineering and Analysis Division (4303), 401 M Street SW, Washington, DC. April 1995 with January 1996 revisions.

———. 1995b. Mercury Study Report to Congress. Office of Research and Development, Washington, DC. External Review Draft. EPA/600/P-94/002Ab.

———. 1995c. Trophic level and exposure analyses for selected piscivorous birds and mammals. Volume I. Analysis for species of the Great Lakes Basin (Draft). U.S. EPA Office of Science and Technology, Washington, DC.

———. 1995d. Final water quality guidance for the Great Lakes system: Final Rule. Fed. Regist. 60(56): 15366-15425 (March 23, 1995).

———. 1995e. Trophic level and exposure analyses for selected piscivorous birds and mammals. Vol. II: Analyses of species in the conterminous United States. Office of Water, Office of Science and Technology Washington, D.C.

———. 1996. Method 1631 - Mercury in water by oxidation, purge and trap, and CVFAS. Office of Water, Office of Science and Technology, Engineering and Analysis Division, Washington, DC.

———. 1997a. Mercury study report to Congress. Vol I: Executive summary. Office of Research and Development. Washington, DC.

———. 1997b. Mercury study report to Congress. Vol. VI: An ecological assessment for anthropogenic mercury emissions in the United States. Office of Research and Development. Washington, DC.

———. 1997c. Mercury study report to Congress. Vol VII: Characterization of human health and wildlife risks from mercury exposure in the United States. Office of Research and Development. Washington, DC.

U.S. NAS (National Academy of Sciences). 1977. An assessment of mercury in the environment. Environmental Studies Board, Nat. Acad. Sci, Washington D.C.

Vandal, G.M., R.P. Mason, and W.F. Fitzgerald. 1991. Cycling of volatile mercury in temperate

lakes. *Water Air Soil Pollut.* 56: 791-803.

Varekamp, J.C., and P.R. Buseck. 1986. Global mercury flux from volcanic and geothermal sources. *Appl. Geochem.* 1: 65-73.

Varshal, G.M., G.A. Sobolev, V.L. Barsukov, A.V. Koltsov, B.I. Kostin, T.F. Kudinova, Y.U. Stakheyev, and S.P. Tretyakova. 1985. Separation of volatile components from rocks under mechanical loadings as the source of hydrogeochemical anomalies preceding earthquakes. *Pure and Applied Geophysics* 122: 463-477.

Veiga, M.M., J.A. Meech, and N. Onate. 1994. Mercury from deforestation. *Nature* 368: 816-817.

Veiga, M.M., J.A. Meech, and D. Tromans. 1995. Mercury emissions and stability in the Amazon region. In: *Waste processing and recycling in the minerals and metallurgical industries*, S.R. Rao, L.M. Amaratunga, G.C. Richards, P.D. Kondos, eds. *Proc. Intern. Symp. At 34<sup>th</sup> Ann. Conf. Of Metallurgists of CIM, Vancouver, B.C.* pp. 523-538.

Verdon, R., and A. Tremblay. 1998. Mercury accumulation in fishes from the La Grande complex: influence of feeding habits and the concentration of mercury in the ingested preys, Ch. 5.2. In: *Mercury in natural environments and hydroelectric reservoirs of northern Québec (Canada)*, M. Lucotte, ed. Hydro-Québec, Environment Canada, Montreal. pp. 206-223.

Verdon, R., D. Brouard, C. Demers, R. Lalumiere, M. Laperle, and R. Schetagne. 1991. Mercury evolution (1978-1988) in fishes of the La Grande hydroelectric complex, Québec, Canada. *Water Air Soil Pollut.* 56: 405-417.

Vermeer, K., and F.A.J. Armstrong. 1972. Mercury in Canadian prairie ducks. *J. Wildl. Manage.* 36: 179-182.

Vermeer, K., F.A.J. Armstrong, and D.R.M. Hatch. 1973. Mercury in aquatic birds at Clay Lake, Western Ontario. *J. Wildl. Manage.* 37: 58-61.

Verschuuren, H.G., R. Kroes, E.M. den Tonkelaar, and others. 1976. Toxicity of methylmercury chloride in rats. III. Long-term toxicity study. *Toxicol.* 6: 107-123.

Verta, M., S. Rekolainen, and K. Kinnunen. 1986. Causes of increased fish mercury levels in Finnish reservoirs. *Vesientutkimuslaitoksen Julk.* 65: 44-58.

Visman, V., G. Mierle, and D.J. McQueen. 1995. Uptake of aqueous methylmercury by larval *Chaoborus americanus*. *Water Air Soil Pollut.* 80: 1007-1010.

Wagemann, R. 1994. Report to the Joint Fisheries Management Committee of the Inuvialuit Settlement Region, Inuvik, NWT. Canada. Dept. of Fisheries and Oceans Canada. pp. 1-16.

———. 1995. Mercury levels and their spatial and temporal trends in Arctic marine mammals. In: *Proceedings of the 1995 Canadian mercury network workshop*, T. Brydges, A. Perez-Fuentetaja, and D.J. McQueen, eds. York University, Toronto. pp. 9-12.

Wagemann, R., W.L. Lockhart, H. Welch, and S. Innes. 1995. Arctic marine mammals as integrators and indicators of mercury in the Arctic. *Water Air Soil Pollut.* 80: 683-693.

## Canadian Tissue Residue Guidelines: Methylmercury

Wagemann, R., S. Innes, and P.R. Richard. 1996. Overview and regional and temporal differences of heavy metals in Arctic whales and ringed seals in the Canadian Arctic. *Sci. Total Environ.* 186: 41-66.

Wagemann, R., E. Trebacz, R. Hunt, and G. Boila. 1997. Percent methylmercury and organic mercury in tissues of marine mammals and fish using different experimental and calculation methods. *Environ. Toxicol. Chem.* 16: 1859-1866.

Waiser, M.J., and R.D. Robarts. 1995. Microbial nutrient limitation in prairie saline lakes with high sulfate concentration. *Limnol. Oceanogr.* 40: 566-574.

Watras, C.J., and N.S. Bloom. 1992. Mercury and methylmercury in individual zooplankton: implications for bioaccumulation. *Limnol. Oceanogr.* 37: 1313-1318.

Watras, C.J., and J.W. Huckabee 1994. Mercury pollution: Integration and synthesis. Lewis Publishers. Boca Raton. 727 pp.

Watras, C.J., N.S. Bloom, R.J.M. Hudson, S. Gherini, R. Munson, S.A. Claus, K.A. Morrison, J. Hurley, J.G. Wiener, W.F. Fitzgerald, R. Mason, G. Vandal, D. Powell, R. Rada, L. Rislor, M. Winfrey, J. Elder, and J.W. Huckabee. 1994. Sources and fates of mercury and methylmercury in Wisconsin lakes. In: Mercury pollution: Integration and synthesis, C.J. Watras and J.W. Huckabee, eds. Lewis Publishers. pp. 153-177.

Watras, C.J., K.A. Morrison, and J.S. Host. 1995. Concentration of mercury species in relationship to other site-specific factors in the surface waters of northern Wisconsin lakes. *Limnol. Oceanogr.* 40: 556-565.

Watson, T. 1992. Evaluation of mercury concentration in selected environmental receptors in the Williston Lake and Peace River areas of British Columbia. Prepared for BC Hydro by Triton Environmental Consultants Ltd. Vancouver, BC.

Weseloh, D.V.C., S.M. Teeple, and M. Gilbertson. 1983. Double-crested cormorants of the Great Lakes: egg-laying parameters, reproductive failure, and contaminant residues in eggs, Lake Huron 1972-1973. *Can. J. Zool.* 61: 427-436.

Westcott, K., and J. Kalff. 1996. Environmental factors affecting methyl mercury accumulation in zooplankton. *Can. J. Fish. Aquat. Sci.* 53: 2221-2228.

WHO (World Health Organization). 1972. Evaluation of certain food additives and the contaminants mercury, lead, and cadmium. Sixteenth report of the joint FAO/WHO Expert Committee on Food Additives. April 4-12 Rome: World Health Organization. WHO Technical Report Series 505. 32 pp.

———. 1976. Mercury. Environmental Health Criteria 1. Geneva, Switzerland. 131 pp.

———. 1989. Mercury- Environmental Aspects. Environmental Health Criteria 86. IPCS - International Programme on Chemical Safety. World Health Organization. Geneva, Switzerland. 115 pp.

———. 1990. Methylmercury. Environmental Health Criteria 101. International Programme on Chemical Safety. Geneva. 144 pp.

## Canadian Tissue Residue Guidelines: Methylmercury

Williamson, D.A. 1986. Mercury in water and sediments in the Churchill and Nelson Rivers, Manitoba, Canada. In: Technical appendices to the summary report of the Canada-Manitoba agreement on the study and monitoring of Hg in the Churchill River diversion, Manitoba Environment and Workplace Safety and Health and Environment Canada, eds. Ottawa, Ontario. V.1, pp. 1-162.

Winfrey, M.R., and J.W.M. Rudd. 1990. Environmental factors affecting the formation of methylmercury in low pH lakes. *Environ. Toxicol. Chem.* 9: 853-870.

Wobeser, G. 1975. Prolonged oral administration of methyl mercury chloride to rainbow trout (*Salmo gairdneri*) fingerlings. *J. Fish. Res. Board Can.* 32: 2015-1023.

Wobeser, G., N.O. Nielsen, and B. Sciefer. 1976a. Mercury and mink I. Mercury contaminated fish as food for ranch mink. *Can. J. Comp. Med.* 40: 30-33.

———. 1976b. Mercury and mink II. Experimental methyl mercury intoxication. *Can. J. Comp. Med.* 40: 34-45.

Wolfe, M., and D. Norman. 1998. Effects of waterborne mercury on terrestrial wildlife at Clear Lake: evaluation and testing of a predictive model. *Environ. Toxicol. Chem.* 17: 214-227.

Wolfe, M.F., S. Schwarzbach, and R.A. Sulaiman. 1998. Effects of mercury on wildlife: a comprehensive review. *Environ. Contam. Toxicol.* 17: 146-160.

Wood, J.M., F.S. Kennedy, and C.G. Rosen. 1968. Synthesis of methylmercury compounds by extracts of a methanogenic bacterium. *Nature* 220: 173-174.

Wren, C.D. 1984. Distribution of metals in tissues of beaver, raccoon and otter from Ontario, Canada. *Sci. Total Environ.* 34: 177-184.

———. 1991. Relationship of mercury levels in sportfish with lake sediment and water quality variables. Prepared for the Research and Technology Branch, Ontario Ministry of the Environment. Toronto, Ontario. 36 pp.

Wren, C.D., and H.R. MacCrimmon. 1983. Mercury levels in the sunfish, *Lepomis gibbosus*, relative to pH and other environmental variables of Precambrian Shield Lakes. *Can. J. Fish. Aquat. Sci.* 40: 1737-1744.

Wren, C.D., and P.M. Stokes. 1988. Depressed mercury levels in biota from acid and metal stressed lakes near Sudbury, Ontario. *Ambio* 17: 28-30.

Wren, C.D., P.M. Stokes, and K.L. Fischer. 1986. Mercury levels in Ontario mink and otter relative to food levels and environmental acidification. *Can. J. Zool.* 64: 2854-2859.

Wren, C.D., D.B. Hunter, J.F. Leatherland, and P.M. Stokes. 1987a. The effects of polychlorinated biphenyls and methylmercury, singly and in combination, on mink. I: Uptake and toxic responses. *Arch. Environ. Contam. Toxicol.* 16: 441-447.

———. 1987b. The effects of polychlorinated biphenyls and methylmercury, singly and in combination, on mink. II: Reproduction and kit development. *Arch. Environ. Contam. Toxicol.* 16: 449-454.

## Canadian Tissue Residue Guidelines: Methylmercury


Wren, C.D., W.A. Scheider, D.L. Wales, B.W. Muncaster, and I.M. Gray. 1991. Relation between mercury concentrations in walleye (*Stizostedion vitreum vitreum*) and northern pike (*Esox lucius*) in Ontario lakes and influence of environmental factors. *Can. J. Fish. Aquat. Sci.* 48: 132-139.

Xun, L., N.E.R. Campbell, and J.W.M. Rudd. 1987. Measurements of specific rates of net methyl mercury production in the water column and surface sediments of acidified and circumneutral lakes. *Can. J. Fish. Aquat. Sci.* 44: 750-757.


Zauke, G.P. 1977. Mercury in benthic invertebrates of the Elbe Estuary. *Helgolander wissenschaftliche Meeresuntersuchungen* 29: 357-374.

Zillioux, E.J., D.B. Porcella, and J.M. Benoit. 1993. Mercury cycling and effects in freshwater wetland ecosystems. *Environ. Toxicol. Chem.* 12: 2245-2264.

1990 - 33 Tonnes



1995 - 12 Tonnes




**Figure 1. Mercury emissions in Canada by sector for 1990 and 1995 as estimated by the Transboundary Air Issues Branch of Environment Canada.**

Source: Environment Canada 1997; L. Trip, unpubl. data



## Canadian Tissue Residue Guidelines: Methylmercury




**Figure 2. Tonnes of mercury emitted by each Canadian province in 1990 as estimated by the Transboundary Air Issues Branch of Environment Canada.**

Source: Pilgrim and EMAN 1998


### 1990 Global Mercury Emissions

Total 2214 Tonnes



### 1990 North American Emissions

Total - 330 Tonnes



**Figure 3. Global and North American total mercury emissions for 1990 as reported by the Transboundary Air Issues Branch of Environment Canada.**

Source: Environment Canada 1997; L. Trip 1999, Environment Canada, pers. com.

Canadian Tissue Residue Guidelines: Methylmercury




Figure 4. The Mercury Cycle

**APPENDIX A: CANADIAN STRATEGIES UNDER A BILATERAL STRATEGY TO  
EVENTUALLY ELIMINATE ANTHROPOGENIC TOXIC SUBSTANCES IN THE  
GREAT LAKES BASIN.**

In Canada:

It has been estimated that between 2 700 and 3 450 kg of mercury are released to the atmosphere in Ontario annually from anthropogenic sources, while up to 2 500 kg are released to the waters of the Great Lakes Basin annually. Through an analysis of mercury uses and sources, significant sources of mercury have been identified and prioritised. These sources will be encouraged to develop strategies to reduce their releases by 90 percent from a baseline year of 1988 through adoption of pollution prevention measures.

In partnership with Pollution Probe, Canada and Ontario have identified potential industrial partners to participate in a unique three-way initiative to reduce or eliminate mercury in industrial or commercial applications. Co-ordination of this effort with U.S. partners is being considered, and the findings and approaches are being shared with the U.S. Virtual Elimination Pilot Project.

Activities by companies to date have resulted in significant reductions in mercury content in batteries (60 - 90 percent), fluorescent lamps (44 percent) and switches, while further reductions are planned, such as 70 percent by fluorescent lamp manufacturers by 2000. One impact of past mercury usage is that landfill emissions may be a source of mercury releases in the Great Lakes Basin, but the quantities released and possible control mechanisms need further consideration.

In applying the analytical framework in addressing mercury, relevant information from research projects undertaken by Environment Canada, Natural Resources Canada, and other agencies will be considered.

Canada will work with the U.S. and Mexico in implementing the North American Regional Action Plan for Mercury and will incorporate mercury reduction targets in its partnerships with commercial and industrial sectors in Ontario (Environment Canada and US EPA 1997).

Source: Environment Canada and US EPA 1997

**APPENDIX B-1: LEVELS OF TOTAL MERCURY AND METHYLMERCURY IN FRESHWATER AQUATIC PLANTS IN THE CANADIAN ENVIRONMENT.**

| Location                                | Year | Species  | Tissue | N  | Total Hg<br>$\mu\text{g}\cdot\text{kg}^{-1}$ dw | MeHg<br>$\mu\text{g}\cdot\text{kg}^{-1}$ dw | %<br>MeHg | Reference                    |
|---|------|--|--------|----|---|---|-----------|------------------------------|
| BC: Nahounli L.                         | 1982 | Macrophyte<br><i>Myriophyllum<br/>spicatum</i>   | Shoot  | 8  | 38 ± 6  | NR  | NR        | Siegel et al.<br>1985        |
| BC: Nahounli L.                         | 1982 | Macrophyte<br><i>Potamogeton</i> spp.  | Shoot  | 8  | 61 ± 11   | NR  | NR        | Siegel et al.<br>1985        |
| BC: Pinchi L.                           | 1982 | Macrophyte<br><i>Myriophyllum<br/>spicatum</i>   | Shoot  | 8  | 37 to 176                                       | NR  | NR        | Siegel et al.<br>1985        |
| BC: Pinchi L.                           | 1982 | Macrophyte<br><i>Potamogeton</i> spp.  | Shoot  | 8  | 15 to 132                                       | NR  | NR        | Siegel et al.<br>1985        |
| ON: Bentshoe L.<br>Muskoka-Haliburton   | 1988 | Macrophyte<br><i>Eriocaulon<br/>septangulare</i>   | Root   | 13 | 90 to 130                                       | NR  | NR        | Coquery and<br>Welbourn 1995 |
| ON: Bentshoe L.<br>Muskoka-Haliburton   | 1988 | Macrophyte<br><i>Eriocaulon<br/>septangulare</i>   | Leaf   | 13 | 40 to 110                                       | NR  | NR        | Coquery and<br>Welbourn 1995 |
| ON: Chub L.<br>South-central Ontario    | 1981 | Filamentous Algae<br><i>Mougeotia</i> spp. And<br><i>Spirogyra</i> spp.                            | Whole  | 3  | 90 to 330                                       | NR  | NR        | Stokes et al.<br>1983        |
| ON: Crosson L.<br>South-central Ontario | 1981 | Filamentous Algae<br><i>Mougeotia</i> spp. And<br><i>Spirogyra</i> spp.                            | Whole  | 3  | 40 to 180                                       | NR  | NR        | Stokes et al.<br>1983        |
| ON: Dickie L.<br>South-central Ontario  | 1981 | Filamentous Algae<br><i>Mougeotia</i> spp. and<br><i>Spirogyra</i> spp.                            | Whole  | 2  | 120; 120  | < 50  | NR        | Stokes et al.<br>1983        |
| ON: Experimental Lakes<br>Area          | 1992 | Macrophyte/sedge<br><i>Carex aquatilis</i> ,<br><i>C. rostrata</i> , and<br><i>Sparganium</i> spp. | Whole  | 4  | 6 to 18.8                                       | 1.3 to 6.4                                  | 37        | Moore et al.<br>1995         |

## Canadian Tissue Residue Guidelines: Methylmercury

| Location                                | Year | Species   | Tissue     | N  | Total Hg<br>$\mu\text{g}\cdot\text{kg}^{-1}$ dw | MeHg<br>$\mu\text{g}\cdot\text{kg}^{-1}$ dw | %<br>MeHg | Reference          |
|---|------|---|------------|----|---|---|-----------|--------------------|
| ON: Experimental Lakes Area Hummocks    | 1992 | Moss/bryophyte<br><i>Sphagnum</i> spp.                                  | Whole      | 2  | $36.3 \pm 10.7$                                 | $0.45 \pm 0.35$                             | 1.4       | Moore et al. 1995  |
| ON: Experimental Lakes Area Hollow/lawn | 1992 | Moss/bryophyte<br><i>Sphagnum</i> spp.                                  | Whole      | 5  | $46.7 \pm 18.8$                                 | $1.02 \pm 0.63$                             | 2.5       | Moore et al. 1995  |
| ON: Experimental Lakes Area Pool/stream | 1992 | Moss/bryophyte<br><i>Sphagnum</i> spp.                                  | Whole      | 2  | $92.5 \pm 38.1$                                 | $20.4 \pm 22.1$                             | 19        | Moore et al. 1995  |
| ON: Experimental Lakes Area             | 1992 | Moss/bryophyte<br><i>Sphagnum</i><br><i>angustifolium</i>               | Capitulum  | NR | 91.9  | 2.5   | 2.7       | Moore et al. 1995  |
| ON: Experimental Lakes Area             | 1992 | Moss/bryophyte<br><i>Sphagnum</i><br><i>angustifolium</i>               | Upper Stem | NR | 116.8   | 1.6   | 1.4       | Moore et al. 1995  |
| ON: Experimental Lakes Area             | 1992 | Moss/bryophyte<br><i>Sphagnum</i><br><i>angustifolium</i>               | Lower Stem | NR | 125.3   | 0.3   | 0.3       | Moore et al. 1995  |
| ON: Fawn L. South-central Ontario       | 1981 | Filamentous Algae<br><i>Mougeotia</i> spp. and<br><i>Spirogyra</i> spp. | Whole      | 2  | 87; 96  | NR  | NR        | Stokes et al. 1983 |
| ON: Harp L. South-central Ontario       | 1981 | Filamentous Algae<br><i>Mougeotia</i> spp. and<br><i>Spirogyra</i> spp. | Whole      | 2  | 160; 220  | NR  | NR        | Stokes et al. 1983 |
| ON: Heney L. South-central Ontario      | 1981 | Filamentous Algae<br><i>Mougeotia</i> spp. and<br><i>Spirogyra</i> spp. | Whole      | 3  | 80 to 93  | 50  | 55        | Stokes et al. 1983 |
| ON: Ottawa R., near Ottawa              | 1975 | Macrophyte<br><i>Elodea canadensis</i>                                  | Shoot      | 2  | 65 to 225                                       | NR  | 30        | Mortimer 1985      |
| ON: Ottawa R., near Ottawa              | 1975 | Macrophyte<br><i>Elodea canadensis</i>                                  | Root       | 2  | 159 to 260                                      | NR  | NR        | Mortimer 1985      |
| ON: Ottawa R., near Ottawa              | 1976 | Macrophyte<br><i>Elodea canadensis</i>                                  | Shoot      | 6  | NR  | NR  | 24        | Mortimer 1985      |

Canadian Tissue Residue Guidelines: Methylmercury

| Location                      | Year | Species   | Tissue | N | Total Hg<br>$\mu\text{g}\cdot\text{kg}^{-1}$ dw | MeHg<br>$\mu\text{g}\cdot\text{kg}^{-1}$ dw | %<br>MeHg | Reference     |
|-------------------------------|------|---|--------|---|---|---|-----------|---------------|
| ON: Ottawa R.,<br>near Ottawa | 1976 | Macrophyte<br><i>Elodea canadensis</i>            | Root   | 6 | NR  | NR  | 11        | Mortimer 1985 |
| ON: Ottawa R.,<br>near Ottawa | 1975 | Macrophyte<br><i>Phalaris<br/>arundinacea</i>     | Shoot  | 5 | 79 to 254                                       | NR  | NR        | Mortimer 1985 |
| ON: Ottawa R.,<br>near Ottawa | 1975 | Macrophyte<br><i>Phalaris<br/>arundinacea</i>     | Root   | 4 | 211 to 448                                      | NR  | NR        | Mortimer 1985 |
| ON: Ottawa R.,<br>near Ottawa | 1975 | Macrophyte<br><i>Sagittaria latifolia</i>         | Shoot  | 4 | 54 to 94  | NR  | 39        | Mortimer 1985 |
| ON: Ottawa R.,<br>near Ottawa | 1975 | Macrophyte<br><i>Sagittaria latifolia</i>         | Root   | 4 | 111 to 207                                      | NR  | 9         | Mortimer 1985 |
| ON: Ottawa R.,<br>near Ottawa | 1976 | Macrophyte<br><i>Sagittaria latifolia</i>         | Shoot  | 9 | NR  | NR  | 27        | Mortimer 1985 |
| ON: Ottawa R.,<br>near Ottawa | 1976 | Macrophyte<br><i>Sagittaria latifolia</i>         | Root   | 9 | NR  | NR  | 12        | Mortimer 1985 |
| ON: Ottawa R.,<br>near Ottawa | 1975 | Macrophyte<br><i>Sparganium<br/>augustifolium</i> | Shoot  | 3 | 71 to 215                                       | NR  | 30        | Mortimer 1985 |
| ON: Ottawa R.,<br>near Ottawa | 1975 | Macrophyte<br><i>Sparganium<br/>augustifolium</i> | Root   | 3 | 97 to 252                                       | NR  | 14        | Mortimer 1985 |
| ON: Ottawa R.,<br>near Ottawa | 1976 | Macrophyte<br><i>Sparganium<br/>augustifolium</i> | Shoot  | 5 | NR  | NR  | 33        | Mortimer 1985 |
| ON: Ottawa R.,<br>near Ottawa | 1976 | Macrophyte<br><i>Sparganium<br/>augustifolium</i> | Root   | 8 | NR  | NR  | 8         | Mortimer 1985 |
| ON: Ottawa R.,<br>near Ottawa | 1975 | Macrophyte<br><i>Sparganium<br/>eurycarpum</i>    | Shoot  | 3 | 50 to 164                                       | NR  | 48        | Mortimer 1985 |
| ON: Ottawa R.,<br>near Ottawa | 1975 | Macrophyte<br><i>Sparganium<br/>eurycarpum</i>    | Root   | 3 | 48 to 550                                       | NR  | 11        | Mortimer 1985 |

Canadian Tissue Residue Guidelines: Methylmercury

| Location                                  | Year | Species   | Tissue          | N       | Total Hg<br>$\mu\text{g}\cdot\text{kg}^{-1}$ dw | MeHg<br>$\mu\text{g}\cdot\text{kg}^{-1}$ dw | %<br>MeHg    | Reference                           |
|---|------|---|-----------------|---------|---|---|--------------|-------------------------------------|
| ON: Ottawa R.,<br>near Ottawa             | 1976 | Macrophyte<br><i>Sparganium<br/>eurycarpum</i>                          | Shoot           | 4       | NR  | NR  | 23           | Mortimer 1985                       |
| ON: Ottawa R.,<br>near Ottawa             | 1976 | Macrophyte<br><i>Sparganium<br/>eurycarpum</i>                          | Root            | 4       | NR  | NR  | 8            | Mortimer 1985                       |
| ON: Plastic L.<br>South-central Ontario   | 1981 | Filamentous Algae<br><i>Mougeotia</i> spp. and<br><i>Spirogyra</i> spp. | Whole           | 2       | 160; 180  | NR  | NR           | Stokes et al.<br>1983               |
| ON: Red Chalk L.<br>South-central Ontario | 1981 | Filamentous Algae<br><i>Mougeotia</i> spp. and<br><i>Spirogyra</i> spp. | Whole           | 1       | 270   | NR  | NR           | Stokes et al.<br>1983               |
| ON: St. Lawrence R.,<br>near Cornwall     | 1994 | Macrophyte<br><i>Elodea canadensis</i>                                  | Shoot           | 40      | 58 to 225                                       | NR  | NR           | Thompson-<br>Roberts et al.<br>1999 |
| ON: St. Lawrence R.,<br>near Cornwall     | 1994 | Macrophyte<br><i>Myriophyllum<br/>spicatum</i>                          | Shoot           | 55      | 63 to 240                                       | NR  | NR           | Thompson-<br>Roberts et al.<br>1999 |
| ON: St. Lawrence R.,<br>near Cornwall     | 1976 | Macrophyte<br><i>Myriophyllum<br/>spicatum</i>                          | Shoot           | NR      | 68 to 247                                       | NR  | 0 to<br>30 % | Mortimer 1985                       |
| ON: St. Lawrence R.,<br>near Cornwall     | 1978 | Macrophyte<br><i>Myriophyllum<br/>spicatum</i>                          | Shoot           | NR      | 17 to 1600                                      | NR  | 0 to<br>30 % | Mortimer 1985                       |
| ON: St. Lawrence R.,<br>near Cornwall     | 1994 | Macrophyte<br><i>Nuphar variegatum</i>                                  | Leaf<br>Petiole | 11<br>5 | 8 to 21<br>6 to 37                              | NR  | NR           | Thompson-<br>Roberts et al.<br>1999 |
| ON: St. Lawrence R.,<br>Near Cornwall     | 1994 | Macrophyte<br><i>Potamogeton<br/>crispus</i>                            | Shoot           | 15      | 80 to 85  | NR  | NR           | Thompson-<br>Roberts et al.<br>1999 |
| ON: St. Lawrence R.,<br>Near Cornwall     | 1975 | Macrophyte<br><i>Sagittaria latifolia</i>                               | Shoot<br>Root   | NR      | NR  | NR  | 38.6<br>8.7  | Mortimer 1985                       |



Canadian Tissue Residue Guidelines: Methylmercury

| Location                                   | Year | Species   | Tissue          | N       | Total Hg<br>$\mu\text{g}\cdot\text{kg}^{-1}$ dw | MeHg<br>$\mu\text{g}\cdot\text{kg}^{-1}$ dw | %<br>MeHg | Reference             |
|--|------|---|-----------------|---------|---|---|-----------|-----------------------|
| ON: Swan L.<br>South-central Ontario       | 1981 | Filamentous Algae<br><i>Mougeotia</i> spp. and<br><i>Spirogyra</i> spp. | Whole           | 2       | 120; 130  | < 50  | NR        | Stokes et al.<br>1983 |
| ON: Wetlands (22)<br>South-central Ontario | 1993 | Macrophyte<br><i>Nuphar variegatum</i>                                  | Leaf<br>Petiole | 11<br>0 | 13 to 29<br>14 to 31                            | NR  | NR        | Thompson 1996         |

NR = Not reported

**APPENDIX B-2A: LEVELS OF TOTAL MERCURY AND METHYLMERCURY IN FRESHWATER INVERTEBRATES IN THE CANADIAN ENVIRONMENT.**

| Location                                  | Year | Species                          | Life Stage or Size | Tissue | N | Total Hg $\mu\text{g}\cdot\text{g}^{-1}$ ww <sup>a</sup> | MeHg $\mu\text{g}\cdot\text{g}^{-1}$ ww <sup>a</sup> | % MeHg | Reference     |
|---|------|----------------------------------|--------------------|--------|---|--|--|--------|---------------|
| MB: Mynarksi L. Notigi Reservoir          | 1981 | Chironomids                      | Larvae             | Whole  | 3 | 0.014 - 0.025  | NR   | NR     | Jackson 1988b |
| MB: Mynarksi L. Notigi Reservoir          | 1981 | Clams<br><i>Pelecypoda</i>       | Adult              | Whole  | 1 | 0.015  | NR   | NR     | Jackson 1988b |
| MB: Mynarksi L. Notigi Reservoir          | 1981 | Earthworms<br><i>Oligochaeta</i> | Adult              | Whole  | 1 | 0.337  | NR   | NR     | Jackson 1988b |
| MB: Mynarksi L. Notigi Reservoir          | 1981 | Roundworms<br><i>Nematoda</i>    | Adult              | Whole  | 1 | 0.164  | NR   | NR     | Jackson 1988b |
| MB: Notigi L. Notigi Reservoir            | 1981 | Chironomids                      | Larvae<br>Pupae    | Whole  | 1 | 0.024<br>0.026   | NR   | NR     | Jackson 1988b |
| MB: Notigi L. Notigi Reservoir            | 1981 | Earthworms<br><i>Oligochaeta</i> | Adult              | Whole  | 1 | 0.051  | NR   | NR     | Jackson 1988b |
| MB: Notigi L. Notigi Reservoir            | 1981 | Roundworms<br><i>Nematoda</i>    | Adult              | Whole  | 1 | 0.032  | NR   | NR     | Jackson 1988b |
| MB: Rat L. Notigi Reservoir               | 1981 | Chironomids                      | Larvae             | Whole  | 4 | 0.017 - 0.028  | NR   | NR     | Jackson 1988b |
| MB: Rat L. Notigi Reservoir               | 1981 | Clams<br><i>Pelecypoda</i>       | Adult              | Whole  | 1 | 0.078  | NR   | NR     | Jackson 1988b |
| MB: Rat L. Notigi Reservoir               | 1981 | Earthworms<br><i>Oligochaeta</i> | Adult              | Whole  | 1 | 0.043  | NR   | NR     | Jackson 1988b |
| MB: Rat L. Notigi Reservoir               | 1981 | Roundworms<br><i>Nematoda</i>    | Adult              | Whole  | 1 | 0.097  | NR   | NR     | Jackson 1988b |
| MB: Southern Indian L. Long Bay – shallow | 1981 | Chironomids                      | Larvae             | Whole  | 1 | 0.208  | NR   | NR     | Jackson 1988b |
| MB: Southern Indian L. Long Bay – shallow | 1981 | Clams<br><i>Pelecypoda</i>       | Adult              | Whole  | 1 | 0.307  | NR   | NR     | Jackson 1988b |

## Canadian Tissue Residue Guidelines: Methylmercury

| Location  | Year        | Species                                | Life Stage or Size | Tissue | N  | Total Hg<br>$\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | MeHg<br>$\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | % MeHg | Reference              |
|---|-------------|--|--------------------|--------|----|--|--|--------|------------------------|
| MB: Southern Indian L. Long Bay – shallow               | 1981        | Earthworms<br><i>Oligochaeta</i>       | Adult              | Whole  | 1  | 0.698  | NR   | NR     | Jackson 1988b          |
| MB: Southern Indian L. Long Bay – shallow               | 1981        | Roundworms<br><i>Nematoda</i>          | Adult              | Whole  | 1  | 0.459  | NR   | NR     | Jackson 1988b          |
| MB: Southern Indian L. South Bay                        | 1981        | Chironomids                            | Larvae             | Whole  | 1  | 0.115  | NR   | NR     | Jackson 1988b          |
| MB: Southern Indian L. South Bay                        | 1981        | Clams<br><i>Pelecypoda</i>             | Adult              | Whole  | 1  | 0.023  | NR   | NR     | Jackson 1988b          |
| ON & QC: Lakes from 44°12N – 48°07N and 70°52W - 80°17W | 1990 - 1991 | Zooplankton <sup>b</sup><br>- unsorted | >225 $\mu\text{m}$ | Whole  | 24 | 0.108 $\pm$ 0.076 dw   | NR   | NR     | Tremblay et al. 1995   |
| ON: Bear L.   | 1986        | Crayfish<br><i>Cambarus bartoni</i>    | 7.1 $\pm$ 2.2 g    | Muscle | 39 | 0.124 $\pm$ 0.044  | NR   | NR     | Allard and Stokes 1989 |
| ON: Blue Chalk L.                                       | 1991        | Crayfish<br><i>Orconectes virilis</i>  | 2.1-3.5 cm         | Muscle | 11 | 0.022 to 0.055   | NR   | NR     | Headon et al. 1996     |
| ON: Clay L. East basin                                  | 1979        | Crayfish<br><i>Orconectes virilis</i>  | Adult              | Muscle | 20 | 2.2 $\pm$ 0.66   | NR   | NR     | Parks et al. 1991      |
| ON: Clay L. East basin                                  | 1981        | Crayfish<br><i>Orconectes virilis</i>  | Adult              | Muscle | 19 | 0.96 $\pm$ 0.34  | NR   | NR     | Parks et al. 1991      |

Canadian Tissue Residue Guidelines: Methylmercury

| Location                     | Year | Species  | Life Stage or Size | Tissue | N  | Total Hg<br>$\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | MeHg<br>$\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | % MeHg | Reference              |
|------------------------------|------|--|--------------------|--------|----|--|--|--------|------------------------|
| ON: Cygnet L.                | 1980 | Crayfish<br><i>Orconectes virilis</i>                      | Adult              | Muscle | 18 | $0.06 \pm 0.02$  | NR   | NR     | Parks et al. 1991      |
| ON: Dickie L.                | 1986 | Crayfish<br><i>Cambarus robustus</i>                       | $4.9 \pm 5.5$ g    | Muscle | 19 | $0.087 \pm 0.033$  | NR   | NR     | Allard and Stokes 1989 |
| ON: Eagle L./ Eagle R.       | 1980 | Crayfish<br><i>Orconectes virilis</i>                      | Adult              | muscle | 31 | $0.07 \pm 0.03$  | NR   | NR     | Parks et al. 1991      |
| ON: Eagle L./ Eagle R.       | 1981 | Crayfish<br><i>Orconectes virilis</i>                      | Adult              | Muscle | 6  | $0.06 \pm 0.05$  | NR   | NR     | Parks et al. 1991      |
| ON: Eaglenest L.             | 1979 | Crayfish<br><i>Orconectes virilis</i>                      | Adult              | Muscle | 18 | $0.43 \pm 0.12$  | NR   | NR     | Parks et al. 1991      |
| ON: Eaglenest L.             | 1980 | Crayfish<br><i>Orconectes virilis</i>                      | Adult              | Muscle | 26 | $0.21 \pm 0.10$  | NR   | NR     | Parks et al. 1991      |
| ON: ELARP Reservoir (L. 104) | 1992 | Freshwater mussel<br><i>Pyganodon grandis</i>              | Adult              | Foot   | 12 | $0.422 \pm 0.033$ dw   | $0.250 \pm 0.018$ dw   | 62%    | Malley et al. 1996     |
| ON: ELARP Reservoir (L. 104) | 1992 | Freshwater mussel<br><i>Pyganodon grandis</i> <sup>c</sup> | Adult              | Gill   | 11 | $0.315 \pm 0.044$ dw   | $0.171 \pm 0.019$ dw   | 58%    | Malley et al. 1996     |

## Canadian Tissue Residue Guidelines: Methylmercury

| Location                           | Year | Species  | Life Stage or Size | Tissue  | N  | Total Hg<br>$\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | MeHg<br>$\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | % MeHg | Reference          |
|------------------------------------|------|--|--------------------|---------|----|--|--|--------|--------------------|
| ON: ELARP Reservoir (L. 104)       | 1992 | Freshwater mussel<br><i>Pyganodon grandis</i>              | Adult              | Kidney  | 6  | $1.992 \pm 0.250$<br>dw  | $0.288 \pm 0.023$<br>dw                                      | 15%    | Malley et al. 1996 |
| ON: ELARP Reservoir (L. 104)       | 1992 | Freshwater mussel<br><i>Pyganodon grandis</i> <sup>c</sup> | Adult              | Mantle  | 12 | $0.245 \pm 0.014$<br>dw  | $0.108 \pm 0.009$<br>dw                                      | 45%    | Malley et al. 1996 |
| ON: ELARP Reservoir (L. 104)       | 1992 | Freshwater mussel<br><i>Pyganodon grandis</i>              | Adult              | Viscera | 12 | $0.322 \pm 0.014$<br>dw  | $0.151 \pm 0.009$<br>dw                                      | 47%    | Malley et al. 1996 |
| ON: ELARP Reservoir (L. 104)       | 1992 | Freshwater mussel<br><i>Pyganodon grandis</i>              | Adult              | Whole   | 12 | $0.368 \pm 0.018$<br>dw  | $0.161 \pm 0.009$<br>dw                                      | 44%    | Malley et al. 1996 |
| ON: English R., inflow to Ball L.  | 1979 | Crayfish<br><i>Orconectes virilis</i>                      | Adult              | Muscle  | 30 | 0.25   | NR   | NR     | Parks et al. 1991  |
| ON: English R., inflow to Ball L.  | 1980 | Crayfish<br><i>Orconectes virilis</i>                      | Adult              | Muscle  | 17 | $0.15 \pm 0.06$  | NR   | NR     | Parks et al. 1991  |
| ON: English R., inflow to Ball L.  | 1981 | Crayfish<br><i>Orconectes virilis</i>                      | Adult              | Muscle  | 42 | $0.12 \pm 0.06$  | NR   | NR     | Parks et al. 1991  |
| ON: English R., outflow of Ball L. | 1979 | Crayfish<br><i>Orconectes virilis</i>                      | Adult              | Muscle  | 30 | $0.68 \pm 0.36$  | NR   | NR     | Parks et al. 1991  |
| ON: English R., outflow of Ball L. | 1980 | Crayfish<br><i>Orconectes virilis</i>                      | Adult              | Muscle  | 18 | $0.42 \pm 0.20$  | NR   | NR     | Parks et al. 1991  |

## Canadian Tissue Residue Guidelines: Methylmercury

| Location                              | Year | Species                                  | Life Stage or Size                 | Tissue | N  | Total Hg<br>$\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | MeHg<br>$\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | % MeHg | Reference                  |
|---------------------------------------|------|--|------------------------------------|--------|----|--|--|--------|----------------------------|
| ON: English R.,<br>outflow of Ball L. | 1981 | Crayfish<br><i>Orconectes virilis</i>    | Adult                              | Muscle | 40 | $0.48 \pm 0.23$  | NR   | NR     | Parks et al.<br>1991       |
| ON: Four Mile L.                      | 1986 | Crayfish<br><i>Orconectes propinquus</i> | $2.8 \pm 1.1$ g                    | Muscle | 27 | $0.042 \pm 0.018$  | NR   | NR     | Allard and<br>Stokes 1989  |
| ON: Four Mile L.                      | 1986 | Crayfish<br><i>Orconectes virilis</i>    | $4.9 \pm 2.3$ g                    | Muscle | 14 | $0.095 \pm 0.025$  | NR   | NR     | Allard and<br>Stokes 1989  |
| ON: Galeairy L.                       | 1986 | Crayfish<br><i>Orconectes obscurus</i>   | $4.4 \pm 2.3$ g                    | Muscle | 31 | $0.073 \pm 0.029$  | NR   | NR     | Allard and<br>Stokes 1989  |
| ON: Gull L.                           | 1986 | Crayfish<br><i>Cambarus robustus</i>     | $16.8 \pm 9.7$ g                   | Muscle | 22 | $0.062 \pm 0.036$  | NR   | NR     | Allard and<br>Stokes 1989  |
| ON: Kimball L.                        | 1986 | Crayfish<br><i>Cambarus bartoni</i>      | $7.3 \pm 3.5$ g                    | Muscle | 43 | $0.089 \pm 0.057$  | NR   | NR     | Allard and<br>Stokes 1989  |
| ON: L. Vernon                         | 1986 | Crayfish<br><i>Cambarus bartoni</i>      | $10.2 \pm 3.4$ g                   | Muscle | 30 | $0.227 \pm 0.102$  | NR   | NR     | Allard and<br>Stokes 1989  |
| ON: L. Vernon,<br>inlet               | 1986 | Crayfish<br><i>Cambarus bartoni</i>      | $4.9 \pm 3.8$ g                    | Muscle | 8  | $0.304 \pm 0.073$  | NR   | NR     | Allard and<br>Stokes 1989  |
| ON: Lakes<br>South-central<br>Ontario | 1993 | Daphnia spp.                             | Adult<br>>363<br>$\mu\text{m}$     | Whole  | 24 | NR   | 0.019 to<br>0.448 dw   | NR     | Westcott and<br>Kalff 1996 |
| ON: Little Hawk<br>L.                 | 1986 | Crayfish<br><i>Cambarus bartoni</i>      | $7.1 \pm 2.8$ g<br>$8.7 \pm 3.5$ g | Muscle | 16 | $0.073 \pm 0.038$<br>$0.072 \pm 0.020$                           | NR   | NR     | Allard and<br>Stokes 1989  |

## Canadian Tissue Residue Guidelines: Methylmercury

| Location                  | Year | Species                               | Life Stage or Size         | Tissue | N      | Total Hg<br>$\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | MeHg<br>$\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | % MeHg | Reference              |
|---------------------------|------|---------------------------------------|----------------------------|--------|--------|--|--|--------|------------------------|
| ON: Little Hawk L., inlet | 1986 | Crayfish<br><i>Cambarus bartoni</i>   | 2.9 ± 1.6 g<br>2.7 ± 3.1 g | Muscle | 6<br>8 | 0.197 ± 0.101<br>0.123 ± 0.020                                   | NR   | NR     | Allard and Stokes 1989 |
| ON: Opeongo L.            | 1986 | Crayfish<br><i>Cambarus bartoni</i>   | 4.5 ± 2.2 g                | Muscle | 31     | 0.077 ± 0.043  | NR   | NR     | Allard and Stokes 1989 |
| ON: Peninsula L.          | 1986 | Crayfish<br><i>Cambarus bartoni</i>   | 5.4 ± 1.8 g                | Muscle | 40     | 0.069 ± 0.030  | NR   | NR     | Allard and Stokes 1989 |
| ON: Rebecca L.            | 1986 | Crayfish<br><i>Cambarus bartoni</i>   | 10.0 ± 4.0 g               | Muscle | 33     | 0.145 ± 0.070  | NR   | NR     | Allard and Stokes 1989 |
| ON: Rebecca L.            | 1986 | Crayfish<br><i>Orconectes virilis</i> | 9.3 ± 6.5 g                | Muscle | 27     | 0.059 ± 0.035  | NR   | NR     | Allard and Stokes 1989 |
| ON: Roughrock L.          | 1980 | Crayfish<br><i>Orconectes virilis</i> | Adult                      | Muscle | 11     | 0.16 ± 0.06  | NR   | NR     | Parks et al. 1991      |
| ON: Round L.              | 1986 | Crayfish<br><i>Cambarus bartoni</i>   | 4.5 ± 2.5 g                | Muscle | 35     | 0.081 ± 0.047  | NR   | NR     | Allard and Stokes 1989 |
| ON: Sand L.               | 1979 | Crayfish<br><i>Orconectes virilis</i> | Adult                      | Muscle | 14     | 0.26 ± 0.11  | NR   | NR     | Parks et al. 1991      |
| ON: Umfreville L.         | 1980 | Crayfish<br><i>Orconectes virilis</i> | Adult                      | Muscle | 4      | 0.25 ± 0.08  | NR   | NR     | Parks et al. 1991      |
| ON: Wabigoon L.           | 1987 | Crayfish<br><i>Orconectes virilis</i> | Adult                      | Muscle | 28     | 0.15 <sup>d</sup>  | NR   | NR     | Parks et al. 1988      |

Canadian Tissue Residue Guidelines: Methylmercury

| Location                            | Year | Species                                  | Life Stage or Size                                 | Tissue | N              | Total Hg<br>$\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | MeHg<br>$\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | % MeHg | Reference              |
|-------------------------------------|------|--|--|--------|----------------|--|--|--------|------------------------|
| ON: Wabigoon L.                     | 1979 | Crayfish<br><i>Orconectes virilis</i>    | Adult  | Muscle | 28             | $0.16 \pm 0.16$  | NR   | NR     | Parks et al. 1991      |
| ON: Wabigoon L.                     | 1980 | Crayfish<br><i>Orconectes virilis</i>    | Adult  | Muscle | 30             | $0.09 \pm 0.04$  | NR   | NR     | Parks et al. 1991      |
| ON: Wabigoon L.                     | 1981 | Crayfish<br><i>Orconectes virilis</i>    | Adult  | Muscle | 4              | $0.07 \pm 0.03$  | NR   | NR     | Parks et al. 1991      |
| ON: Wabigoon R., inflow to Ball L.  | 1979 | Crayfish<br><i>Orconectes virilis</i>    | Adult  | Muscle | 21             | $0.80 \pm 0.20$  | NR   | NR     | Parks et al. 1991      |
| ON: Wabigoon R., inflow to Ball L.  | 1980 | Crayfish<br><i>Orconectes virilis</i>    | Adult  | Muscle | 29             | $1.7 \pm 0.49$   | NR   | NR     | Parks et al. 1991      |
| ON: Wabigoon R., inflow to Ball L.  | 1981 | Crayfish<br><i>Orconectes virilis</i>    | Adult  | Muscle | 40             | $0.91 \pm 0.25$  | NR   | NR     | Parks et al. 1991      |
| ON: Wabigoon R., inflow to Clay L.  | 1980 | Crayfish<br><i>Orconectes virilis</i>    | Adult  | Muscle | 30             | $0.79 \pm 0.26$  | NR   | NR     | Parks et al. 1991      |
| ON: Wabigoon R., inflow to Clay L.  | 1981 | Crayfish<br><i>Orconectes virilis</i>    | Adult  | Muscle | 13             | $0.76 \pm 0.20$  | NR   | NR     | Parks et al. 1991      |
| ON: Wabigoon R., outflow of Clay L. | 1981 | Crayfish<br><i>Orconectes virilis</i>    | Adult  | Muscle | 15             | $0.75 \pm 0.25$  | NR   | NR     | Parks et al. 1991      |
| ON: Waseosa L.                      | 1986 | Crayfish<br><i>Orconectes propinquus</i> | 2.5 ±<br>1.0 g<br>8.6 ±<br>3.9 g<br>3.4 ±<br>2.3 g | Muscle | 34<br>23<br>23 | $0.095 \pm 0.052$<br>$0.082 \pm 0.034$<br>$0.083 \pm 0.036$      | NR   | NR     | Allard and Stokes 1989 |



## Canadian Tissue Residue Guidelines: Methylmercury

| Location                                       | Year | Species                                  | Life Stage or Size | Tissue | N  | Total Hg<br>$\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | MeHg<br>$\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | % MeHg | Reference            |
|--|------|--|--------------------|--------|----|--|--|--------|----------------------|
| QC: Duncan L., control for La Grande Reservoir | 1992 | Coleoptera<br>Dytiscidae                 | Larvae/<br>Pupae   | Whole  | 11 | 0.175 dw   | 0.107 dw   | 61%    | Tremblay et al. 1996 |
| QC: Duncan L., control for La Grande Reservoir | 1992 | Corixidae<br><i>Sigara</i> spp.          | Larvae/<br>Pupae   | Whole  | 17 | 0.256 dw   | 0.124 dw   | 49%    | Tremblay et al. 1996 |
| QC: Duncan L., control for La Grande Reservoir | 1992 | Ephemeroptea<br><i>Leptophlebia</i> spp. | Larvae/<br>Pupae   | Whole  | 11 | 0.129 dw   | 0.014 dw   | 11%    | Tremblay et al. 1996 |
| QC: Duncan L., control for La Grande Reservoir | 1992 | Odonata<br><i>Somatochlora</i> spp.      | Larvae/<br>Pupae   | Whole  | 4  | 0.136 dw   | 0.102 dw   | 75%    | Tremblay et al. 1996 |
| QC: Duncan L., control for La Grande Reservoir | 1992 | Trichoptera<br><i>Agrypnia</i> spp       | Larvae/<br>Pupae   | Whole  | 5  | 0.124 dw   | 0.061 dw   | 50%    | Tremblay et al. 1996 |
| QC: Duncan L., control for La Grande Reservoir | 1992 | Trichoptera<br><i>Grammotaulius</i> spp. | Larvae/<br>Pupae   | Whole  | 4  | 0.120 dw   | 0.056 dw   | 47%    | Tremblay et al. 1996 |
| QC: Duncan L., control for La Grande Reservoir | 1992 | Trichoptera<br><i>Asynarchus</i> spp     | Larvae/<br>Pupae   | Whole  | 10 | 0.143 dw   | 0.029 dw   | 20%    | Tremblay et al. 1996 |
| QC: La Grande-1 Reservoir                      | 1992 | Chironomids                              | Larvae/<br>Pupae   | Whole  | 12 | 1.020 dw   | 0.064 dw   | 6%     | Tremblay et al. 1996 |
| QC: La Grande-1 Reservoir                      | 1992 | Corixidae<br><i>Sigara</i> spp.          | Larvae/<br>Pupae   | Whole  | 10 | 0.514 dw   | 0.411 dw   | 80%    | Tremblay et al. 1996 |

Canadian Tissue Residue Guidelines: Methylmercury

| Location                            | Year | Species                                   | Life Stage or Size | Tissue | N  | Total Hg<br>$\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | MeHg<br>$\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | % MeHg | Reference            |
|-------------------------------------|------|---|--------------------|--------|----|--|--|--------|----------------------|
| QC: La Grande-2 Reservoir (Dike 14) | 1992 | Coleoptera<br>Dytiscidae                  | Larvae/<br>Pupae   | Whole  | 12 | 0.621 dw   | 0.473 dw   | 76%    | Tremblay et al. 1996 |
| QC: La Grande-2 Reservoir (Dike 14) | 1992 | Corixidae<br><i>Sigara</i> spp.           | Larvae/<br>Pupae   | Whole  | 14 | 0.551 dw   | NR   | NR     | Tremblay et al. 1996 |
| QC: La Grande-2 Reservoir (Dike 14) | 1992 | Ephemeroptera<br><i>Leptophlebia</i> spp. | Larvae/<br>Pupae   | Whole  | 32 | 0.139 dw   | 0.069 dw   | 49%    | Tremblay et al. 1996 |
| QC: La Grande-2 Reservoir (Dike 14) | 1992 | Trichoptera<br><i>Agrypnia</i> spp.       | Larvae/<br>Pupae   | Whole  | 4  | 0.292 dw   | 0.059 dw   | 20%    | Tremblay et al. 1996 |
| QC: La Grande-2 Reservoir North arm | 1992 | Chironomids                               | Larvae/<br>Pupae   | Whole  | 8  | 0.285 dw   | 0.073 dw   | 26%    | Tremblay et al. 1996 |
| QC: La Grande-2 Reservoir North arm | 1992 | Corixidae<br><i>Sigara</i> spp.           | Larvae/<br>Pupae   | Whole  | 13 | 1.371 dw   | 1.043 dw   | 76%    | Tremblay et al. 1996 |
| QC: La Grande-2 Reservoir North arm | 1992 | Odonata<br><i>Somatochlora</i> spp.       | Larvae/<br>Pupae   | Whole  | 5  | 0.283 dw   | 0.211 dw   | 75%    | Tremblay et al. 1996 |
| QC: La Grande-2 Reservoir North arm | 1992 | Trichoptera<br><i>Asynarchus</i> spp.     | Larvae/<br>Pupae   | Whole  | 19 | 0.330 dw   | 0.081 dw   | 24%    | Tremblay et al. 1996 |
| QC: La Grande-2 Reservoir South arm | 1992 | Chironomids                               | Larvae/<br>Pupae   | Whole  | 66 | 0.501 dw   | 0.076 dw   | 15%    | Tremblay et al. 1996 |
| QC: La Grande-2 Reservoir South arm | 1992 | Corixidae<br><i>Sigara</i> spp.           | Larvae/<br>Pupae   | Whole  | 44 | 1.675 dw   | 1.519 dw   | 91%    | Tremblay et al. 1996 |

Canadian Tissue Residue Guidelines: Methylmercury

| Location                            | Year | Species  | Life Stage or Size | Tissue | N  | Total Hg<br>$\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | MeHg<br>$\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | % MeHg | Reference            |
|-------------------------------------|------|--|--------------------|--------|----|--|--|--------|----------------------|
| QC: La Grande-2 Reservoir South arm | 1992 | Ephemeroptea<br><i>Leptophlebia</i> spp.                   | Larvae/<br>Pupae   | Whole  | 7  | 0.224 dw   | 0.081 dw   | 36%    | Tremblay et al. 1996 |
| QC: La Grande-2 Reservoir South arm | 1992 | Odonata<br><i>Somatochlora</i> spp.                        | Larvae/<br>Pupae   | Whole  | 3  | 0.615 dw   | 0.615 dw   | 100%   | Tremblay et al. 1996 |
| QC: La Grande-2 Reservoir South arm | 1992 | Trichoptera<br><i>Asynarchus</i> spp.                      | Larvae/<br>Pupae   | Whole  | 10 | 0.544 dw   | 0.106 dw   | 20%    | Tremblay et al. 1996 |
| SK: Buffalo Pound L.                | 1985 | Freshwater mussel<br><i>Pyganodon grandis</i> <sup>y</sup> | Adult              | Muscle | 12 | $0.20 \pm 0.07$  | NR   | NR     | Hammer et al. 1988   |

NR = Not Reported

Notes:

<sup>a</sup> Unless noted otherwise, i.e., dw = dry weight.

<sup>b</sup> Non-zooplankton debris was not separated from bulk samples.

<sup>c</sup> Formerly *Anodonta grandis grandis*

<sup>d</sup> Authors assumed the majority of THg to be MeHg.

**APPENDIX B-2B: LEVELS OF TOTAL MERCURY AND METHYLMERCURY IN MARINE INVERTEBRATES IN THE CANADIAN ENVIRONMENT.**

| Location                       | Year      | Species                         | Life Stage or size | Tissue      | N  | Total Hg $\mu\text{g}\cdot\text{g}^{-1}$ ww | MeHg $\mu\text{g}\cdot\text{g}^{-1}$ ww | % MeHg | Reference              |
|--------------------------------|-----------|---------------------------------|--------------------|-------------|----|---|---|--------|------------------------|
| QC: Saguenay Fjord, mouth      | 1976-1977 | Mussel<br><i>Mytilus edulis</i> | Adult              | Soft tissue | NR | 0.629<br>0.165                              | NR                                      | NR     | Cossa and Rondeau 1985 |
| QC: South shore, lower estuary | 1976-1977 | Mussel<br><i>Mytilus edulis</i> | Adult              | Soft tissue | NR | $0.459 \pm 0.041$<br>$0.267 \pm 0.034$      | NR                                      | NR     | Cossa and Rondeau 1985 |
| QC: Gaspé Peninsula            | 1976-1977 | Mussel<br><i>Mytilus edulis</i> | Adult              | Soft tissue | NR | $0.301 \pm 0.071$<br>$0.120 \pm 0.015$      | NR                                      | NR     | Cossa and Rondeau 1985 |

## APPENDIX B-3A: LEVELS OF TOTAL MERCURY AND METHYL MERCURY IN FRESHWATER FISH IN THE CANADIAN ENVIRONMENT.

| Location  | Year | Species  | Life Stage, Age or Size | Sex | Tissue | N  | Total Hg $\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | MeHg $\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | Other information       | Reference                 |
|---|------|--|-------------------------|-----|--------|----|---|---|-------------------------|---------------------------|
| AB: Giauque L. 63° 10 N                           | 1977 | Lake trout<br><i>Salvelinus namaycush</i>                    | Adult 10-22 years       | NR  | Muscle | 31 | 3.79 mean<br>1.38 to 12.30                                    | NR  | Contaminated            | Moore and Sutherland 1980 |
| AB: Giauque L. 63° 10 N                           | 1977 | Northern pike<br><i>Esox lucius</i>                          | Adult 8-22 years        | NR  | Muscle | 27 | 1.75 mean<br>0.55 to 4.80                                     | NR  | Contaminated            | Moore and Sutherland 1980 |
| AB: Giauque L. 63° 10 N                           | 1977 | Round whitefish<br><i>Prosopium cylindraceum</i>             | Adult 9-20 years        | NR  | Muscle | 35 | 1.22 mean<br>0.44 to 2.64                                     | NR  | Contaminated            | Moore and Sutherland 1980 |
| AB: Hidden L., south of Thompson L.               | 1978 | Lake whitefish<br><i>Coregonus clupeaformis</i>              | Adult 4-17 years        | NR  | Muscle | 13 | 0.20 mean<br>0.02 to 0.39                                     | NR  | Control for Thompson L. | Moore and Sutherland 1980 |
| AB: North Saskatchewan R., downstream of Edmonton | 1982 | Goldeye<br><i>Hiodon alosoides</i>                           | NR                      | NR  | Muscle | 69 | 0.594 ± 0.074<br>(95% CI)<br>0.294 to 1.766                   | NR  | 93% MeHg                | Ramamoorthy et al. 1985   |
| AB: North Saskatchewan R., downstream of Edmonton | 1982 | Longnose sucker<br><i>Catostomus catostomus</i>              | NR                      | NR  | Muscle | 46 | 0.245 ± 0.029<br>(95% CI)<br>0.062 to 0.548                   | NR  | 77% MeHg                | Ramamoorthy et al. 1985   |
| AB: North Saskatchewan R., downstream of Edmonton | 1982 | Northern pike<br><i>Esox lucius</i>                          | NR                      | NR  | Muscle | 43 | 0.314 ± 0.066<br>(95% CI)<br>0.044 to 1.039                   | NR  | 90% MeHg                | Ramamoorthy et al. 1985   |
| AB: North Saskatchewan R., downstream of Edmonton | 1982 | Northern redbhorse sucker<br><i>Moxostoma macrolepidotum</i> | NR                      | NR  | Muscle | 23 | 0.368 ± 0.047<br>(95% CI)<br>0.212 to 0.619                   | NR  | 87% MeHg                | Ramamoorthy et al. 1985   |

## Canadian Tissue Residue Guidelines: Methylmercury

| Location                                       | Year      | Species                                      | Life Stage, Age or Size | Sex | Tissue | N  | Total Hg $\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | MeHg $\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | Other information | Reference                 |
|--|-----------|--|-------------------------|-----|--------|----|---|---|-------------------|---------------------------|
| AB: N. Saskatchewan R., downstream of Edmonton | 1982      | Sauger <i>Stizostedion canadense</i>         | NR                      | NR  | Muscle | 26 | 0.743 ± 0.094 (95% CI)<br>0.271 to 1.585                      | NR  | 92% MeHg          | Ramamoorthy et al. 1985   |
| AB: N. Saskatchewan R. downstream of Edmonton  | 1982      | Walleye <i>Stizostedion vitreum</i>          | NR                      | NR  | Muscle | 35 | 0.645 ± 0.104 (95% CI)<br>0.140 to 1.202                      | NR  | 93% MeHg          | Ramamoorthy et al. 1985   |
| AB: N. Saskatchewan R., downstream of Edmonton | 1982      | White sucker <i>Catostomus commersoni</i>    | NR                      | NR  | Muscle | 27 | 0.313 ± 0.061 (95% CI)<br>0.028 to 0.659                      | NR  | 87% MeHg          | Ramamoorthy et al. 1985   |
| AB: N. Saskatchewan R., upstream of Edmonton   | 1982      | Goldeye <i>Hiodon alosoides</i>              | NR                      | NR  | Muscle | 10 | 0.452 ± 0.072 (95% CI)<br>0.295 to 0.633                      | NR  | 93% MeHg          | Ramamoorthy et al. 1985   |
| AB: N. Saskatchewan R., upstream of Edmonton   | 1982      | Walleye <i>Stizostedion vitreum</i>          | NR                      | NR  | Muscle | 8  | 0.369 ± 0.112 (95% CI)<br>0.228 to 0.625                      | NR  | 95% MeHg          | Ramamoorthy et al. 1985   |
| AB: Thistlethwaite, Unnamed Lakes              | 1977-1978 | Lake trout <i>Salvelinus namaycush</i>       | Adult 5-19 years        | NR  | Muscle | 32 | 0.11 to 1.00  | NR  | Controls          | Moore and Sutherland 1980 |
| AB: Thistlethwaite, Unnamed, Hidden Lakes      | 1977-1978 | Northern pike <i>Esox lucius</i>             | Adult 12-26 years       | NR  | Muscle | 22 | 0.05 to 1.91  | NR  | Controls          | Moore and Sutherland 1980 |
| AB: Thompson L. 62° 37 N                       | 1978      | Lake whitefish <i>Coregonus clupeaformis</i> | Adult 3-15 years        | NR  | Muscle | 33 | 0.20 mean<br>0.03 to 0.62                                     | NR  | Lake Contaminated | Moore and Sutherland 1980 |
| AB: Thompson L. 62° 37 N                       | 1978      | Northern pike <i>Esox lucius</i>             | Adult 5-15 years        | NR  | Muscle | 16 | 1.69 mean<br>1.02 to 3.11                                     | NR  | Contaminated      | Moore and Sutherland 1980 |

## Canadian Tissue Residue Guidelines: Methylmercury

| Location        | Year      | Species  | Life Stage, Age or Size | Sex | Tissue | N  | Total Hg $\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | MeHg $\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | Other information | Reference                         |
|-----------------|-----------|--|-------------------------|-----|--------|----|---|---|-------------------|-----------------------------------|
| BC: Kamloops L. | 1995      | Burbot<br><i>Lota lota</i>                         | NR                      | NR  | Liver  | 5  | < 0.05  | 0.01<br>0.01 to<br>0.02                                   | --                | McDonald et al. 1999 <sup>c</sup> |
| BC: Lakes (5)   | 1982-1987 | Arctic grayling<br><i>Thymallus arcticus</i>       | Adult                   | NR  | Liver  | 1  | 0.08  | NR  | --                | Rieberger 1992                    |
| BC: Lakes (5)   | 1982-1987 | Arctic grayling<br><i>Thymallus arcticus</i>       | Adult                   | NR  | Muscle | 21 | 0.08 $\pm$ 0.04 S.D   | NR  | --                | Rieberger 1992                    |
| BC: Lakes (11)  | 1982-1987 | Cutthroat trout<br><i>Salmo clarki</i>             | Adult                   | NR  | Liver  | 13 | 0.35 $\pm$ 0.25 S.D.  | NR  | --                | Rieberger 1992                    |
| BC: Lakes (11)  | 1982-1987 | Cutthroat trout<br><i>Salmo clarki</i>             | Adult                   | NR  | Muscle | 54 | 0.29 $\pm$ 0.25 S.D.  | NR  | --                | Rieberger 1992                    |
| BC: Lakes (12)  | 1982-1987 | Dolly Varden Char<br><i>Salvelinus malma</i>       | Adult                   | Nr  | Liver  | 9  | 0.11 $\pm$ 0.12 S.D.  | NR  | --                | Rieberger 1992                    |
| BC: Lakes (12)  | 1982-1987 | Dolly Varden Char<br><i>Salvelinus malma</i>       | Adult                   | NR  | Muscle | 46 | 0.15 $\pm$ 0.22 S.D.  | NR  | --                | Rieberger 1992                    |
| BC: Lakes (5)   | 1982-1987 | Lake trout<br><i>Salvelinus namaycush</i>          | Adult                   | NR  | Liver  | 3  | 0.09 $\pm$ 0.04 S.D.  | NR  | --                | Rieberger 1992                    |
| BC: Lakes (5)   | 1982-1987 | Lake trout<br><i>Salvelinus namaycush</i>          | Adult                   | NR  | Muscle | 25 | 0.26 $\pm$ 0.26 S.D.  | NR  | --                | Rieberger 1992                    |
| BC: Lakes (4)   | 1982-1987 | Mountain whitefish<br><i>Prosopium williamsoni</i> | Adult                   | NR  | Liver  | 3  | 0.12 $\pm$ 0.02 S.D.  | NR  | --                | Rieberger 1992                    |
| BC: Lakes (4)   | 1982-1987 | Mountain whitefish<br><i>Prosopium williamsoni</i> | Adult                   | NR  | Muscle | 15 | 0.11 $\pm$ 0.09 S.D.  | NR  | --                | Rieberger 1992                    |

## Canadian Tissue Residue Guidelines: Methylmercury

| Location                | Year           | Species   | Life Stage, Age or Size | Sex | Tissue          | N   | Total Hg $\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | MeHg $\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | Other information | Reference                         |
|-------------------------|----------------|---|-------------------------|-----|-----------------|-----|---|---|-------------------|-----------------------------------|
| BC: Lakes (24)          | 1982-1987      | Rainbow trout<br><i>Oncorhynchus mykiss</i>     | Adult                   | NR  | Liver           | 17  | 0.11 ± 0.11<br>S.D.   | NR  | --                | Rieberger 1992                    |
| BC: Lakes (24)          | 1982-1987      | Rainbow trout<br><i>Oncorhynchus mykiss</i>     | Adult                   | NR  | Muscle          | 97  | 0.09 ± 0.05<br>S.D.   | NR  | --                | Rieberger 1992                    |
| BC: Moose L.            | 1995           | Burbot<br><i>Lota lota</i>                      | NR                      | NR  | Liver           | 5   | 0.03<br>0.02 to 0.08  | 0.04<br>< 0.05 to 0.12                                    | --                | McDonald et al. 1999 <sup>c</sup> |
| BC: Nicola L            | 1995           | Burbot<br><i>Lota lota</i>                      | NR                      | NR  | Liver           | 5   | 0.04<br>< 0.05 to 0.08  | 0.03<br>< 0.05 to 0.07                                    | --                | McDonald et al. 1999              |
| BC: Northern lakes (5)  | 1970s          | Burbot<br><i>Lota lota</i>                      | 51 – 71<br>cm           | NR  | Muscle or Whole | Var | 0.25 to 0.46<br>range of means                                | NR  | Pinchi Fault area | Watson 1992                       |
| BC: Northern lakes (3)  | 1970s          | Kokanee<br><i>Oncorhynchus nerka kennerlyi</i>  | 1000–3500 g             | NR  | Muscle or Whole | Var | 0.03 to 0.07<br>range of means                                | NR  | Pinchi Fault area | Watson 1992                       |
| BC: Northern lakes (14) | 1970s<br>1980s | Lake trout<br><i>Salvelinus namaycush</i>       | 28 – 99<br>cm           | NR  | Muscle or Whole | Var | 0.17 to 0.63<br>range of means                                | NR  | Pinchi Fault area | Watson 1992                       |
| BC: Northern lakes (10) | 1970s<br>1980s | Rainbow trout<br><i>Oncorhynchus mykiss</i>     | 23 – 77<br>cm           | NR  | Muscle or Whole | Var | 0.10 to 0.27<br>range of means                                | NR  | Pinchi Fault area | Watson 1992                       |
| BC: Northern rivers     | 1970s<br>1980s | Bull trout<br><i>Salvelinus confluentus</i>     | 20 – 85<br>cm           | NR  | Muscle or Whole | Var | 0.02 to 0.78<br>range of means                                | NR  | --                | Watson 1992                       |
| BC: Northern rivers     | 1970s<br>1980s | Lake whitefish<br><i>Coregonus clupeaformis</i> | 32 – 51<br>cm           | NR  | Muscle or Whole | 21  | 0.07 to 0.26<br>range of means                                | NR  | --                | Watson 1992                       |
| BC: Northern rivers     | 1970s<br>1980s | Rainbow trout<br><i>Oncorhynchus mykiss</i>     | 15 – 42<br>cm           | NR  | Muscle or Whole | 37  | 0.03 to 0.12<br>range of means                                | NR  | --                | Watson 1992                       |



## Canadian Tissue Residue Guidelines: Methylmercury

| Location                             | Year          | Species   | Life Stage, Age or Size | Sex | Tissue          | N   | Total Hg $\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | MeHg $\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | Other information | Reference            |
|--------------------------------------|---------------|---|-------------------------|-----|-----------------|-----|---|---|-------------------|----------------------|
| BC: Pinchi L.                        | 1986          | Bull trout<br><i>Salvelinus confluentus</i>     | 30 – 36 cm              | NR  | Muscle or Whole | 3   | 0.65 mean<br>0.49 to 0.75                                     | NR  | High natural Hg   | Watson 1992          |
| BC: Pinchi L.                        | 1975          | Kokanee<br><i>Oncorhynchus nerka kennerlyi</i>  | 22 – 26 cm              | NR  | Muscle or Whole | 6   | 0.48 mean<br>0.32 to 0.76                                     | NR  | High natural Hg   | Watson 1992          |
| BC: Pinchi L.                        | 1970s<br>1986 | Lake trout<br><i>Salvelinus namaycush</i>       | 24 – 72 cm              | NR  | Muscle or Whole | Var | 1.06 to 5.78<br>range of means                                | NR  | High natural Hg   | Watson 1992          |
| BC: Pinchi L.                        | 1975-<br>1986 | Lake whitefish<br><i>Coregonus clupeaformis</i> | 26 – 46 cm              | NR  | Muscle or Whole | Var | 0.65 to 0.90<br>range of means                                | NR  | High natural Hg   | Watson 1992          |
| BC: Pinchi L.                        | 1970s<br>1986 | Rainbow trout<br><i>Oncorhynchus mykiss</i>     | 14 – 32 cm              | NR  | Muscle or Whole | 16  | 0.20 to 0.40<br>range of means                                | NR  | High natural Hg   | Watson 1992          |
| BC: Stuart L.                        | 1995          | Burbot<br><i>Lota lota</i>                      | NR                      | NR  | Liver           | 5   | 0.04<br>< 0.05 to 0.10  | 0.03<br>0.02 to 0.09                                      | --                | McDonald et al. 1999 |
| BC: Williston L., behind Bennett Dam | 1980s         | Bull trout<br><i>Salvelinus confluentus</i>     | 20 – 88 cm              | NR  | Muscle or Whole | Var | 0.25 to 1.62<br>range of site means                           | NR  | --                | Watson 1992          |
| BC: Williston L., behind Bennett Dam | 1980s         | Burbot<br><i>Lota lota</i>                      | 24 – 83 cm              | NR  | Muscle or Whole | Var | 0.26 to 0.33<br>range of site means                           | NR  | --                | Watson 1992          |
| BC: Williston L., behind Bennett Dam | 1980s         | Kokanee<br><i>Oncorhynchus nerka kennerlyi</i>  | 20 – 31 cm              | NR  | Muscle or Whole | Var | 0.04 to 0.31<br>range of site means                           | NR  | --                | Watson 1992          |
| BC: Williston L., behind Bennett Dam | 1980          | Lake trout<br><i>Salvelinus namaycush</i>       | 39 – 49 cm              | NR  | Muscle or Whole | Var | 0.24 to 0.32<br>range of site means                           | NR  | --                | Watson 1992          |

## Canadian Tissue Residue Guidelines: Methylmercury

| Location   | Year      | Species                                      | Life Stage, Age or Size | Sex | Tissue          | N   | Total Hg $\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | MeHg $\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | Other information         | Reference                  |
|--|-----------|--|-------------------------|-----|-----------------|-----|---|---|---------------------------|----------------------------|
| BC: Williston L., behind Bennett Dam             | 1980s     | Rainbow trout <i>Oncorhynchus mykiss</i>     | 22 – 41 cm              | NR  | Muscle or Whole | Var | 0.04 to 0.08 range of site means                              | NR  | --                        | Watson 1992                |
| MB: Cliff L.                                     | 1976      | Northern pike <i>Esox lucius</i>             | 1-7 years               | NR  | Liver           | 23  | 0.06 - 0.72   | NR  | Near a base metal smelter | McFarlane and Franzin 1980 |
| MB: Footprint L., downstream of Notigi Reservoir | 1978-1981 | Northern pike <i>Esox lucius</i>             | 45.7 – 52.6 cm          | NR  | Muscle          | 58  | 0.81 mean of all site means for 3 years.                      | NR  | Churchill River flooding  | Bodaly et al. 1984         |
| MB: Footprint L., downstream of Notigi Reservoir | 1978-1981 | Walleye <i>Stizostedion vitreum</i>          | 37.6 – 39.4 cm          | NR  | Muscle          | 82  | 0.95 mean of all site means for 3 years.                      | NR  | Churchill River flooding  | Bodaly et al. 1984         |
| MB: Hamell L.                                    | 1976      | Northern pike <i>Esox lucius</i>             | 2-5 years               | NR  | Liver           | 18  | 0.08 to 0.86  | NR  | Near a base metal smelter | McFarlane and Franzin 1980 |
| MB: Isset L., Notigi Reservoir                   | 1975-1982 | Lake whitefish <i>Coregonus clupeaformis</i> | 31.0 – 38.4 cm          | NR  | Muscle          | 54  | 0.23 mean of all site means for 3 years.                      | NR  | Churchill River flooding  | Bodaly et al. 1984         |
| MB: Isset L., Notigi Reservoir                   | 1978-1982 | Northern pike <i>Esox lucius</i>             | 57.3 – 59.6 cm          | NR  | Muscle          | 31  | 0.76 mean of all site means for 2 years.                      | NR  | Churchill River flooding  | Bodaly et al. 1984         |
| MB: Isset L., Notigi Reservoir                   | 1978-1982 | Walleye <i>Stizostedion vitreum</i>          | 34.7 – 38.9 cm          | NR  | Muscle          | 24  | 1.16 mean of all site means for 2 years.                      | NR  | Churchill River flooding  | Bodaly et al. 1984         |
| MB: Mystery L., downstream of Notigi Reservoir   | 1979      | Northern pike <i>Esox lucius</i>             | 54.1 cm (mean)          | NR  | Muscle          | 45  | 0.79 mean of all site means for 1 year.                       | NR  | Churchill River flooding  | Bodaly et al. 1984         |
| MB: Mystery L., downstream of Notigi Reservoir   | 1979      | Walleye <i>Stizostedion vitreum</i>          | 46.2 cm                 | NR  | Muscle          | 33  | 1.13  | NR  | Churchill River flooding  | Bodaly et al. 1984         |

## Canadian Tissue Residue Guidelines: Methylmercury

| Location                                  | Year      | Species   | Life Stage, Age or Size | Sex | Tissue | N   | Total Hg $\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | MeHg $\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | Other information         | Reference                  |
|---|-----------|---|-------------------------|-----|--------|-----|---|---|---------------------------|----------------------------|
| MB: Notigi L., Notigi Reservoir           | 1980-1981 | Lake whitefish<br><i>Coregonus clupeaformis</i> | 30.9 – 42.1 cm          | NR  | Muscle | 44  | 0.18 mean of all site means for 2 years.                      | NR  | Churchill River flooding  | Bodaly et al. 1984         |
| MB: Notigi L., Notigi Reservoir           | 1977-1982 | Northern pike<br><i>Esox lucius</i>             | 58.1 – 78.8 cm          | NR  | Muscle | 68  | 1.83 mean of all site means for 4 years.                      | NR  | Churchill River flooding  | Bodaly et al. 1984         |
| MB: Notigi L., Notigi Reservoir           | 1978-1982 | Walleye<br><i>Stizostedion vitreum</i>          | 41.6 – 50.3 cm          | NR  | Muscle | 62  | 1.89 mean of all site means for 4 years.                      | NR  | Churchill River flooding  | Bodaly et al. 1984         |
| MB: Rat L., Notigi Reservoir              | 1978-1980 | Lake whitefish<br><i>Coregonus clupeaformis</i> | 42.2 – 43.9 cm          | NR  | Muscle | 31  | 0.36 mean of all site means for 4 years.                      | NR  | Churchill River flooding  | Bodaly et al. 1984         |
| MB: Rat L., Notigi Reservoir              | 1978-1980 | Northern pike<br><i>Esox lucius</i>             | 69.8 cm                 | NR  | Muscle | 30  | 2.17 mean of all site means for 2 years.                      | NR  | Churchill River flooding  | Bodaly et al. 1984         |
| MB: Rat L., Notigi Reservoir              | 1978-1980 | Walleye<br><i>Stizostedion vitreum</i>          | 40.8 – 45.0 cm          | NR  | Muscle | 79  | 2.14 mean of all site means for 3 years.                      | NR  | Churchill River flooding  | Bodaly et al. 1984         |
| MB: Southern Indian L.                    | 1975-1882 | Lake whitefish<br><i>Coregonus clupeaformis</i> | 31.6 – 42.3 cm          | NR  | Muscle | 700 | 0.13 mean of all site means for 6 years.                      | NR  | --                        | Bodaly et al. 1984         |
| MB: Southern Indian L.                    | 1975-1882 | Northern pike<br><i>Esox lucius</i>             | 49.7 – 66.6 cm means    | NR  | Muscle | 461 | 0.68 mean of all site means for 4-5 years.                    | NR  | --                        | Bodaly et al. 1984         |
| MB: Southern Indian L.                    | 1975-1882 | Walleye<br><i>Stizostedion vitreum</i>          | 33.3 – 43.5 cm means    | NR  | Muscle | 289 | 0.57 mean of all site means for 4-5 years.                    | NR  | --                        | Bodaly et al. 1984         |
| MB: Thompson L.                           | 1976      | Northern pike<br><i>Esox lucius</i>             | 1-6 years.              | NR  | Liver  | 22  | 0.09 to 0.64  | NR  | Near a base metal smelter | McFarlane and Franzin 1980 |
| MB: Threepoint L., downstream Notigi Res. | 1980-1981 | Lake whitefish<br><i>Coregonus clupeaformis</i> | 33.3 – 42.9 cm means    | NR  | Muscle | 26  | 0.40 mean of all site means for 2 years.                      | NR  | Churchill River flooding  | Bodaly et al. 1984         |

## Canadian Tissue Residue Guidelines: Methylmercury

| Location                                       | Year      | Species   | Life Stage, Age or Size | Sex | Tissue | N   | Total Hg $\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | MeHg $\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | Other information        | Reference          |
|--|-----------|---|-------------------------|-----|--------|-----|---|---|--------------------------|--------------------|
| MB: Threepoint L., downstream Notigi Reservoir | 1980-1981 | Northern pike<br><i>Esox lucius</i>             | 51.7 – 68.8 cm means    | NR  | Muscle | 38  | 1.31 mean of all site means for 2 years.                      | NR  | Churchill River flooding | Bodaly et al. 1984 |
| MB: Threepoint L., downstream Notigi Reservoir | 1980-1981 | Walleye<br><i>Stizostedion vitreum</i>          | 38.1 – 40.3 cm means    | NR  | Muscle | 52  | 1.27 mean of all site means for 2 years.                      | NR  | Churchill River flooding | Bodaly et al. 1984 |
| MB: Wapisu L., downstream Notigi Reservoir     | 1977      | Northern pike<br><i>Esox lucius</i>             | 41.1 cm mean            | NR  | Muscle | 41  | 0.81 mean of all site means for 1 year.                       | NR  | Churchill River flooding | Bodaly et al. 1984 |
| MB: Wapisu L., downstream Notigi Reservoir     | 1977      | Walleye<br><i>Stizostedion vitreum</i>          | 41.1 cm mean            | NR  | Muscle | 94  | 1.25 mean of all site means for 1 year.                       | NR  | Churchill River flooding | Bodaly et al. 1984 |
| MB: Wuskwatim L., downstream Notigi Reservoir  | 1979-1982 | Northern pike<br><i>Esox lucius</i>             | 51.4 – 60.5 cm means    | NR  | Muscle | 108 | 1.01 mean of all site means for 4 years.                      | NR  | Churchill River flooding | Bodaly et al. 1984 |
| MB: Wuskwatim L., downstream Notigi Reservoir  | 1979-1981 | Walleye<br><i>Stizostedion vitreum</i>          | 39.9 – 41.5 cm means    | NR  | Muscle | 143 | 0.88 mean of all site means for 3 years.                      | NR  | Churchill River flooding | Bodaly et al. 1984 |
| NB: St. John R., Green River Reservoir         | 1983      | Lake whitefish<br><i>Coregonus clupeaformis</i> | 20 - 25 cm              | NR  | Muscle | 12  | 0.12 mean<br>0.10 to 0.64                                     | NR  | --                       | Bailey 1985        |
| NB: St. John R., Sisson Branch Reservoir       | 1983      | Land-locked salmon<br><i>Salmo salar</i>        | 38 & 58 cm              | NR  | Muscle | 2   | 0.84 and 1.5  | NR  | --                       | Bailey 1985        |

## Canadian Tissue Residue Guidelines: Methylmercury

| Location                         | Year      | Species  | Life Stage, Age or Size     | Sex       | Tissue | N   | Total Hg $\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | MeHg $\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | Other information        | Reference                               |
|----------------------------------|-----------|--|-----------------------------|-----------|--------|-----|---|---|--------------------------|---|
| NF: Labrador                     | 1977-1978 | Brook trout<br><i>Salvelinus fontinalis</i>          | NR                          | NR        | NR     | 96  | 0.01-1.69   | NR  | --                       | Bruce et al. 1979                       |
| NF: Labrador                     | 1977-1978 | Northern pike<br><i>Esox lucius</i>                  | NR                          | NR        | NR     | 117 | 0.01-3.08   | NR  | --                       | Bruce et al. 1979                       |
| NF: Labrador                     | 1977-1978 | Lake trout<br><i>Salvelinus namaycush</i>            | NR                          | NR        | NR     | 107 | 0.12-3.37   | NR  | --                       | Bruce et al. 1979                       |
| NF: Labrador Smallwood Reservoir | 1977-1978 | Land-locked salmon<br><i>Salmo salar</i>             | NR                          | NR        | Muscle | NR  | 1.5 to 2.3  | NR  | --                       | Bruce and Spencer 1978 c.f. Bailey 1985 |
| NS: Annapolis R.                 | NR        | Striped bass <sup>d</sup><br><i>Morone saxatilis</i> | 7.38 kg mean                | NR        | Muscle | NR  | 0.77 mean<br>0.26 to 1.97                                     | NR  | --                       | Ray et al. 1984                         |
| NS: Wreck Cove Reservoir         | 1994      | Brook trout<br><i>Salvelinus fontinalis</i>          | 49-211 g<br>16.8-26.8 cm    | NR        | Muscle | 12  | $0.495 \pm 0.164$<br>0.27 to 0.73                             | NR  | Oldest fish only 3 years | ADI Nolan Davis Inc. 1994               |
| NS: Shubenacadie R.              | NR        | Striped bass<br><i>Morone saxatilis</i>              | 1.81 kg mean                | NR        | Muscle | NR  | 0.40<br>0.16 to 1.44  | NR  | --                       | Ray et al. 1984                         |
| NWT: Hay R.                      | 1989-1990 | Lake whitefish<br><i>Coregonus clupeaformis</i>      | 362 – 1800g<br>7 – 12 years | 7M<br>33F | Muscle | 40  | 0.07 mean<br>0.03 to 0.13                                     | NR  | --                       | Grey et al. 1995                        |
| NWT: Hay R.                      | 1989-1990 | Northern pike<br><i>Esox lucius</i>                  | 775 – 3100g<br>5 – 13 years | 16M<br>5F | Muscle | 21  | 0.32 mean<br>0.19 to 0.59                                     | NR  | --                       | Grey et al. 1995                        |
| NWT: Hay R.                      | 1989-1990 | Walleye<br><i>Stizostedion vitreum</i>               | 450 – 1800g<br>4 – 11 years | 5M<br>30F | Muscle | 35  | 0.22 mean<br>0.1 to 0.32                                      | NR  | --                       | Grey et al. 1995                        |
| NWT: Johnny Hoe R./ Keller L.    | 1993      | Lake trout<br><i>Salvelinus namaycush</i>            | 8-32 years                  | NR        | Muscle | 15  | 0.41 mean<br>0.22 to 1.05                                     | NR  | --                       | Stephens 1995                           |

## Canadian Tissue Residue Guidelines: Methylmercury

| Location                             | Year      | Species   | Life Stage, Age or Size      | Sex        | Tissue          | N  | Total Hg $\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | MeHg $\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | Other information | Reference             |
|--------------------------------------|-----------|---|------------------------------|------------|-----------------|----|---|---|-------------------|-----------------------|
| NWT: Johnny Hoe R./ Keller L.        | 1993      | Lake whitefish<br><i>Coregonus clupeaformis</i> | 7-19 years                   | NR         | Muscle          | 15 | 0.064   | NR  | --                | Stephens 1995         |
| NWT: Johnny Hoe R./ Lac Ste. Thérèse | 1993      | Lake trout<br><i>Salvelinus namaycush</i>       | 32-39 years                  | NR         | Muscle          | 2  | 1.34  | NR  | Unknown source    | Stephens 1995         |
| NWT: Johnny Hoe R./ Lac Ste. Thérèse | 1993      | Northern pike<br><i>Esox lucius</i>             | 6-11 years                   | NR         | Muscle          | 4  | 0.735 mean<br>0.25 to 1.09                                    | NR  | Unknown source    | Stephens 1995         |
| NWT: Johnny Hoe R./ Lac Ste. Thérèse | 1993      | Walleye<br><i>Stizostedion vitreum</i>          | 9-17 years                   | NR         | Muscle          | 30 | 1.49 mean<br>0.29 to 1.99                                     | NR  | Unknown source    | Stephens 1995         |
| NWT: Johnny Hoe R./ Tseepantee L.    | 1993      | Lake whitefish<br><i>Coregonus clupeaformis</i> | 6-21 years                   | NR         | Muscle          | 15 | 0.102   | NR  | --                | Stephens 1995         |
| NWT: Johnny Hoe R./ Tseepantee L.    | 1993      | Northern pike<br><i>Esox lucius</i>             | 6-8 years                    | NR         | Muscle          | 6  | 0.475 mean<br>0.39 to 0.71                                    | NR  | --                | Stephens 1995         |
| NWT: Johnny Hoe R./ Tseepantee L.    | 1993      | Walleye<br><i>Stizostedion vitreum</i>          | 11-23 years                  | NR         | Muscle          | 15 | 0.926 mean<br>0.25 to 1.42                                    | NR  | --                | Stephens 1995         |
| NWT: Kaminak L., West of Hudson Bay  | 1970-1971 | Lake trout<br><i>Salvelinus namaycush</i>       | Adult                        | NR         | Muscle or Whole | 70 | 0.57 to 2.0   | NR  | High natural Hg   | Shilts and Coker 1995 |
| NWT: Leland L.                       | 1989-1990 | Lake whitefish<br><i>Coregonus clupeaformis</i> | 800 – 1600 g<br>7 – 12 years | 10M<br>10F | Muscle          | 20 | 0.11 mean<br>0.05 to 0.23                                     | NR  | --                | Grey et al. 1995      |
| NWT: Leland L.                       | 1989-1990 | Northern pike<br><i>Esox lucius</i>             | 500 – 2880 g<br>4 – 14 years | 14M<br>19F | Muscle          | 32 | 0.34 mean<br>0.16 to 0.59                                     | NR  | --                | Grey et al. 1995      |

## Canadian Tissue Residue Guidelines: Methylmercury

| Location                       | Year      | Species   | Life Stage, Age or Size      | Sex        | Tissue | N  | Total Hg $\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | MeHg $\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | Other information | Reference                  |
|--------------------------------|-----------|---|------------------------------|------------|--------|----|---|---|-------------------|----------------------------|
| NWT: Leland L.                 | 1989-1990 | Walleye<br><i>Stizostedion vitreum</i>          | 776 – 2450 g<br>8 – 18 years | 11M<br>8F  | Muscle | 19 | 0.46 mean<br>0.25 to 0.73                                     | NR  | --                | Grey et al. 1995           |
| NWT: Slave R.                  | 1988-1990 | Lake whitefish<br><i>Coregonus clupeaformis</i> | 524 – 1025 g<br>6 – 12 years | 17M<br>13F | Muscle | 30 | 0.08 mean<br>0.02 to 0.13                                     | NR  | --                | Grey et al. 1995           |
| NWT: Slave R.                  | 1988-1990 | Northern pike<br><i>Esox lucius</i>             | 700 – 2750 g<br>5 – 15 years | 24M<br>39F | Muscle | 63 | 0.34 mean<br>0.17 to 0.60                                     | NR  | --                | Grey et al. 1995           |
| NWT: Slave R.                  | 1988-1990 | Walleye<br><i>Stizostedion vitreum</i>          | 500 – 1600 g<br>6 – 14 years | 72M<br>27F | Muscle | 99 | 0.34 mean<br>0.13 to 0.80                                     | NR  | --                | Grey et al. 1995           |
| ON: Ball L. North-west Ontario | 1970      | Northern pike<br><i>Esox lucius</i>             | 1529 – 2900 g                | NR         | Muscle | 7  | 7.7 mean<br>1.61 to 20.0                                      | NR  | Contaminated      | Fimreite and Reynolds 1973 |
| ON: Ball L. North-west Ontario | 1970      | Rock bass<br><i>Ambloplites rupestris</i>       | 650 – 1685 g                 | NR         | Muscle | 9  | 6.2 mean<br>1.14 to 10.9                                      | NR  | Contaminated      | Fimreite and Reynolds 1973 |
| ON: Ball L. North-west Ontario | 1970      | Walleye<br><i>Stizostedion vitreum</i>          | 410 – 1550 g                 | NR         | Muscle | 9  | 4.5 mean<br>0.5 to 8.7  | NR  | Contaminated      | Fimreite and Reynolds 1973 |
| ON: Clay L. North-west Ontario | 1970      | Burbot<br><i>Lota lota</i>                      | 1355 – 2150 g                | NR         | Muscle | 4  | 21.9 mean<br>19.1 to 24.8                                     | NR  | Contaminated      | Fimreite and Reynolds 1973 |
| ON: Clay L. North-west Ontario | 1970      | Walleye<br><i>Stizostedion vitreum</i>          | 1025 – 2040 g                | NR         | Muscle | 5  | 15.7 mean<br>12.3 to 19.6                                     | NR  | Contaminated      | Fimreite and Reynolds 1973 |
| ON: Clay L. North-west Ontario | 1970      | White sucker<br><i>Catostomus commersoni</i>    | 575 – 1050 g                 | NR         | Muscle | 5  | 3.13 mean<br>2.3 to 3.8                                       | NR  | Contaminated      | Fimreite and Reynolds 1973 |

## Canadian Tissue Residue Guidelines: Methylmercury

| Location   | Year      | Species                                   | Life Stage, Age or Size | Sex | Tissue          | N   | Total Hg $\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | MeHg $\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | Other information                          | Reference                  |
|--|-----------|---|-------------------------|-----|-----------------|-----|---|---|--|----------------------------|
| ON: L. Ontario   | 1977-1988 | Lake trout<br><i>Salvelinus namaycush</i> | 1 – 8 years             | NR  | Whole           | 134 | 0.20 mean of all sites<br>0.18 to 0.23                        | NR  | [Hg] increased with weight                 | Borgmann and Whittle 1991  |
| ON: L. Simcoe  | 1982      | Northern pike<br><i>Esox lucius</i>       | 1 - 11 years            | NR  | Muscle          | 43  | Max of 0.85   | NR  | --   | Mathers and Johansen 1985  |
| ON: L. Simcoe  | 1982      | Walleye<br><i>Stizostedion vitreum</i>    | 1 - 16 years            | NR  | Muscle          | 46  | Max of 2.7  | NR  | --   | Mathers and Johansen 1985  |
| ON: Lakes  | NR        | Lake trout<br><i>Salvelinus namaycush</i> | 44 cm                   | NR  | Muscle          | 41  | 0.303 mean<br>0.070 to 1.033                                  | NR  | --   | Wren 1991                  |
| ON: Lakes (3)<br>Wabigoon/English R. area,<br>North-west Ontario | 1970      | Lake trout<br><i>Salvelinus namaycush</i> | 1195 – 2400 g           | NR  | Muscle          | 13  | 0.52 to 0.82<br>range of means<br>all lakes                   | NR  | Control lakes                              | Fimreite and Reynolds 1973 |
| ON: Lakes (28)<br>from 44°53' to 49°37' Lat.                     | 1975-1984 | Lake trout<br><i>Salvelinus namaycush</i> | 1213 g<br>(geom. mean)  | NR  | Whole or muscle | 569 | 0.47 mean of all lake means                                   | NR  | With pelagic forage fish, no <i>Mysis</i>  | Cabana et al. 1994         |
| ON: Lakes (33)<br>from 44°53' to 49°37' Lat.                     | 1975-1984 | Lake trout<br><i>Salvelinus namaycush</i> | 1615 g<br>(geom. mean)  | NR  | Whole or muscle | 851 | 0.65 mean of all lake means                                   | NR  | With pelagic forage fish and <i>Mysis</i>  | Cabana et al. 1994         |
| ON: Lakes (35)<br>from 45°01' to 49°10' Lat.                     | 1975-1984 | Lake trout<br><i>Salvelinus namaycush</i> | 686 g<br>(geom. mean)   | NR  | whole or muscle | 885 | 0.18 mean of all lake means                                   | NR  | With <i>Mysis</i> , no pelagic forage fish | Cabana et al. 1994         |
| ON: Lakes<br>Koshlong,<br>Rosseau,<br>Tadenac                    | 1977-1981 | Lake trout<br><i>Salvelinus namaycush</i> | 60 cm                   | NR  | Muscle          | 84  | 0.61 to 0.99 means  | NR  | Rated moderate [Hg]                        | MacCrimmon et al. 1983     |
| ON: Lakes<br>Muskoka,<br>Mary, Vernon                            | 1977-1981 | Lake trout<br><i>Salvelinus namaycush</i> | 60 cm                   | NR  | Muscle          | 84  | 2.96 to 4.49 means  | NR  | Rated high [Hg]                            | MacCrimmon et al. 1983     |



## Canadian Tissue Residue Guidelines: Methylmercury

| Location   | Year      | Species                                     | Life Stage, Age or Size | Sex | Tissue             | N                   | Total Hg $\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | MeHg $\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | Other information                      | Reference                  |
|--|-----------|---|-------------------------|-----|--------------------|---------------------|---|---|--|----------------------------|
| ON: Lakes Simcoe, Joseph, Bella                            | 1977-1981 | Lake trout <i>Salvelinus namaycush</i>      | 60 cm                   | NR  | Muscle             | 85                  | 0.39 to 0.55 means  | NR  | Rated low [Hg]                         | MacCrimmon et al. 1983     |
| ON: Lakes (7) Wabigoon/English R. area, North-west Ontario | 1970      | Northern pike <i>Esox lucius</i>            | 580 – 4370 g            | NR  | Muscle             | 27                  | 0.52 to 1.3 range of means all lakes                          | NR  | Control lakes                          | Fimreite and Reynolds 1973 |
| ON: Lakes (170) Province-wide                              | 1978-1984 | Northern pike <i>Esox lucius</i>            | Adult ~54 cm            | NR  | Muscle             | 10+ PL <sup>e</sup> | 0.50 mean 0.07 to 1.28  | NR  | --                                     | Wren et al. 1991           |
| ON: Lakes (16) South-central Ontario                       | 1981      | Pumpkinseed sunfish <i>Lepomis gibbosus</i> | Adult                   | NR  | Muscle             | 20 <sup>f</sup>     | 0.01 to 0.54  | NR  | --                                     | Wren and MacCrimmon 1983   |
| ON: Lakes Koshlong, Rosseau, Tadenac                       | 1977-1981 | Rainbow smelt <i>Osmerus mordax</i>         | 15 cm                   | NR  | Muscle             | 22                  | 0.18 to 0.26 means  | NR  | Rated moderate [Hg] <sup>g</sup>       | MacCrimmon et al. 1983     |
| ON: Lakes Muskoka, Mary, Vernon                            | 1977-1981 | Rainbow smelt <i>Osmerus mordax</i>         | 15 cm                   | NR  | Muscle             | 20                  | 0.34 to 0.49 means  | NR  | Rated high [Hg] <sup>g</sup>           | MacCrimmon et al. 1983     |
| ON: Lakes Simcoe, Joseph, Bella                            | 1977-1981 | Rainbow smelt <i>Osmerus mordax</i>         | 15 cm                   | NR  | Muscle             | 30                  | 0.08 to 0.14 means  | NR  | Rated low [Hg]                         | MacCrimmon et al. 1983     |
| ON: Lakes  | NR        | Smallmouth bass <i>Micropterus dolomieu</i> | 31 cm                   | NR  | Muscle             | NR                  | 0.402 mean 0.132 to 0.943                                     | NR  | --                                     | Wren 1991                  |
| ON: Lakes (9) from 44°55' to 45°24' Lat.                   | 1975-1984 | Smallmouth bass <i>Micropterus dolomieu</i> | 305 g (geom. mean)      | NR  | NR Whole or muscle | 179                 | 0.42 mean of all lake means                                   | NR  | No pelagic forage fish or <i>Mysis</i> | Cabana et al. 1994         |

## Canadian Tissue Residue Guidelines: Methylmercury

| Location   | Year      | Species  | Life Stage, Age or Size | Sex | Tissue             | N      | Total Hg $\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | MeHg $\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | Other information                         | Reference                  |
|--|-----------|--|-------------------------|-----|--------------------|--------|---|---|---|----------------------------|
| ON: Lakes (17) from 44°25' to 47°58' Lat.                  | 1975-1984 | Smallmouth bass<br><i>Micropterus dolomieu</i> | 508 g (geom. mean)      | NR  | NR Whole or muscle | 337    | 0.43 mean of all lake means                                   | NR  | With pelagic forage fish, no <i>Mysis</i> | Cabana et al. 1994         |
| ON: Lakes (30) from 44°50' to 50°02' Lat.                  | 1975-1984 | Smallmouth bass<br><i>Micropterus dolomieu</i> | 489 g (geom. mean)      | NR  | NR Whole or muscle | 605    | 0.53 mean of all lake means                                   | NR  | With pelagic forage fish and <i>Mysis</i> | Cabana et al. 1994         |
| ON: Lakes  | NR        | Walleye<br><i>Stizostedion vitreum</i>         | 41 cm                   | NR  | Muscle             | 44     | 0.517 mean<br>0.128 to 2.216                                  | NR  | --  | Wren 1991                  |
| ON: Lakes (9) Wabigoon/English R. area, North-west Ontario | 1970      | Walleye<br><i>Stizostedion vitreum</i>         | 360 – 2420 g            | NR  | Muscle             | 49     | 0.31 to 0.76 range of means all lakes                         | NR  | Control lakes                             | Fimreite and Reynolds 1973 |
| ON: Lakes (255) province-wide                              | 1978-1984 | Walleye<br><i>Stizostedion vitreum</i>         | Adult ~41 cm            | NR  | Muscle             | 10+ PL | 0.58 mean<br>0.09 to 3.24                                     | NR  | --  | Wren et al. 1991           |
| ON: Lakes(6) North-west Ontario- remote                    | 1987-1989 | Yellow Perch<br><i>Perca flavescens</i>        | Age 1+ 7 – 8.1 cm       | NR  | Muscle             | 246    | 0.04 to 0.14  | NR  | Inverse relation with lake size           | Bodaly et al. 1993         |
| ON: Lakes (16) Muskoka-Haliburton area                     | 1978-1987 | Yellow perch<br><i>Perca flavescens</i>        | Yearling                | NR  | Whole              | <11 PL | 0.031 to 0.233 range of lake means                            | NR  | --  | Suns and Hitchin 1990      |
| ON: Ottawa R.  | 1974      | Yellow Perch<br><i>Perca flavescens</i>        | Yearling                | NR  | Whole              | 32     | 0.13 to 0.17  | NR  | 68 to 74% organic Hg                      | Rodgers and Qadri 1982     |
| ON: Ottawa R.  | 1975      | Yellow Perch<br><i>Perca flavescens</i>        | Yearling                | NR  | Whole              | 16     | 0.13 ± 0.005  | NR  | 83% organic Hg                            | Rodgers and Qadri 1982     |

## Canadian Tissue Residue Guidelines: Methylmercury

| Location   | Year      | Species                                      | Life Stage, Age or Size | Sex | Tissue | N   | Total Hg $\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^a$ | MeHg $\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^a$ | Other information | Reference                  |
|--|-----------|--|-------------------------|-----|--------|-----|--|--|-------------------|----------------------------|
| ON: Tide-Indian-Grassy Narrows L. North-west Ontario | 1970      | Walleye <i>Stizostedion vitreum</i>          | 715 – 4220 g            | NR  | Muscle | 18  | 2.7 to 3.4 range of means of 3 lakes                 | NR   | Contaminated      | Fimreite and Reynolds 1973 |
| ON: Wabigoon R. North-west Ontario                   | 1970      | Northern pike <i>Esox lucius</i>             | 510 – 2700 g            | NR  | Muscle | 3   | 15.2 mean<br>8.6 to 27.8                             | NR   | Contaminated      | Fimreite and Reynolds 1973 |
| ON: Wabigoon R. North-west Ontario                   | 1970      | Walleye <i>Stizostedion vitreum</i>          | 160 – 1490 g            | NR  | Muscle | 4   | 6.8 mean<br>0.5 to 10.4                              | NR   | Contaminated      | Fimreite and Reynolds 1973 |
| ON: Wabigoon R. North-west Ontario                   | 1970      | White sucker <i>Catostomus commersoni</i>    | 570 – 1060 g            | NR  | Muscle | 3   | 4.2 mean<br>0.6 to 8.9                               | NR   | Contaminated      | Fimreite and Reynolds 1973 |
| QC: Great Whale area, Northern Québec                | 1989-1990 | Northern pike <i>Esox lucius</i>             | NR                      | NR  | Muscle | 75  | 0.63 ± 0.46  | NR   | --                | Langlois et al. 1995       |
| QC: Great Whale area, Northern Québec                | 1989-1990 | Lake trout <i>Salvelinus namaycush</i>       | NR                      | NR  | Muscle | 504 | 0.77 ± 0.66  | NR   | --                | Langlois et al. 1995       |
| QC: La Grande- 2 Reservoir                           | 1982-1988 | Lake whitefish <i>Coregonus clupeaformis</i> | 40 cm                   | NR  | Muscle | NR  | 0.48 to 0.57 – High value after 5 years.             | NR   | Flooded 1978/9    | Verdon et al. 1991         |
| QC: La Grande- 2 Reservoir                           | 1982-1988 | Longnose sucker <i>Catostomus catostomus</i> | 40 cm                   | NR  | Muscle | NR  | 0.41 to 0.67 – High value after 5 years.             | NR   | Flooded 1978/9    | Verdon et al. 1991         |
| QC: La Grande -2 Reservoir                           | 1982-1988 | Northern pike <i>Esox lucius</i>             | 70 cm                   | NR  | Muscle | NR  | 1.31 to 2.99 – High value after 9 years.             | NR   | Flooded 1978/9    | Verdon et al. 1991         |

## Canadian Tissue Residue Guidelines: Methylmercury

| Location   | Year      | Species                                      | Life Stage, Age or Size        | Sex | Tissue | N   | Total Hg $\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | MeHg $\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | Other information                       | Reference            |
|--|-----------|--|--------------------------------|-----|--------|-----|---|---|---|----------------------|
| QC: La Grande- 2 Reservoir                                     | 1982-1988 | Walleye <i>Stizostedion vitreum</i>          | 40 cm                          | NR  | Muscle | NR  | 1.92 to 2.80 – High level after 9 years.                      | NR  | Flooded 1978/9                          | Verdon et al. 1991   |
| QC: Lakes (29), near La Grande complex                         | 1978-1988 | Lake whitefish <i>Coregonus clupeaformis</i> | 40 cm                          | NR  | Muscle | NR  | 0.07 to 0.30  | NR  | Control                                 | Verdon et al. 1991   |
| QC: Lakes (29), near La Grande complex                         | 1978-1988 | Longnose sucker <i>Catostomus catostomus</i> | 40 cm                          | NR  | Muscle | NR  | 0.06 to 0.32  | NR  | Control                                 | Verdon et al. 1991   |
| QC: Lakes (29), near La Grande complex                         | 1978-1988 | Northern pike <i>Esox lucius</i>             | 70 cm                          | NR  | Muscle | NR  | 0.25 to 0.90  | NR  | Control                                 | Verdon et al. 1991   |
| QC: Lakes (29), near La Grande complex                         | 1978-1988 | Walleye <i>Stizostedion vitreum</i>          | 40 cm                          | NR  | Muscle | NR  | 0.32 to 1.26  | NR  | Control                                 | Verdon et al. 1991   |
| QC: Lakes (235) and Rivers (214) across QC                     | 1995      | Northern pike <i>Esox lucius</i>             | 40-50 cm<br>55-70 cm<br>>70 cm | NR  | NR     | NR  | 0.41<br>0.66<br>1.12  | NR  | Info from I. Guay 1999, MEF, pers. com. | Laliberté 1996       |
| QC: Lakes (235) and Rivers (214) across QC                     | 1995      | Walleye <i>Stizostedion vitreum</i>          | 30-40 cm<br>40-50 cm<br>>50 cm | NR  | NR     | NR  | 0.53<br>0.78<br>1.26  | NR  | Info from I. Guay 1999, MEF, pers. com. | Laliberté 1996       |
| QC: Nottaway-Broadback-Rupert area, South-east coast James Bay | 1990-1991 | Northern pike <i>Esox lucius</i>             | NR                             | NR  | Muscle | 636 | 0.93 ± 0.51   | NR  | --                                      | Langlois et al. 1995 |

Canadian Tissue Residue Guidelines: Methylmercury

| Location   | Year      | Species                             | Life Stage, Age or Size | Sex | Tissue | N   | Total Hg $\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | MeHg $\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | Other information | Reference            |
|--|-----------|-------------------------------------|-------------------------|-----|--------|-----|---|---|-------------------|----------------------|
| QC: Nottaway-Broadback-Rupert area, South-east coast James Bay | 1990-1991 | Walleye <i>Stizostedion vitreum</i> | NR                      | NR  | Muscle | 789 | 0.78 ± 0.46   | NR  | --                | Langlois et al. 1995 |

NR = Not Reported

Var = Variation in the number of samples

Notes:

<sup>a</sup> Unless noted specifically, most mercury in fish is assumed to be MeHg.

<sup>b</sup> Lake had Hg-laden mine tailings discharged to it; this species had low [Hg] compared with Northern pike.

<sup>c</sup> Data unpublished.

<sup>d</sup> Anadromous on Atlantic coast. Introduced on Pacific coast (Eddy and Underhill 1984).

<sup>e</sup> PL = per lake.

<sup>f</sup> Number of fish per lake, except Georgian Bay where 5 fish were obtained.

<sup>g</sup> Hg rating is for lake trout.

**APPENDIX B-3B: LEVELS OF TOTAL MERCURY IN MARINE FISH IN THE CANADIAN ENVIRONMENT.**

| Location                     | Year      | Species   | Life Stage, Age or Size | Sex | Tissue | N      | Total Hg $\mu\text{g}\cdot\text{g}^{-1}$ ww <sup>a</sup> | Other information         | Reference   |
|------------------------------|-----------|---|-------------------------|-----|--------|--------|--|---------------------------|---|
| NB: Bay of Fundy, SW         | 1981      | Atlantic herring<br><i>Clupea harengus harengus</i>   | 1-2 years               | NR  | Muscle | 61     | 0.005 $\pm$ 0.001  | --                        | Braune 1987a  |
| NB: Bay of Fundy, SW         | 1981      | Atlantic herring<br><i>Clupea harengus harengus</i>   | 3-5 years               | NR  | Muscle | 54     | 0.008 to 0.015 range of year class means                 | Hg increased with age     | Braune 1987a  |
| NB: Bay of Fundy, SW         | 1981      | Atlantic herring<br><i>Clupea harengus harengus</i>   | 1-2 years               | NR  | Whole  | 76     | 0.005 to 0.006 range of year class means                 | --                        | Braune 1987a  |
| NB: Bay of Fundy, SW         | 1981      | Atlantic herring<br><i>Clupea harengus harengus</i>   | 3-5 years               | NR  | Whole  | 54     | 0.007 to 0.015 range of year class means                 | [Hg] ~same as muscle only | Braune 1987a  |
| NS: Medway R., at Greenfield | 1972      | American eel<br><i>Anuilla rostrata</i>               | NR                      | NR  | Muscle | 23     | 0.72 THg, 0.40 MeHg                                      | MeHg 50% on average       | Freeman and Horne 1972                              |
| QC: coastal James Bay        | 1987-1994 | Greenland cod<br><i>Gadus ogac</i>                    | 400 mm                  | NR  | NR     | N<br>R | 0.14-0.42  | Standardized length       | Hydro-Québec and Groupe-conseil Génivar Inc. (1997) |
| QC: coastal James Bay        | 1987-1994 | Fourhorn sculpin<br><i>Myoxocephalus quadricornis</i> | 250 mm                  | NR  | NR     | N<br>R | 0.10-0.55  | Standardized length       | Hydro-Québec and Groupe-conseil Génivar Inc. (1997) |

Notes:

<sup>a</sup> Unless noted specifically, most mercury in fish is assumed to be MeHg.

**APPENDIX B-4: LEVELS OF TOTAL MERCURY AND METHYLMERCURY IN AMPHIBIANS AND REPTILES IN THE CANADIAN ENVIRONMENT.**

| Location                                      | Year | Species  | Life Stage       | Tissue                | N | Total Hg<br>$\mu\text{g}\cdot\text{g}^{-1}$ ww | MeHg<br>$\mu\text{g}\cdot\text{g}^{-1}$ ww | %<br>MeHg | Reference         |
|---|------|--|------------------|-----------------------|---|--|--|-----------|-------------------|
| ON: Gray's Cr.,<br>Cornwall area              | 1990 | Snapping Turtle<br>(Reptile)<br><i>Chelydra<br/>serpentina</i> | Eggs             | Whole                 | 5 | 0.110  | NR   | NR        | Bonin et al. 1995 |
| ON: Raquette R.,<br>Cornwall-<br>Massena area | 1990 | Snapping Turtle<br>(Reptile)<br><i>Chelydra<br/>serpentina</i> | Eggs             | Whole                 | 5 | 0.140  | NR   | NR        | Bonin et al. 1995 |
| ON: St. Lawrence<br>R., Dundee area           | 1989 | Snapping Turtle<br>(Reptile)<br><i>Chelydra<br/>serpentina</i> | Eggs             | Whole                 | 7 | 0.090  | NR   | NR        | Bonin et al. 1995 |
| QC: St. Lawrence<br>R., Beauharnois           | 1990 | Snapping Turtle<br>(Reptile)<br><i>Chelydra<br/>serpentina</i> | Eggs             | Whole                 | 3 | 0.150  | NR   | NR        | Bonin et al. 1995 |
| QC: Ottawa R.,<br>Rigaud                      | 1992 | Mudpuppy<br>(Amphibia)<br><i>Necturus<br/>maculosus</i>        | Adult<br>females | Whole<br>no<br>gonads | 6 | $0.087 \pm 0.011$<br>max 0.104                 | NR   | NR        | Bonin et al. 1995 |
| QC: Ottawa R.,<br>Rigaud                      | 1992 | Mudpuppy<br>(Amphibia)<br><i>Necturus<br/>maculosus</i>        | Adult<br>females | Gonads                | 6 | 0.068  | NR   | NR        | Bonin et al. 1995 |
| QC: St. Lawrence<br>R.,<br>Beauharnois        | 1992 | Mudpuppy<br>(Amphibia)<br><i>Necturus<br/>maculosus</i>        | Adult<br>females | Whole<br>no<br>gonads | 6 | $0.239 \pm 0.137$<br>max 0.445                 | NR   | NR        | Bonin et al. 1995 |

Canadian Tissue Residue Guidelines: Methylmercury

| Location                                       | Year | Species  | Life Stage    | Tissue | N  | Total Hg<br>$\mu\text{g}\cdot\text{g}^{-1}$ ww | MeHg<br>$\mu\text{g}\cdot\text{g}^{-1}$ ww | %<br>MeH<br>g | Reference         |
|--|------|--|---------------|--------|----|--|--|---------------|-------------------|
| QC: St. Lawrence R., Beauharnois               | 1992 | Mudpuppy (Amphibia)<br><i>Necturus maculosus</i> | Adult females | Gonads | 6  | 0.193  | NR   | NR            | Bonin et al. 1995 |
| QC: Ottawa R., Vaudreuil                       | 1992 | Mudpuppy (Amphibia)<br><i>Necturus maculosus</i> | Adult         | Whole  | 3  | 0.110  | NR   | NR            | Bonin et al. 1995 |
| QC: Ottawa R./ St. Lawrence R. Montreal East   | 1992 | Mudpuppy (Amphibia)<br><i>Necturus maculosus</i> | Adult         | Whole  | 2  | 0.110  | NR   | NR            | Bonin et al. 1995 |
| QC: Ottawa R. /St. Lawrence R., Illes de Sorel | 1992 | Mudpuppy (Amphibia)<br><i>Necturus maculosus</i> | Adult         | Whole  | 6  | 0.110  | NR   | NR            | Bonin et al. 1995 |
| QC: St. Lawrence R., Pte. Leblanc              | 1992 | Mudpuppy (Amphibia)<br><i>Necturus maculosus</i> | Adult         | Whole  | 4  | 0.130  | NR   | NR            | Bonin et al. 1995 |
| QC: St. Lawrence R., Port Lewis                | 1992 | Mudpuppy (Amphibia)<br><i>Necturus maculosus</i> | Adult         | Whole  | 2  | 0.100  | NR   | NR            | Bonin et al. 1995 |
| QC: St. Lawrence R., Nicolet                   | 1992 | Mudpuppy (Amphibia)<br><i>Necturus maculosus</i> | Adult         | Whole  | 3  | 0.100  | NR   | NR            | Bonin et al. 1995 |
| QC: St. Lawrence R., Ste Anne                  | 1992 | Mudpuppy (Amphibia)<br><i>Necturus maculosus</i> | Adult         | Whole  | 14 | 0.290  | NR   | NR            | Bonin et al. 1995 |

NR = Not Reported



**APPENDIX B-5: LEVELS OF TOTAL MERCURY AND METHYLMERCURY IN FRESHWATER MAMMALS IN THE CANADIAN ENVIRONMENT.**

| Location                                | Year | Species                 | Life Stage   | Tissue | N  | Total Hg<br>$\mu\text{g}\cdot\text{g}^{-1}$ ww | MeHg<br>$\mu\text{g}\cdot\text{g}^{-1}$ ww | % MeHg | Reference   |
|---|------|-------------------------|--------------|--------|----|--|--|--------|-------------|
| MB: Burntwood R. North-central Manitoba | 1979 | Mink                    | <1 – 6 years | Brain  | 21 | 0.80   | NR   | NR     | Kucera 1983 |
|   | -    | <i>Mustela vison</i>    | 88% <2 years | Kidney |    | 1.76   |  |        |             |
|   | 1981 |                         |              | Liver  |    | 2.35   |  |        |             |
| MB: Duck Mountain, near SK border       | 1979 | Mink                    | <1 – 6 years | Brain  | 43 | 0.19   | NR   | NR     | Kucera 1983 |
|   | -    | <i>Mustela vison</i>    | 88% <2 years | Kidney |    | 0.67   |  |        |             |
|   | 1981 |                         |              | Liver  |    | 0.82   |  |        |             |
| MB: Wekusko L. North-central Manitoba   | 1979 | Mink                    | <1 – 6 years | Brain  | 19 | 0.39   | NR   | NR     | Kucera 1983 |
|   | -    | <i>Mustela vison</i>    | 88% <2 years | Kidney |    | 1.04   |  |        |             |
|   | 1981 |                         |              | Liver  |    | 1.46   |  |        |             |
| MB: Wekusko L. North-central Manitoba   | 1979 | Otter                   | <1 – 6 years | Brain  | 17 | 0.28   | NR   | NR     | Kucera 1983 |
|   | -    | <i>Lutra canadensis</i> | 75% <2 years | Kidney |    | 1.07   |  |        |             |
|   | 1981 |                         |              | Liver  |    | 1.75   |  |        |             |
| MB: Whiteshell R., South-east Manitoba  | 1979 | Mink                    | <1 – 6 years | Brain  | 27 | 0.42   | NR   | NR     | Kucera 1983 |
|   | -    | <i>Mustela vison</i>    | 88% <2 years | Kidney |    | 1.47   |  |        |             |
|   | 1981 |                         |              | Liver  |    | 1.71   |  |        |             |
| MB: Whiteshell R., South-east Manitoba  | 1979 | Otter                   | <1 – 6 years | Brain  | 6  | 0.38   | NR   | NR     | Kucera 1983 |
|   | -    | <i>Lutra canadensis</i> | 75% <2 years | Kidney |    | 1.66   |  |        |             |
|   | 1981 |                         |              | Liver  |    | 2.27   |  |        |             |
| MB: Winnipeg R.                         | 1979 | Mink                    | <1 – 6 years | Brain  | 62 | 0.97   | NR   | NR     | Kucera 1983 |
|   | -    | <i>Mustela vison</i>    | 88% <2 years | Kidney |    | 2.98   |  |        |             |
|   | 1981 |                         |              | Liver  |    | 3.93   |  |        |             |

## Canadian Tissue Residue Guidelines: Methylmercury

| Location                                       | Year              | Species                          | Life Stage                         | Tissue                             | N        | Total Hg<br>$\mu\text{g}\cdot\text{g}^{-1}$ ww               | MeHg<br>$\mu\text{g}\cdot\text{g}^{-1}$ ww | % MeHg   | Reference           |
|--|-------------------|----------------------------------|------------------------------------|------------------------------------|----------|--|--|----------|---------------------|
| MB: Winnipeg R.                                | 1979<br>-<br>1981 | Otter<br><i>Lutra canadensis</i> | <1 – 6<br>years<br>75% <2<br>years | Brain<br>Kidney<br>Liver           | 13       | 0.91<br>2.37<br>3.67   | NR   | NR       | Kucera 1983         |
| ON: English R.<br>North-west<br>Ontario        | 1983<br>-<br>1985 | Mink<br><i>Mustela vison</i>     | NR                                 | Brain<br>Kidney<br>Liver<br>Muscle | 11       | 0.45 to 0.64<br>0.48 to 5.54<br>0.56 to 6.9<br>0.16 to 1.62  | NR   | NR       | Wren et al.<br>1986 |
| ON: English R.<br>North-west<br>Ontario        | 1983<br>-<br>1985 | Otter<br><i>Lutra canadensis</i> | NR                                 | Brain<br>Kidney<br>Liver<br>Muscle | 20       | 0.23 to 7.15<br>0.58 to 12.6<br>0.71 to 17.4<br>0.21 to 12.6 | NR   | NR       | Wren et al.<br>1986 |
| ON: English R.<br>Muskoka and<br>Sudbury       | 1983<br>-<br>1985 | Mink<br><i>Mustela vison</i>     | NR                                 | Kidney<br>Liver                    | 35<br>38 | NR   | NR   | 43<br>53 | Wren et al.<br>1986 |
| ON: English R.<br>Muskoka and<br>Sudbury       | 1983<br>-<br>1985 | Otter<br><i>Lutra canadensis</i> | NR                                 | Kidney                             | 32       | NR   | NR   | 19       | Wren et al.<br>1986 |
| ON: English R.<br>Muskoka and<br>Sudbury       | 1983<br>-<br>1985 | Otter<br><i>Lutra canadensis</i> | NR                                 | Muscle                             | 10       | NR   | NR   | 90       | Wren et al.<br>1986 |
| ON: Cambridge                                  | 1983<br>-<br>1985 | Mink<br><i>Mustela vison</i>     | NR                                 | Liver                              | 9        | 0.14 ± 0.08<br>ND to 0.28                                    | NR   | NR       | Wren et al.<br>1986 |
| ON: Herschel<br>Twp. South-<br>central Ontario | 1990<br>s         | Otter<br><i>Lutra canadensis</i> | <1 – 12<br>years                   | Kidney<br>Liver                    | 19       | 1.67 ± 1.09<br>1.54 ± 0.72                                   | NR   | NR       | Evans 1995          |
| ON: Muskoka<br>lakes                           | 1983<br>-<br>1985 | Mink<br><i>Mustela vison</i>     | NR                                 | Brain                              | 4        | 0.55 ± 0.10<br>0.46 to 0.66                                  | NR   | NR       | Wren et al.<br>1986 |

## Canadian Tissue Residue Guidelines: Methylmercury

| Location             | Year              | Species                            | Life Stage | Tissue                                 | N  | Total Hg<br>$\mu\text{g}\cdot\text{g}^{-1}$ ww                   | MeHg<br>$\mu\text{g}\cdot\text{g}^{-1}$ ww | % MeHg | Reference        |
|----------------------|-------------------|------------------------------------|------------|--|----|--|--|--------|------------------|
| ON: Muskoka lakes    | 1983<br>-<br>1985 | Mink<br><i>Mustela vison</i>       | NR         | Kidney                                 | 23 | $2.49 \pm 0.94$<br>0.41 to 4.49                                  | NR   | NR     | Wren et al. 1986 |
| ON: Muskoka lakes    | 1983<br>-<br>1985 | Mink<br><i>Mustela vison</i>       | NR         | Liver                                  | 24 | $2.17 \pm 1.24$<br>0.40 to 6.25                                  | NR   | NR     | Wren et al. 1986 |
| ON: Muskoka lakes    | 1983<br>-<br>1985 | Mink<br><i>Mustela vison</i>       | NR         | Muscle                                 | 11 | $1.62 \pm 1.07$<br>0.01 to 4.08                                  | NR   | NR     | Wren et al. 1986 |
| ON: Muskoka lakes    | 1983<br>-<br>1985 | Otter<br><i>Lutra canadensis</i>   | NR         | Brain                                  | 2  | $0.52 \pm 0.18$<br>0.40 to 0.63                                  | NR   | NR     | Wren et al. 1986 |
| ON: Muskoka lakes    | 1983<br>-<br>1985 | Otter<br><i>Lutra canadensis</i>   | NR         | Kidney                                 | 11 | $1.28 \pm 0.72$<br>0.34 to 2.64                                  | NR   | NR     | Wren et al. 1986 |
| ON: Muskoka lakes    | 1983<br>-<br>1985 | Otter<br><i>Lutra canadensis</i>   | NR         | Liver                                  | 12 | $1.50 \pm 0.75$<br>0.20 to 2.46                                  | NR   | NR     | Wren et al. 1986 |
| ON: Muskoka lakes    | 1983<br>-<br>1985 | Otter<br><i>Lutra canadensis</i>   | NR         | Muscle                                 | 6  | $0.55 \pm 0.32$<br>0.25 to 1.21                                  | NR   | NR     | Wren et al. 1986 |
| ON: Near Parry Sound | NR                | Beaver<br><i>Castor canadensis</i> | NR         | Kidney<br>Liver<br>Muscle              | NR | 0.03<br>0.03<br>0.03   | NR   | NR     | Wren 1984        |
| ON: Near Parry Sound | NR                | Otter<br><i>Lutra canadensis</i>   | NR         | Intestine<br>Kidney<br>Liver<br>Muscle | NR | $0.4 \pm 0.2$<br>$1.4 \pm 0.3$<br>$3.0 \pm 2.2$<br>$0.9 \pm 0.2$ | NR   | NR     | Wren 1984        |

## Canadian Tissue Residue Guidelines: Methylmercury

| Location                                   | Year              | Species                          | Life Stage | Tissue                    | N  | Total Hg<br>$\mu\text{g}\cdot\text{g}^{-1}$ ww | MeHg<br>$\mu\text{g}\cdot\text{g}^{-1}$ ww | % MeHg | Reference           |
|--|-------------------|----------------------------------|------------|---------------------------|----|--|--|--------|---------------------|
| ON: Near Parry Sound                       | NR                | Racoon<br><i>Procyon lotor</i>   | NR         | Kidney<br>Liver<br>Muscle | NR | 1.1 ± 0.4<br>4.5 ± 3.5<br>0.3 ± 0.1            | NR   | NR     | Wren 1984           |
| ON: Sudbury                                | 1983<br>-<br>1985 | Mink<br><i>Mustela vison</i>     | NR         | Brain                     | 2  | 0.36 ± 0.11<br>0.28 to 0.44                    | NR   | NR     | Wren et al.<br>1986 |
| ON: Sudbury                                | 1983<br>-<br>1985 | Mink<br><i>Mustela vison</i>     | NR         | Liver                     | 31 | 0.56 ± 0.40<br>0.07 to 1.75                    | NR   | NR     | Wren et al.<br>1986 |
| ON: Sudbury                                | 1983<br>-<br>1985 | Mink<br><i>Mustela vison</i>     | NR         | Muscle                    | 19 | 0.38 ± 0.26<br>ND to 0.88                      | NR   | NR     | Wren et al.<br>1986 |
| ON: Sudbury                                | 1983<br>-<br>1985 | Mink<br><i>Mustela vison</i>     | NR         | Kidney                    | 21 | 0.59 ± 0.23<br>0.13 to 1.00                    | NR   | NR     | Wren et al.<br>1986 |
| ON: Sudbury                                | 1983<br>-<br>1985 | Otter<br><i>Lutra canadensis</i> | NR         | Brain                     | 3  | 0.24<br>0.23 to 0.25                           | NR   | NR     | Wren et al.<br>1986 |
| ON: Sudbury                                | 1983<br>-<br>1985 | Otter<br><i>Lutra canadensis</i> | NR         | Kidney                    | 23 | 0.59 ± 0.26<br>0.05 to 1.31                    | NR   | NR     | Wren et al.<br>1986 |
| ON: Sudbury                                | 1983<br>-<br>1985 | Otter<br><i>Lutra canadensis</i> | NR         | Liver                     | 36 | 0.86 ± 0.92<br>ND to 4.14                      | NR   | NR     | Wren et al.<br>1986 |
| ON: Sudbury                                | 1983<br>-<br>1985 | Otter<br><i>Lutra canadensis</i> | NR         | Muscle                    | 17 | 0.30 ± 0.17<br>0.07 to 0.17                    | NR   | NR     | Wren et al.<br>1986 |
| ON: Turkey L.,<br>Sault Ste. Marie<br>area | 1983<br>-<br>1985 | Mink<br><i>Mustela vison</i>     | NR         | Liver                     | 16 | 2.36 ± 1.63<br>1.12 to 7.50                    | NR   | NR     | Wren et al.<br>1986 |

## Canadian Tissue Residue Guidelines: Methylmercury

| Location                                     | Year              | Species                          | Life Stage | Tissue | N  | Total Hg<br>$\mu\text{g}\cdot\text{g}^{-1}$ ww | MeHg<br>$\mu\text{g}\cdot\text{g}^{-1}$ ww | % MeHg | Reference                      |
|--|-------------------|----------------------------------|------------|--------|----|--|--|--------|--------------------------------|
| ON: Turkey L.,<br>Sault Ste. Marie<br>area   | 1983<br>-<br>1985 | Mink<br><i>Mustela vison</i>     | NR         | Kidney | 14 | $1.86 \pm 0.73$<br>0.80 to 2.82                | NR   | NR     | Wren et al.<br>1986            |
| ON: Turkey L.,<br>Sault Ste. Marie<br>area   | 1983<br>-<br>1985 | Mink<br><i>Mustela vison</i>     | NR         | Muscle | 14 | $1.04 \pm 0.46$<br>0.44 to 1.91                | NR   | NR     | Wren et al.<br>1986            |
| ON: Turkey L.,<br>Sault Ste. Marie<br>area   | 1983<br>-<br>1985 | Mink<br><i>Mustela vison</i>     | NR         | Brain  | 16 | $2.36 \pm 1.63$<br>1.12 to 7.50                | NR   | NR     | Wren et al.<br>1986            |
| ON: Turkey L.,<br>Sault Ste. Marie<br>area   | 1983<br>-<br>1985 | Otter<br><i>Lutra canadensis</i> | NR         | Liver  | 8  | $1.99 \pm 0.52$<br>1.18 to 2.60                | NR   | NR     | Wren et al.<br>1986            |
| ON: Turkey L.,<br>Sault Ste. Marie<br>area   | 1983<br>-<br>1985 | Otter<br><i>Lutra canadensis</i> | NR         | Kidney | 6  | $1.90 \pm 0.76$<br>1.42 to 2.52                | NR   | NR     | Wren et al.<br>1986            |
| ON: Turkey L.,<br>Sault Ste. Marie<br>area   | 1983<br>-<br>1985 | Otter<br><i>Lutra canadensis</i> | NR         | Muscle | 7  | $1.02 \pm 0.34$<br>0.36 to 1.39                | NR   | NR     | Wren et al.<br>1986            |
| QC: natural areas<br>of N. Québec            | 1990<br>s         | Mink<br><i>Mustela vison</i>     | NR         | Liver  | 38 | 4.85<br>1.17 to 15.60                          | NR   | NR     | Bélanger and<br>Larivière 1997 |
| QC: natural areas<br>of N. Québec            | 1990<br>s         | Mink<br><i>Mustela vison</i>     | NR         | Kidney | 38 | 2.45<br>0.73 to 6.32                           | NR   | NR     | Bélanger and<br>Larivière 1997 |
| QC: natural areas<br>of N. Québec            | 1990<br>s         | Mink<br><i>Mustela vison</i>     | NR         | Brain  | 37 | 0.93<br>0.27 to 2.57                           | NR   | NR     | Bélanger and<br>Larivière 1997 |
| QC: Great Whale<br>area, Northern.<br>Québec | 1989<br>-<br>1990 | Mink<br><i>Mustela vison</i>     | NR         | Muscle | 6  | $2.40 \pm 2.24$                                | NR   | NR     | Langlois et al.<br>1995        |

NR= Not Reported

ND = Not Detected

**APPENDIX B-6: LEVELS OF TOTAL MERCURY AND METHYLMERCURY IN MARINE MAMMALS IN THE CANADIAN ENVIRONMENT.**

| Location           | Year          | Species                                      | Life Stage, Age or Sex | Tissue | N  | Total Hg<br>$\mu\text{g}\cdot\text{g}^{-1}$ ww | MeHg<br>$\mu\text{g}\cdot\text{g}^{-1}$ ww | % MeHg             | Reference                              |
|--------------------|---------------|--|------------------------|--------|----|--|--|--------------------|--|
| Arctic:<br>Eastern | 1990          | Beluga whale<br><i>Delphinapterus leucas</i> | NR                     | Liver  | 66 | 6  | NR   | NR                 | Wagemann et al. 1995                   |
| Arctic:<br>Eastern | 1993-<br>1994 | Beluga whale<br><i>Delphinapterus leucas</i> | 2.5 to 24.5 years      | Liver  | 73 | 10.2 ± 8.0<br>1.24 to 38.6                     | NR   | NR                 | Wagemann 1995;<br>Wagemann et al. 1996 |
| Arctic:<br>Eastern | 1993-<br>1994 | Beluga whale<br><i>Delphinapterus leucas</i> | 2.5 to 24.5 years      | Muktuk | 45 | 0.59 ± 0.22<br>0.32 to 1.37                    | NR   | NR                 | Wagemann 1995;<br>Wagemann et al. 1996 |
| Arctic:<br>Eastern | 1993-<br>1994 | Beluga whale<br><i>Delphinapterus leucas</i> | 2.5 to 24.5 yrs        | Muscle | 74 | 1.04 ± 0.43<br>0.44 to 2.7                     | NR   | NR                 | Wagemann 1995;<br>Wagemann et al. 1996 |
| Arctic:<br>Eastern | 1983          | Narwhal<br><i>Monodon monoceros</i>          | NR                     | Liver  | 38 | 5.9  | NR   | NR                 | Wagemann et al. 1995                   |
| Arctic:<br>Eastern | 1992-<br>1994 | Narwhal<br><i>Monodon monoceros</i>          | 280-470 cm             | Kidney | 55 | 1.93 ± 1.12<br>0.29 to 6.92                    | NR   | NR                 | Wagemann et al. 1996                   |
| Arctic:<br>Eastern | 1992-<br>1994 | Narwhal<br><i>Monodon monoceros</i>          | 280-470 cm             | Liver  | 55 | 10.8 ± 8.05<br>0.32 to 37.2                    | NR   | NR                 | Wagemann et al. 1996                   |
| Arctic:<br>Eastern | 1992-<br>1994 | Narwhal<br><i>Monodon monoceros</i>          | 280-470 cm             | Muktuk | 48 | 0.59 ± 0.18<br>0.16 to 1.27                    | NR   | NR                 | Wagemann et al. 1996                   |
| Arctic:<br>Eastern | 1992-<br>1994 | Narwhal<br><i>Monodon monoceros</i>          | 280-470 cm             | Muscle | 56 | 1.03 ± 0.37<br>0.41 to 1.94                    | NR   | NR                 | Wagemann et al. 1996                   |
| Arctic:<br>Eastern | 1989          | Ringed seal<br><i>Phoca hispida</i>          | NR                     | Liver  | 54 | 8.8  | NR   | 15-30 <sup>a</sup> | Wagemann et al. 1995                   |

## Canadian Tissue Residue Guidelines: Methylmercury

| Location            | Year      | Species  | Life Stage, Age or Sex        | Tissue | N   | Total Hg $\mu\text{g}\cdot\text{g}^{-1}$ ww | MeHg $\mu\text{g}\cdot\text{g}^{-1}$ ww | % MeHg | Reference                              |
|---------------------|-----------|--|-------------------------------|--------|-----|---|---|--------|--|
| Arctic: Eastern     | 1989-1993 | Ringed seal<br><i>Phoca hispida</i>            | Young of the year to 22 years | Liver  | 115 | 8.34 $\pm$ 7.03<br>0.36 to 38.7             | NR                                      | NR     | Wagemann 1995;<br>Wagemann et al. 1996 |
| Arctic: Eastern     | 1989-1993 | Ringed seal<br><i>Phoca hispida</i>            | Young of the year to 22 years | Muscle | 61  | 0.39 $\pm$ 0.17                             | NR                                      | NR     | Wagemann 1995;<br>Wagemann et al. 1996 |
| Arctic: Hudson Bay  | 1994      | Atlantic walrus<br><i>Odobenus r. rosmarus</i> | NR                            | Liver  | 143 | 1.2   | NR                                      | NR     | Wagemann et al. 1995                   |
| Arctic: location NR | NR        | Narwhal<br><i>Monodon monoceros</i>            | NR                            | Muscle | 2   | 1.4, 1.6 (approx) <sup>b</sup>              | 1.4, 1.6 (approx) <sup>b</sup>          | 96     | Wagemann et al. 1997                   |
| Arctic: location NR | NR        | Ringed seal<br><i>Phoca hispida</i>            | NR                            | Muscle | 12  | 0.6 $\pm$ 0.4 (approx) <sup>b</sup>         | 0.6 $\pm$ 0.4 (approx) <sup>b</sup>     | 100    | Wagemann et al. 1997                   |
| Arctic: location NR | NR        | Beluga whale<br><i>Delphinapterus leucas</i>   | NR                            | Muscle | 2   | 2.3 (approx) <sup>b</sup>                   | 2.3 (approx) <sup>b</sup>               | 98     | Wagemann et al. 1997                   |
| Arctic: Pond inlet  | 1978-1979 | Narwhal<br><i>Monodon monoceros</i>            | 262-485 cm                    | Kidney | 55  | 1.71 $\pm$ 1.01<br>0.31 to 4.67             | NR                                      | NR     | Wagemann et al. 1996                   |
| Arctic: Pond inlet  | 1978-1979 | Narwhal<br><i>Monodon monoceros</i>            | 262-485 cm                    | Liver  | 38  | 6.10 $\pm$ 3.10<br>0.57 to 13.1             | NR                                      | NR     | Wagemann et al. 1996                   |
| Arctic: Pond inlet  | 1978-1979 | Narwhal<br><i>Monodon monoceros</i>            | 262-485 cm                    | Muscle | 58  | 0.85 $\pm$ 0.28<br>0.22 to 1.59             | NR                                      | NR     | Wagemann et al. 1996                   |
| Arctic: Western     | 1993-1994 | Beluga whale<br><i>Delphinapterus leucas</i>   | 7 to 35 years                 | Liver  | 77  | 27.1 $\pm$ 24.7<br>0.31 to 116              | NR                                      | NR     | Wagemann 1995;<br>Wagemann et al. 1996 |

## Canadian Tissue Residue Guidelines: Methylmercury

| Location                              | Year      | Species   | Life Stage, Age or Sex        | Tissue                   | N   | Total Hg<br>$\mu\text{g}\cdot\text{g}^{-1}$ ww          | MeHg<br>$\mu\text{g}\cdot\text{g}^{-1}$ ww | % MeHg           | Reference                              |
|---------------------------------------|-----------|---|-------------------------------|--------------------------|-----|---|--|------------------|--|
| Arctic: Western                       | 1993-1994 | Beluga whale<br><i>Delphinapterus leucas</i>              | 7 to 35 years                 | Muktuk                   | 65  | $0.78 \pm 0.41$<br>0.19 to 1.93                         | NR   | NR               | Wagemann 1995;<br>Wagemann et al. 1996 |
| Arctic: Western                       | 1993-1994 | Beluga whale<br><i>Delphinapterus leucas</i>              | 7 to 35 years                 | Muscle                   | 76  | $1.34 \pm 0.67$<br>0.41 to 3.44                         | NR   | NR               | Wagemann 1995;<br>Wagemann et al. 1996 |
| Arctic: Western                       | 1987-1993 | Ringed seal<br><i>Phoca hispida</i>                       | Young of the year to 38 years | Liver <sup>c</sup>       | 145 | $32.6 \pm 35.2$<br>0.23 to 219                          | NR   | NR               | Wagemann 1995;<br>Wagemann et al. 1996 |
| Arctic: Western                       | 1987-1993 | Ringed seal<br><i>Phoca hispida</i>                       | Young of the year to 38 years | Muscle                   | 133 | $0.41 \pm 0.29$   | NR   | NR               | Wagemann 1995;<br>Wagemann et al. 1996 |
| Atlantic coast                        | 1988      | Harp seal<br><i>Phoca groenlandica</i>                    | Adult females & pups          | Liver                    | 20  | 10  | NR   | NR               | Wagemann et al. 1995                   |
| Atlantic coast                        | 1988      | Pilot whale<br><i>Globicephala malaena</i>                | NR                            | Liver                    | 39  | 23  | NR   | NR               | Wagemann et al. 1995                   |
| Atlantic coast                        | 1988      | White-beaked dolphin<br><i>Lagenorhynchus albirostris</i> | NR                            | Liver                    | 27  | 0.6   | NR   | NR               | Wagemann et al. 1995                   |
| QC: Great Whale area, Northern Québec | 1989-1990 | Beluga whale<br><i>Delphinapterus leucas</i>              | NR                            | Brain<br>Liver<br>Muscle | 6   | $2.63 \pm 2.87$<br>$20.34 \pm 19.60$<br>$2.60 \pm 2.06$ | NR<br>$7.26 \pm 5.21$<br>$1.01 \pm 0.58$   | NR<br>36%<br>39% | Langlois et al. 1995                   |
| QC: Great Whale area, Northern Québec | 1989-1990 | Freshwater seal<br><i>Phoca vitulina</i> <sup>d</sup>     | NR                            | Liver<br>Muscle          | 2   | $28.8 \pm 37.06$<br>$1.10 \pm 0.37$                     | $4.55 \pm 4.6$<br>$1.04 \pm 0.37$          | 16%<br>95%       | Langlois et al. 1995                   |



Canadian Tissue Residue Guidelines: Methylmercury

| Location                              | Year      | Species                                      | Life Stage, Age or Sex | Tissue                             | N  | Total Hg $\mu\text{g}\cdot\text{g}^{-1}$ ww              | MeHg $\mu\text{g}\cdot\text{g}^{-1}$ ww | % MeHg                 | Reference            |
|---------------------------------------|-----------|--|------------------------|------------------------------------|----|--|---|------------------------|----------------------|
| QC: Great Whale area, Northern Québec | 1989-1990 | Ringed seal<br><i>Phoca hispida</i>          | NR                     | Brain<br>Kidney<br>Liver<br>Muscle | 8  | 0.19 ± 0.09<br>0.49 ± 0.07<br>5.12 ± 4.64<br>0.32 ± 0.19 | NR<br>NR<br>1.01 ± 0.87<br>0.11 ± 0.08  | NR<br>NR<br>20%<br>34% | Langlois et al. 1995 |
| QC: St. Lawrence R.                   | 1982-1987 | Beluga whale<br><i>Delphinapterus leucas</i> | YOY to 30 years        | Muscle                             | 9  | 2.46 ± 1.46<br>0.89 to 5.54                              | NR                                      | NR                     | Wagemann et al. 1996 |
| QC: St. Lawrence R.                   | 1982-1987 | Beluga whale<br><i>Delphinapterus leucas</i> | YOY to 30 years        | Liver                              | 30 | 33.6 ± 43.0<br>0.38 to 202                               | NR                                      | NR                     | Wagemann et al. 1996 |

NR = Not Reported

Notes:

<sup>a</sup> Or lower than 10% if calculated using regression rather than an average of percentages (Wagemann et al. 1997; R. Wagemann 1999, DFO, pers. com.).

<sup>b</sup> Numbers estimated from graphical presentation.

<sup>c</sup> Ringed seal liver, but not beluga liver, is consumed by Indigenous People. Most Hg in livers was inorganic Hg.

<sup>d</sup> The freshwater seal is a population of harbour seals captive of inland waters in Des Loups-Marins Lake and vicinity, 74° 00W, 56° 30N approx (L-G de Repentigny 1999, Environment Canada, pers.com.)

## APPENDIX B-7: LEVELS OF TOTAL MERCURY AND METHYLMERCURY IN BIRDS IN THE CANADIAN ENVIRONMENT.

| Location                                | Year      | Species                                       | Life Stage | Tissue                    | N            | Total Hg<br>$\mu\text{g}\cdot\text{g}^{-1}$ ww <sup>a</sup> | MeHg<br>$\mu\text{g}\cdot\text{g}^{-1}$<br>ww | %<br>MeHg      | Reference              |
|---|-----------|---|------------|---------------------------|--------------|---|---|----------------|------------------------|
| BC: Lower mainland, Lower Fraser Valley | 1990-1992 | Bald eagle<br><i>Haliaeetus leucocephalus</i> | Eggs       | Albumen and Yolk          | 6<br>7<br>11 | 0.186 to 0.358<br>0.174 to 0.296<br>NR                      | NR  | NR<br>NR<br>88 | Elliott et al. 1996    |
| BC: Vancouver Island, Crofton           | 1990-1992 | Bald eagle<br><i>Haliaeetus leucocephalus</i> | Eggs       | Albumen and Yolk          | 3            | 0.096 to 0.384  | NR  | NR             | Elliott et al. 1996    |
| BC: Vancouver Island, Johnstone Strait  | 1990-1992 | Bald eagle<br><i>Haliaeetus leucocephalus</i> | Eggs       | Albumen and Yolk          | 7            | 0.236 to 0.367  | NR  | NR             | Elliott et al. 1996    |
| BC: Vancouver Island, Nanaimo           | 1990-1992 | Bald eagle<br><i>Haliaeetus leucocephalus</i> | Eggs       | Albumen and Yolk          | 8            | 0.110 to 0.198  | NR  | NR             | Elliott et al. 1996    |
| NB: Quoddy region                       | 1984      | Bonaparte's gull<br><i>Larus philadephia</i>  | Adult      | Feathers                  | 8<br>4<br>4  | 4.14 ± 2.22<br>all<br>4.82 ± 2.99<br>F<br>3.45 ± 1.15<br>M  | NR  | NR             | Braune and Gaskin 1987 |
| NB: Quoddy region, Bay of Fundy         | 1978-1984 | Arctic tern<br><i>Sterna paradisaea</i>       | Adult      | Brain                     | 31           | 0.094   | NR  | NR             | Braune 1987b           |
| NB: Quoddy region, Bay of Fundy         | 1978-1984 | Arctic tern<br><i>Sterna paradisaea</i>       | Adult      | Kidney<br>Liver<br>Muscle | 36           | 0.453<br>0.470<br>0.089                                     | NR  | NR             | Braune 1987b           |

Canadian Tissue Residue Guidelines: Methylmercury

| Location                        | Year      | Species   | Life Stage | Tissue | N   | Total Hg<br>$\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | MeHg<br>$\mu\text{g}\cdot\text{g}^{-1}\text{ ww}$ | %<br>MeHg | Reference    |       |
|---------------------------------|-----------|---|------------|--------|-----|--|---|-----------|--------------|-------|
| NB: Quoddy region, Bay of Fundy | 1978-1984 | Black guillemots<br><i>Cephus grylle</i>          | Adult      | Brain  | 4   | 0.123  | NR  | NR        | Braune 1987b |       |
|                                 |           |   |            | Kidney |     | 0.491  |   |           |              |       |
|                                 |           |   |            | Liver  |     | 0.513  |   |           |              |       |
|                                 |           |   |            | Muscle |     | 0.113  |   |           |              |       |
| NB: Quoddy region, Bay of Fundy | 1978-1984 | Black-legged kittiwake<br><i>Rissa tridactyla</i> | Adult      | Brain  | 14  | 0.038  | NR  | NR        | Braune 1987b |       |
|                                 |           |   |            | Kidney |     | 18   |   |           |              | 0.242 |
|                                 |           |   |            | Liver  |     | 19   |   |           |              | 0.372 |
|                                 |           |   |            | Muscle |     | 20   |   |           |              | 0.037 |
| NB: Quoddy region, Bay of Fundy | 1978-1984 | Bonaparte's gull<br><i>Larus philadelphia</i>     | Adult      | Brain  | 137 | 0.101  | NR  | NR        | Braune 1987b |       |
| NB: Quoddy region, Bay of Fundy | 1978-1984 | Bonaparte's gull<br><i>Larus philadelphia</i>     | Adult      | Kidney | 145 | 0.418  | NR  | NR        | Braune 1987b |       |
|                                 |           |   |            | Liver  |     | 0.450  |   |           |              |       |
|                                 |           |   |            | Muscle |     | 0.075  |   |           |              |       |
| NB: Quoddy region, Bay of Fundy | 1978-1984 | Common eider duck<br><i>Somateria mollissima</i>  | Adult      | Kidney | 11  | 0.358  | NR  | NR        | Braune 1987b |       |
|                                 |           |   |            | Liver  |     | 0.987  |   |           |              |       |
|                                 |           |   |            | Muscle |     | 0.153  |   |           |              |       |
| NB: Quoddy region, Bay of Fundy | 1978-1984 | Common tern<br><i>Sterna hirundo</i>              | Adult      | Brain  | 29  | 0.190  | NR  | NR        | Braune 1987b |       |
| NB: Quoddy region, Bay of Fundy | 1978-1984 | Common tern<br><i>Sterna hirundo</i>              | Adult      | Kidney | 30  | 1.505  | NR  | NR        | Braune 1987b |       |
|                                 |           |   |            | Liver  |     | 1.249  |   |           |              |       |
|                                 |           |   |            | Muscle |     | 0.166  |   |           |              |       |

Canadian Tissue Residue Guidelines: Methylmercury

| Location                                      | Year      | Species  | Life Stage | Tissue                             | N        | Total Hg<br>$\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | MeHg<br>$\mu\text{g}\cdot\text{g}^{-1}\text{ ww}$ | %<br>MeHg | Reference          |
|---|-----------|--|------------|------------------------------------|----------|--|---|-----------|--------------------|
| NB: Quoddy region, Bay of Fundy               | 1978-1984 | Double-crested cormorant<br><i>Phalacrocorax auritus</i> | Adult      | Brain                              | 1        | 0.360  | NR  | NR        | Braune 1987b       |
| NB: Quoddy region, Bay of Fundy               | 1978-1984 | Double-crested cormorant<br><i>Phalacrocorax auritus</i> | Adult      | Kidney<br>Liver<br>Muscle          | 3        | 5.35<br>7.05<br>0.606  | NR  | NR        | Braune 1987b       |
| NB: Quoddy region, Bay of Fundy               | 1978-1984 | Herring gull<br><i>Larus argentatus</i>                  | Adult      | Brain<br>Kidney<br>Liver<br>Muscle | 4        | 0.056<br>0.350<br>0.482<br>0.101                                 | NR  | NR        | Braune 1987b       |
| NB: Quoddy region, Bay of Fundy               | 1978-1984 | Red-necked phalarope<br><i>Phalaropus lobatus</i>        | Adult      | Liver<br>Muscle                    | 12<br>13 | 0.225<br>0.046   | NR  | NR        | Braune 1987b       |
| NB: Quoddy region, Bay of Fundy               | 1978-1984 | Six bird species   | Adult      | Feathers                           | 968      | 0 to ~12<br>$\mu\text{g}\cdot\text{g}^{-1}\text{ dw}$            | NR  | NR        | Braune 1987b       |
| ON: Akwesasne Reserve, North-east of Cornwall | 1991      | Tree swallow<br><i>Tachycineta bicolor</i>               | Eggs       | Albumen and yolk                   | 5        | 0.043  | NR  | NR        | Bishop et al. 1995 |
| ON: Algonquin/Parry Sound                     | 1970-1978 | Common Loon<br><i>Gavia immer</i>                        | Eggs       | Albumen and Yolk                   | 61       | 0.81 to 1.11 avg   | 0.79  | 71        | Frank et al. 1983  |

Canadian Tissue Residue Guidelines: Methylmercury

| Location                                | Year          | Species   | Life Stage                         | Tissue                   | N   | Total Hg<br>$\mu\text{g}\cdot\text{g}^{-1}$ ww <sup>a</sup>      | MeHg<br>$\mu\text{g}\cdot\text{g}^{-1}$<br>ww | %<br>MeHg | Reference                  |
|---|---------------|---|------------------------------------|--------------------------|-----|--|---|-----------|----------------------------|
| ON: Algonquin/<br>Parry Sound           | 1971-<br>1974 | Common Loon<br><i>Gavia immer</i>               | Healthy<br>Adults &<br>Juveniles   | Brain<br>Liver<br>Muscle | 128 | 0.44 to<br>0.57 1.92<br>to 6.35<br>0.79 to<br>1.14<br>(means)    | NR  | NR        | Frank et al. 1983          |
| ON: Algonquin/<br>Parry Sound           | 1971-<br>1974 | Common Loon<br><i>Gavia immer</i>               | Emaciated<br>Adults &<br>Juveniles | Brain<br>Liver<br>Muscle | 29  | 1.03 to<br>1.82<br>16.3 to<br>26.4<br>2.71 to<br>5.41<br>(means) | NR  | NR        | Frank et al. 1983          |
| ON: Central<br>Ontario                  | 1992          | Common Loon<br><i>Gavia immer</i>               | Chick                              | Feather                  | 17  | 2.31 ± 0.65  | NR  | NR        | Scheuhammer et<br>al. 1998 |
| ON: Central<br>Ontario                  | 1992          | Common Loon<br><i>Gavia immer</i>               | Chick                              | Blood                    | 35  | 0.14 ± 0.12  | NR  | NR        | Scheuhammer et<br>al. 1998 |
| ON: Central<br>Ontario                  | 1992          | Common Loon<br><i>Gavia immer</i>               | Adult                              | Blood                    | 22  | 2.06 ± 0.94  | NR  | NR        | Scheuhammer et<br>al. 1998 |
| ON: Central<br>Ontario                  | 1992          | Common Loon<br><i>Gavia immer</i>               | Adult                              | Feather                  | 20  | 13.3 ± 4.19  | NR  | NR        | Scheuhammer et<br>al. 1998 |
| ON: Clay L.<br>North-western<br>Ontario | 1971          | American<br>wigeon<br><i>Anas<br/>americana</i> | 3 Immature<br>2 Adult              | Breast                   | 5   | 0.48 ± 0.30<br>0.3 to 0.9  | NR  | NR        | Vermeer et al.<br>1973     |
| ON: Clay L.<br>North-western<br>Ontario | 1971          | Blue-winged<br>teal<br><i>Anas discors</i>      | Immature                           | Breast                   | 1   | 10.4   | 7.4   | 71        | Vermeer et al.<br>1973     |

Canadian Tissue Residue Guidelines: Methylmercury

| Location                                | Year          | Species   | Life Stage | Tissue              | N   | Total Hg<br>$\mu\text{g}\cdot\text{g}^{-1}$ ww <sup>a</sup> | MeHg<br>$\mu\text{g}\cdot\text{g}^{-1}$<br>ww | %<br>MeHg | Reference              |
|---|---------------|---|------------|---------------------|-----|---|---|-----------|------------------------|
| ON: Clay L.<br>North-western<br>Ontario | 1971          | Common<br>goldeneye<br><i>Bucephala<br/>clangula</i>    | Immature   | Breast              | 1   | 16.8  | 12.3  | 73        | Vermeer et al.<br>1973 |
| ON: Clay L.<br>North-western<br>Ontario | 1971          | Common<br>merganser<br><i>Mergus<br/>merganser</i>      | Immature   | Breast              | 1   | 19.4  | 14.9  | 77        | Vermeer et al.<br>1973 |
| ON: Clay L.<br>North-western<br>Ontario | 1971          | Herring gull<br><i>Larus<br/>argentatus</i>             | Egg        | Albumen<br>and yolk | 14  | 8.4 ± 3.8<br>S.D  | NR  | NR        | Vermeer et al.<br>1973 |
| ON: Clay L.<br>North-western<br>Ontario | 1971          | Herring gull<br><i>Larus<br/>argentatus</i>             | Egg        | Albumen             | 4   | 12.0 ± 7.8<br>S.D.  | NR  | NR        | Vermeer et al.<br>1973 |
| ON: Clay L.<br>North-western<br>Ontario | 1971          | Herring gull<br><i>Larus<br/>argentatus</i>             | Egg        | Yolk                | 4   | 2.3 ± 1.1<br>S.D.   | NR  | NR        | Vermeer et al.<br>1973 |
| ON: Clay L.<br>North-western<br>Ontario | 1971          | Hooded<br>merganser<br><i>Lophodytes<br/>cucullatus</i> | Immature   | Breast              | 1   | 17.9  | 17.8  | 99        | Vermeer et al.<br>1973 |
| ON: Clay L.<br>North-western<br>Ontario | 1971          | Mallard<br><i>Anas<br/>platyrhynchos</i>                | Immature   | Breast              | 1   | 0.16  | 0.11  | 69        | Vermeer et al.<br>1973 |
| ON: Detroit R.                          | 1973-<br>1992 | Herring gull<br><i>Larus<br/>argentatus</i>             | Eggs       | Albumen<br>and Yolk | Var | 0.14 to<br>0.24   | NR  | NR        | Koster et al. 1996     |
| ON: Georgian<br>Bay<br>Wye Marsh        | 1991          | Tree swallow<br><i>Tachycineta<br/>bicolor</i>          | Eggs       | Albumen<br>and yolk | 10  | 0.066   | NR  | NR        | Bishop et al. 1995     |

Canadian Tissue Residue Guidelines: Methylmercury

| Location                         | Year          | Species   | Life Stage | Tissue              | N               | Total Hg<br>$\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | MeHg<br>$\mu\text{g}\cdot\text{g}^{-1}\text{ ww}$ | %<br>MeHg | Reference               |
|----------------------------------|---------------|---|------------|---------------------|-----------------|--|---|-----------|-------------------------|
| ON: L. Erie<br>2 locations       | 1973-<br>1992 | Herring gull<br><i>Larus<br/>argentatus</i>                       | Eggs       | Albumen<br>and Yolk | Var             | 0.13 to<br>0.54  | NR  | NR        | Koster et al. 1996      |
| ON: L. Erie<br>Mud Creek         | 1991          | Tree swallow<br><i>Tachycineta<br/>bicolor</i>                    | Eggs       | Albumen<br>and yolk | 10              | 0.079  | NR  | NR        | Bishop et al. 1995      |
| ON: L. Huron                     | 1972-<br>1973 | Double-crested<br>cormorant<br><i>Phalacrocorax<br/>auritus</i>   | Eggs       | Albumen<br>and Yolk | UK <sup>b</sup> | 0.32 to<br>0.83  | NR  | NR        | Weseloh et al.<br>1983  |
| ON: L. Huron<br>2 locations      | 1973-<br>1992 | Herring gull<br><i>Larus<br/>argentatus</i>                       | Eggs       | Albumen<br>and Yolk | Var             | 0.13 to<br>0.39  | NR  | NR        | Koster et al. 1996      |
| ON: L. Ontario                   | 1971          | Common tern<br><i>Sterna hirundo</i>                              | Eggs       | Albumen<br>and Yolk | UK <sup>b</sup> | 1.08   | NR  | NR        | Gilbertson 1974         |
| ON: L. Ontario<br>6 locations    | 1973-<br>1992 | Herring gull<br><i>Larus<br/>argentatus</i>                       | Eggs       | Albumen<br>and Yolk | Var             | 0.18 to<br>0.88  | NR  | NR        | Koster et al. 1996      |
| ON: L. Ontario<br>Long Point Tip | 1991          | Tree swallow<br><i>Tachycineta<br/>bicolor</i>                    | Eggs       | Albumen<br>and yolk | 10              | 0.076  | NR  | NR        | Bishop et al. 1995      |
| ON: L. St. Clair                 | 1973          | Black-crowned<br>night-heron,<br><i>Nycticorax<br/>nycticorax</i> | Eggs       | Albumen<br>and Yolk | UK <sup>b</sup> | 0.31 to<br>0.73  | NR  | NR        | Stendell et al.<br>1976 |
| ON: L. St. Clair                 | 1973          | Common tern<br><i>Sterna hirundo</i>                              | Eggs       | Albumen<br>and Yolk | UK <sup>b</sup> | 0.73 to<br>1.09  | NR  | NR        | Stendell et al.<br>1976 |

## Canadian Tissue Residue Guidelines: Methylmercury

| Location   | Year          | Species   | Life Stage | Tissue                   | N               | Total Hg<br>$\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | MeHg<br>$\mu\text{g}\cdot\text{g}^{-1}\text{ ww}$ | %<br>MeHg | Reference               |
|--|---------------|---|------------|--------------------------|-----------------|--|---|-----------|-------------------------|
| ON: L. St. Clair   | 1973          | Great egret<br><i>Casmerodius<br/>albus</i>       | Eggs       | Albumen<br>and Yolk      | UK <sup>b</sup> | 0.24 to<br>0.43  | NR  | NR        | Stendell et al.<br>1976 |
| ON: L. St. Clair   | 1973          | Mallard<br><i>Anas<br/>platyrhynchos</i>          | Eggs       | Albumen<br>and Yolk      | UK <sup>b</sup> | 0.05 to<br>0.20  | NR  | NR        | Stendell et al.<br>1976 |
| ON: L. Superior<br>4 locations                                 | 1973-<br>1992 | Herring gull<br><i>Larus<br/>argentatus</i>       | Eggs       | Albumen<br>and Yolk      | Var             | 0.15 to<br>0.59  | NR  | NR        | Koster et al. 1996      |
| ON: L. Superior  | 1973          | Herring gull<br><i>Larus<br/>argentatus</i>       | Eggs       | Albumen<br>and Yolk      | UK <sup>b</sup> | 0.41 to<br>0.62  | NR  | NR        | Ryder 1974              |
| ON: L. Superior  | 1973          | Ring-billed gull<br><i>Larus<br/>delawarensis</i> | Eggs       | Albumen<br>and Yolk      | UK <sup>b</sup> | 0.45 to<br>0.75  | NR  | NR        | Ryder 1974              |
| ON: Lakes (12)<br>disconnected<br>from Wabigoon-<br>English R. | 1974-<br>1975 | Common Loon<br><i>Gavia immer</i>                 | Eggs       | Albumen<br>and Yolk      | 13              | $0.59 \pm 0.10$  | $0.58 \pm$<br>$0.10$                              | 99        | Barr 1985               |
| ON: Lakes (12)<br>disconnected<br>from Wabigoon-<br>English R. | 1976          | Common Loon<br><i>Gavia immer</i>                 | Adult      | Brain<br>Liver<br>Muscle | 10              | $0.4 \pm 0.1$<br>$8.8 \pm 4.6$<br>$1.2 \pm 0.6$                  | $0.2 \pm$<br>$0.4$                                | 4         | Barr 1985               |
| ON: Lakes (12)<br>disconnected<br>from Wabigoon-<br>English R. | 1976          | Common Loon<br><i>Gavia immer</i>                 | Chick      | Brain<br>Liver<br>Muscle | 4               | $0.4 \pm 0.1$<br>$0.8 \pm 0.3$<br>$0.4 \pm 0.1$                  | $0.7 \pm$<br>$0.2$                                | 87        | Barr 1985               |



## Canadian Tissue Residue Guidelines: Methylmercury

| Location  | Year          | Species                                     | Life Stage | Tissue                   | N   | Total Hg<br>$\mu\text{g}\cdot\text{g}^{-1}$ ww <sup>a</sup> | MeHg<br>$\mu\text{g}\cdot\text{g}^{-1}$<br>ww | %<br>MeHg | Reference          |
|---|---------------|---|------------|--------------------------|-----|---|---|-----------|--------------------|
| ON: Lakes (6)<br>downstream of<br>Wabigoon R,<br>Dryden         | 1974-<br>1975 | Common Loon<br><i>Gavia immer</i>           | Eggs       | Albumen<br>and Yolk      | 5   | 1.39 ± 0.55   | 1.34 ±<br>0.53                                | 97        | Barr 1985          |
| ON: Lakes (6)<br>downstream of<br>Wabigoon R,<br>Dryden         | 1976          | Common Loon<br><i>Gavia immer</i>           | Adult      | Brain<br>Liver<br>Muscle | 5   | 1.5 ± 0.6<br>29.7 ± 12.4<br>4.6 ± 1.6                       | 1.5 ±<br>2.1                                  | 9         | Barr 1985          |
| ON: Lakes (10)<br>upstream of<br>Maynard Falls<br>on English R. | 1974-<br>1975 | Common Loon<br><i>Gavia immer</i>           | Eggs       | Albumen<br>and Yolk      | 10  | 0.54 ± 0.09   | 0.53 ±<br>0.10                                | 97        | Barr 1985          |
| ON: Lakes (10)<br>upstream of<br>Maynard Falls<br>on English R. | 1976          | Common Loon<br><i>Gavia immer</i>           | Adult      | Brain<br>Liver<br>Muscle | 12  | 0.4 ± 0.1<br>5.1 ± 3.2<br>1.2 ± 0.5                         | 1.0 ±<br>0.7                                  | 27        | Barr 1985          |
| ON: Lakes (10)<br>upstream of<br>Maynard Falls<br>on English R. | 1976          | Common Loon<br><i>Gavia immer</i>           | Chick      | Brain<br>Liver<br>Muscle | 6   | 0.3 ± 0.1<br>0.9 ± 0.4<br>0.4 ± 0.2                         | 0.8 ±<br>0.4                                  | 92        | Barr 1985          |
| ON: Lakes (6)<br>Upstream of<br>Wabigoon-<br>English R.         | 1974-<br>1975 | Common Loon<br><i>Gavia immer</i>           | Eggs       | Albumen<br>and Yolk      | 6   | 0.72 ± 0.23   | 0.71 ±<br>0.22                                | 99        | Barr 1985          |
| ON: Lakes (6)<br>Upstream of<br>Wabigoon-<br>English R.         | 1976          | Common Loon<br><i>Gavia immer</i>           | Adult      | Brain<br>Liver<br>Muscle | 4   | 1.2 ± 0.6<br>16.7 ± 10.9<br>3.4 ± 2.0                       | 1.1 ±<br>2.2                                  | 7         | Barr 1985          |
| ON: Niagara R.<br>Un-named island                               | 1973-<br>1992 | Herring gull<br><i>Larus<br/>argentatus</i> | Eggs       | Albumen<br>and Yolk      | Var | 0.15 to<br>0.35   | NR  | NR        | Koster et al. 1996 |

## Canadian Tissue Residue Guidelines: Methylmercury

| Location                              | Year      | Species                                     | Life Stage | Tissue       | N  | Total Hg<br>$\mu\text{g}\cdot\text{g}^{-1}\text{ ww}^{\text{a}}$ | MeHg<br>$\mu\text{g}\cdot\text{g}^{-1}\text{ ww}$ | %<br>MeHg | Reference              |
|---------------------------------------|-----------|---|------------|--------------|----|--|---|-----------|------------------------|
| QC: Great Whale area, Northern Québec | 1989-1990 | Common merganser<br><i>Mergus merganser</i> | Adult      | Liver Muscle | 13 | 17.53 ± 12.06<br>1.27 ± 0.48                                     | NR  | NR        | Langlois et al. 1995   |
| QC: Great Whale area, Northern Québec | 1989-1990 | Herring gull<br><i>Larus argentatus</i>     | Adult      | Muscle       | 28 | 1.03 ± 0.80  | NR  | NR        | Langlois et al. 1995   |
| QC: Great Whale area, Northern Québec | 1989-1990 | Herring gull<br><i>Larus argentatus</i>     | Adult      | Liver        | 27 | 2.91 ± 2.35  | NR  | NR        | Langlois et al. 1995   |
| QC: Great Whale area, Northern Québec | 1989-1990 | Herring gull<br><i>Larus argentatus</i>     | Eggs       | NR           | 2  | 0.28 ± 0.12  | NR  | NR        | Langlois et al. 1995   |
| QC: Lakes & rivers North-west Québec  | 1989-1991 | Osprey<br><i>Pandion haliaetus</i>          | Eggs       | Whole        | 33 | 0.2 ± 0.1  | NR  | NR        | DesGranges et al. 1998 |
| QC: Lakes & rivers North-west Québec  | 1989-1991 | Osprey<br><i>Pandion haliaetus</i>          | Nestling   | Feather      | 63 | 7.0 ± 4.3  | NR  | NR        | DesGranges et al. 1998 |
| QC: Lakes & rivers North-west Québec  | 1989-1991 | Osprey<br><i>Pandion haliaetus</i>          | Adult      | Feather      | 29 | 16.5 ± 12.8  | NR  | NR        | DesGranges et al. 1998 |

## Canadian Tissue Residue Guidelines: Methylmercury

| Location   | Year          | Species  | Life Stage | Tissue             | N  | Total Hg<br>$\mu\text{g}\cdot\text{g}^{-1}$ ww <sup>a</sup> | MeHg<br>$\mu\text{g}\cdot\text{g}^{-1}$<br>ww | %<br>MeHg | Reference                 |
|--|---------------|--|------------|--------------------|----|---|---|-----------|---------------------------|
| QC: Lakes & rivers<br>North-west<br>Québec                                   | 1989-<br>1991 | Osprey<br><i>Pandion<br/>haliaetus</i>             | NR         | Stomach<br>Content | 6  | $0.3 \pm 0.2$   | NR  | NR        | DesGranges et al.<br>1998 |
| QC: Nottaway-<br>Broadback-<br>Rupert area,<br>South-east coast<br>James Bay | 1990-<br>1991 | Common<br>merganser<br><i>Mergus<br/>merganser</i> | Adult      | Muscle             | 19 | $1.41 \pm 1.26$   | NR  | NR        | Langlois et al.<br>1995   |
| QC: Nottaway-<br>Broadback-<br>Rupert area,<br>South-east coast<br>James Bay | 1990-<br>1991 | Common<br>merganser<br><i>Mergus<br/>merganser</i> | Adult      | Liver              | 15 | $10.9 \pm 7.5$  | NR  | NR        | Langlois et al.<br>1995   |
| QC: Nottaway-<br>Broadback-<br>Rupert area,<br>South-east coast<br>James Bay | 1990-<br>1991 | Herring gull<br><i>Larus<br/>argentatus</i>        | Adult      | Muscle             | 13 | $1.59 \pm 1.32$   | NR  | NR        | Langlois et al.<br>1995   |
| QC: Reservoirs<br>North-west<br>Québec                                       | 1989-<br>1991 | Osprey<br><i>Pandion<br/>haliaetus</i>             | Eggs       | Whole              | 18 | $0.2 \pm 0.1$   | NR  | NR        | DesGranges et al.<br>1998 |
| QC: Reservoirs<br>North-west<br>Québec                                       | 1989-<br>1991 | Osprey<br><i>Pandion<br/>haliaetus</i>             | Nestling   | Feather            | 78 | $37.4 \pm 20.1$   | NR  | NR        | DesGranges et al.<br>1998 |
| QC: Reservoirs<br>North-west<br>Québec                                       | 1989-<br>1991 | Osprey<br><i>Pandion<br/>haliaetus</i>             | Adult      | Feather            | 31 | $58.1 \pm 51.3$   | NR  | NR        | DesGranges et al.<br>1998 |

Canadian Tissue Residue Guidelines: Methylmercury

| Location                               | Year          | Species                                | Life Stage | Tissue             | N | Total Hg<br>$\mu\text{g}\cdot\text{g}^{-1}$ ww <sup>a</sup> | MeHg<br>$\mu\text{g}\cdot\text{g}^{-1}$<br>ww | %<br>MeHg | Reference                 |
|--|---------------|--|------------|--------------------|---|---|---|-----------|---------------------------|
| QC: Reservoirs<br>North-west<br>Québec | 1989-<br>1991 | Osprey<br><i>Pandion<br/>haliaetus</i> | NR         | Stomach<br>Content | 6 | $0.8 \pm 0.33$  | NR  | NR        | DesGranges et al.<br>1998 |

NR = Not Reported

Var = Variation in the number of samples

Notes:

<sup>a</sup> Unless noted otherwise, i.e., dw = dry weight.

<sup>b</sup> UK = unknown, information cited from Koster et al. 1996

**APPENDIX B-8: LEVELS OF TOTAL MERCURY AND METHYLMERCURY IN SELECTED NON-BIOLOGICAL MEDIA IN THE CANADIAN ENVIRONMENT.**

| Location                    | Year      | Media                             | N  | Total Hg<br>ng·L <sup>-1</sup> or µg·g <sup>-1</sup><br>dw <sup>a</sup> | MeHg<br>ng·L <sup>-1</sup> or µg·g <sup>-1</sup><br>dw | %<br>MeHg | Reference             |
|-----------------------------|-----------|-----------------------------------|----|---|--|-----------|-----------------------|
| ON/QC: Black L.             | 1987      | Water, lake                       | 5  | 3.5   | 1.3  | 37%       | Schintu et al. 1989   |
| ON/QC: Gatineau R.          | 1987      | Water, river                      | 5  | 11.4  | 4.1  | 36%       | Schintu et al. 1989   |
| ON/QC: Ottawa R.            | 1987      | Water, river                      | 5  | 9.0   | 2.3  | 26%       | Schintu et al. 1989   |
| ON: Experimental Lakes Area | 1992      | Rain, August & Sept. samples      | 4  | 4.62  | 0.017  | 0.4%      | St. Louis et al. 1995 |
| ON: Experimental Lakes Area | 1993      | Rain, May – August                | 6  | 5.33  | 0.049  | 0.9%      | St. Louis et al. 1995 |
| ON: Experimental Lakes Area | 1995      | Peat porewater                    | NR | 6   | 0.6  | 10%       | Moore et al. 1995     |
| ON: Experimental Lakes Area | 1993      | Snow on lake surface              | 2  | 0.95  | 0.008  | 0.8%      | St. Louis et al. 1995 |
| ON: Experimental Lakes Area | 1994      | Snow on lake surface              | 2  | 1.96  | 0.032  | 1.7%      | St. Louis et al. 1995 |
| ON: Experimental Lakes Area | 1992      | Snow, Oct. & Nov. samples         | 2  | 3.58  | 0.104  | 2.9%      | St. Louis et al. 1995 |
| ON: Experimental Lakes Area | 1991      | Water from upland area            | 9  | 13.11 ± 0.67  | 0.03 ± 0.006   | 0.2%      | St. Louis et al. 1994 |
| ON: Experimental Lakes Area | 1991      | Water from wetland dominated area | 19 | 5.02 ± 0.36   | 0.626 ± 0.126  | 13.3%     | St. Louis et al. 1994 |
| ON: Experimental Lakes Area | 1991      | Water from wetland/upland area#1  | 15 | 11.77 ± 0.73  | 0.176 ± 0.012  | 1.6%      | St. Louis et al. 1994 |
| ON: Experimental Lakes Area | 1991      | Water from wetland/upland area#2  | 9  | 11.40 ± 0.87  | 0.228 ± 0.033  | 2.0%      | St. Louis et al. 1994 |
| ON: Experimental Lakes Area | 1991-1993 | Water, Pond 979 before flooding   | NR | 2.6 ± 1.5   | 0.09 ± 0.04  | 4.4%      | Kelly et al. 1995     |

## Canadian Tissue Residue Guidelines: Methylmercury

| Location                                | Year                   | Media  | N   | Total Hg<br>ng·L <sup>-1</sup> or µg·g <sup>-1</sup><br>dw <sup>a</sup> | MeHg<br>ng·L <sup>-1</sup> or µg·g <sup>-1</sup><br>dw | %<br>MeHg | Reference             |
|---|------------------------|--|-----|---|--|-----------|-----------------------|
| ON: Experimental Lakes Area             | 1991-1993              | Water, Pond 979 after flooding                   | NR  | 4.5 ± 1.7   | 1.38 ± 0.51  | 31%       | Kelly et al. 1995     |
| ON: Experimental Lakes Area             | 1991                   | Water, upland/wetland & lake area                | 16  | 1.69 ± 0.27   | 0.040 ± 0.003  | 4.4%      | St. Louis et al. 1994 |
| ON: Experimental Lakes Area             | 1991                   | Wet deposition                                   | 5   | 3.81 ± 0.85   | 0.019 ± 0.004  | 0.6%      | St. Louis et al. 1994 |
| ON: St. Lawrence R., Cornwall area      | 1994                   | Surface sediments from riverine wetlands         | 69  | 0.015 to 0.882  | NR   | 1%        | Thompson 1996         |
| ON: Wabigoon/English R.                 | 1978-1979 <sup>b</sup> | Water, Clay L.                                   | 26  | NR  | 1.3  | NR        | Parks et al. 1989     |
| ON: Wabigoon/English R.                 | 1978-1979 <sup>b</sup> | Water, English R., Ball L. outflow               | 12  | NR  | 0.22   | NR        | Parks et al. 1989     |
| ON: Wabigoon/English R.                 | 1978-1979 <sup>b</sup> | Water, Wabigoon L.                               | 25  | NR  | 0.27   | NR        | Parks et al. 1989     |
| ON: Wabigoon/English R.                 | 1978-1979 <sup>b</sup> | Water, Wabigoon R., Clay L. outflow              | 50  | NR  | 1.3  | NR        | Parks et al. 1989     |
| ON: Wabigoon/English R.                 | 1978-1979 <sup>b</sup> | Water, Wabigoon R., inflow to Wainwright Dam     | 25  | NR  | 0.43   | NR        | Parks et al. 1989     |
| ON: Wetlands (22) South-central Ontario | 1993                   | Surface sediments from various types of wetlands | 198 | 0.081 to 0.316  | NR   | 1%        | Thompson 1996         |
| QC: Great Whale area, Northern Québec   | 1989-1990              | Sediments  | 12  | 0.07 ± 0.05   | NR   | NR        | Langlois et al. 1995  |
| QC: Great Whale area, Northern Québec   | 1989-1990              | Seston – fall samples                            | 7   | 0.11 ± 0.07   | 0.02 ± 0.02  | 18.2%     | Langlois et al. 1995  |
| QC: Great Whale area, Northern Québec   | 1989-1990              | Water – summer samples                           | 22  | 4.8 ± 6.62  | 0.453 ± 0.37   | 9.4%      | Langlois et al. 1995  |
| QC: Natural lakes, Northern Québec      | 1994-1996              | Water – dissolved Hg                             | NR  | 1.51 ± 0.06<br>0.4 - 2.6  | 0.049 ± 0.004<br>0.018 - 0.115                         | 3.2%      | Lucotte et al. 1998   |

Canadian Tissue Residue Guidelines: Methylmercury

| Location   | Year              | Media                          | N  | Total Hg<br>ng·L <sup>-1</sup> or µg·g <sup>-1</sup><br>dw <sup>a</sup> | MeHg<br>ng·L <sup>-1</sup> or µg·g <sup>-1</sup><br>dw | %<br>MeHg | Reference                 |
|--|-------------------|--------------------------------|----|---|--|-----------|---------------------------|
| QC: R. Bourassa and Laforge-1 Reservoirs               | 1994-1996         | Water                          | 87 | 2.35 ± 0.12<br>0.96 - 5.34  | 0.28 ± 0.02<br>0.03-0.85                               | 12%       | Thérien and Morrison 1998 |
| QC: Lakes near R. Bourassa and Laforge-1 Reservoirs    | 1994-1996         | Water                          | 30 | 1.51 ± 0.06<br>0.4 - 2.6  | 0.05 ± 0.004<br>0.02 - 0.11                            | 3%        | Thérien and Morrison 1998 |
| QC: La Grande 2 Reservoirs                             | 1992              | Soils, flooded, 0-4 cm depth   | NR | 0.129   | 0.0085   | 6.6%      | Mucci et al. 1995         |
| QC: La Grande 2 Reservoirs                             | 1992              | Soils, flooded, 11-19 cm depth | NR | 0.032   | 0.0019   | 5.9%      | Mucci et al. 1995         |
| QC: La Grande 2 Reservoirs                             | 1992              | Soils, flooded, 4-11 cm depth  | NR | 0.142   | 0.043  | 30%       | Mucci et al. 1995         |
| QC: Northern reservoirs/lake                           | 1992              | Sediments, surface             | NR | 0.036 to 0.059  | 0.0001 to 0.0006                                       | 1%        | Tremblay et al. 1996      |
| QC: Nottaway-Broadback-Rupert area, SE coast James Bay | 1990-1991         | Water – summer samples         | 25 | 3.07 ± 1.71   | 0.94 ± 0.54  | 31%       | Langlois et al. 1995      |
| QC: Nottaway-Broadback-Rupert area, SE coast James Bay | 1990-1991         | Water – winter samples         | 20 | 4.45 ± 1.73   | 0.46 ± 0.83  | 10.3%     | Langlois et al. 1995      |
| SK: Pasqua L.  | 1982 <sup>b</sup> | Water                          | 4  | 51.6  | 0.081  | 0.16%     | Jackson 1993              |

NR = Not Reported

Notes:

<sup>a</sup> ng·L<sup>-1</sup> is unit for water-based media; µg·g<sup>-1</sup> is unit for solid media.

<sup>b</sup> Samples taken at a time prior to realization that “ultra-clean” techniques were necessary.

**APPENDIX C-1: SUMMARY OF DATA ON THE ACUTE OR CHRONIC TOXICITY OF ORALLY-ADMINISTERED METHYLMERCURY IN MAMMALS.**

| Species  | Life Stage      | Sex | N  | Daily dose<br>mg·kg <sup>-1</sup> in<br>diet<br>µg·kg <sup>-1</sup> bw·d <sup>-1</sup> | Length of<br>exposure                                      | Total<br>dose<br>mg·kg <sup>-1</sup><br>bw | [Tissue]<br>mg·kg <sup>-1</sup><br>ww | Endpoint and Effects   | Reference                      |
|--|-----------------|-----|----|--|--|--|---------------------------------------|--|--------------------------------|
| Cat<br><i>Felis catus</i>                                | Adult           | NR  | NR | 0.25 (fed 5 of<br>7 d)<br>250  | 90 d   | 22.5<br>(16.1/cat)                         | NR                                    | First convulsions at<br>68d; mean survival<br>78d  | Eaton et al.<br>1980           |
| Cat<br><i>Felis catus</i>                                | Adult           | NR  | NR | 0.55±0.29<br>S.E.  | 188 d  | 18-29/cat                                  | 15.1<br>Muscle<br>170 Hair            | Behavioural effects<br>following tuna fish diet  | Houpt et al.<br>1988           |
| Cat<br><i>Felis catus</i>                                | Adult           | NR  | NR | 0.5  | 7-11 mos   | NR   | NR                                    | Proliferation of smooth<br>ER; degeneration of<br>hepatic mitochondria<br>(tuna fish diet)       | Chang et<br>al. 1974           |
| Cat<br><i>Felis catus</i>                                | Adult           | NR  | NR | 0.05, 0.14,<br>0.3, 0.76, 1.2,<br>2.9<br>3, 8.4, 20, 46,<br>74, 176                    | 39 weeks<br>(273 d)  | NR   | NR                                    | Neurologic damage at<br>46 µg·kg <sup>-1</sup> bw (0.76<br>mg·kg <sup>-1</sup> ) in diet of fish | Charbonne<br>au et al.<br>1976 |
| Crab-eating<br>macaque<br><i>Macaca<br/>fascicularis</i> | 0 - 7<br>years  | NR  | 5  | NR<br>50   | 7 years  | 128  | NR                                    | Lost sensitivity to<br>touch and tactile<br>response   | Rice 1989                      |
| Crab-eating<br>macaque<br><i>Macaca<br/>fascicularis</i> | 0 - 13<br>years | NR  | 5  | NR<br>50   | Exposed<br>age 0 to 7;<br>Not<br>exposed<br>age 7 to<br>13 | 128  | NR                                    | Clumsier and slower<br>to react than controls  | Rice 1989                      |



Canadian Tissue Residue Guidelines: Methylmercury

| Species  | Life Stage | Sex | N       | Daily dose<br>mg·kg <sup>-1</sup> in<br>diet<br>µg·kg <sup>-1</sup> bw·d <sup>-1</sup> | Length of<br>exposure | Total<br>dose<br>mg·kg <sup>-1</sup><br>bw | [Tissue]<br>mg·kg <sup>-1</sup><br>ww | Endpoint and Effects  | Reference                                |
|--|------------|-----|---------|--|-----------------------|--|---------------------------------------|---|--|
| Crab-eating<br>macaque<br><i>Macaca<br/>fascicularis</i> | Adult      | F   | 7       | NR<br>90   | 1 year                | 32.4                                       | NR                                    | 4/7 suffered<br>neurologic damage   | Burbacher<br>et al. 1984                 |
| Crab-eating<br>macaque<br><i>Macaca<br/>fascicularis</i> | Adult      | NR  | NR      | NR<br>0.4 – 50   | 150 d                 | 0.06 to<br>7.5                             | NR                                    | No clinical symptoms,<br>no difference in<br>cholinesterase activity                        | Petruccioli<br>and<br>Turillazzi<br>1991 |
| Harp seal<br><i>Phoca<br/>groenlandica</i>               | Adult      | NR  | NR      | 0.25<br>NR   | 60 d                  | NR   | 8.9 Blood                             | Decline in appetite<br>and body weight  | Ronald et<br>al. 1977                    |
| Harp seal<br><i>Phoca<br/>groenlandica</i>               | Adult      | NR  | NR      | 0.25<br>NR   | 90 d                  | NR   | 12.5<br>Blood                         | Reduced activity after<br>60 d  | Ronald et<br>al. 1977                    |
| Harp seal<br><i>Phoca<br/>groenlandica</i>               | Adult      | NR  | NR      | 25   | 26 d                  | NR   | 21.3<br>Blood                         | Lethargy; weight loss;<br>death on day 20 to 26   | Ronald et<br>al. 1977                    |
| Mice<br><i>Mus musculus</i><br>Strain: B6C3-<br>F1       | Adult      | M/F | 12<br>0 | 0.4<br>40  | 2 years               | 28.8                                       | NR                                    | NOAEL   | Mitsumori<br>et al. 1990                 |
| Mice<br><i>Mus musculus</i><br>Strain: B6C3-<br>F1       | Adult      | M/F | 12<br>0 | 2<br>170   | 2 years               | 122  | NR                                    | LOAEL - chronic<br>kidney disease   | Mitsumori<br>et al. 1990                 |
| Mice<br><i>Mus musculus</i><br>Strain: B6C3-<br>F1       | Adult      | F   | 60      | 10<br>830  | 2 years               | 598  | NR                                    | Posterior paralysis in<br>3/60 after 80 weeks,<br>brain necrosis; chronic<br>kidney disease | Mitsumori<br>et al. 1990                 |

## Canadian Tissue Residue Guidelines: Methylmercury

| Species  | Life Stage | Sex       | N  | Daily dose<br>mg·kg <sup>-1</sup> in<br>diet<br>µg·kg <sup>-1</sup> bw·d <sup>-1</sup> | Length of<br>exposure                | Total<br>dose<br>mg·kg <sup>-1</sup><br>bw       | [Tissue]<br>mg·kg <sup>-1</sup><br>ww | Endpoint and Effects   | Reference                |
|--|------------|-----------|----|--|--------------------------------------|--|---------------------------------------|--|--------------------------|
| Mice<br><i>Mus musculus</i><br>Strain: B6C3-<br>F1 | Adult      | M         | 60 | 10<br>830  | 2 years                              | 598  | NR                                    | Posterior paralysis in<br>33/60 after 59 weeks,<br>decreased bw gain;<br>brain necrosis; chronic<br>kidney disease | Mitsumori<br>et al. 1990 |
| Mink<br><i>Mustela vison</i>                       | Adult      | 12F<br>4M | 16 | 1.0<br>100 M; 180 F  | 11.5 wks<br>(81 d)                   | 8.1 M<br>14.6 F                                  | 15.3 F<br>Brain                       | 4/12 F died, dose cut<br>in half for remainder<br>(see next table row).<br>Brain lesions in F that<br>died.        | Wren et al.<br>1987a     |
| Mink<br><i>Mustela vison</i>                       | Adult      | 4F<br>3M  | 7  | 1.0 reduced to<br>0.5 <sup>a</sup><br>100 to 50 M;<br>180 to 90 F                      | 11.5 wks<br>@ 1.0<br>15 wks @<br>0.5 | 8.1 +5.3<br>= 13.4 M<br>14.6 +<br>9.5<br>=24.1 F | 18 M<br>Brain<br>4.2 F<br>Brain       | No clinical effects (C.<br>Wren, pers. com.) -<br>elevated Hg in<br>offspring livers<br>measured.                  | Wren et al.<br>1987a     |
| Mink<br><i>Mustela vison</i>                       | Adult      | NR        | NR | 5.0<br>500 M; 700 F <sup>b</sup>   | 30-37 d                              | 15-18.5<br>M<br>21-26 F                          | 25.2<br>Muscle<br>20 Brain            | LT <sub>50</sub> - Ataxia,<br>anorexia, paralysis,<br>death  | Aulerich et<br>al. 1974  |
| Mink<br><i>Mustela vison</i>                       | Adult      | F         | 10 | 0.25<br>38 <sup>c</sup>  | 120 d                                | 4.5  | 0.5 Brain                             | No clinical or<br>pathologic effect  | Wobeser et<br>al. 1976a  |
| Mink<br><i>Mustela vison</i>                       | Adult      | F         | 10 | 0.33<br>53   | 120 d                                | 6.4  | 3.4 Brain                             | No clinical or<br>pathologic effect<br>(NOAEL)   | Wobeser et<br>al. 1976a  |
| Mink<br><i>Mustela vison</i>                       | Adult      | F         | 5  | <0.1<br><16  | 93 d                                 | <1.5   | 0.2<br>Muscle<br>0.1 Brain            | Control  | Wobeser et<br>al. 1976b  |
| Mink<br><i>Mustela vison</i>                       | Adult      | F         | 5  | 1.1<br>176   | 93 d                                 | 16.4   | 7.8<br>Muscle<br>8.2 Brain            | LOAEL: pale livers,<br>nerve tissue lesions,<br>2/5 showed reduced<br>movement.                                    | Wobeser et<br>al. 1976b  |

Canadian Tissue Residue Guidelines: Methylmercury

| Species                          | Life Stage | Sex | N  | Daily dose<br>mg·kg <sup>-1</sup> in<br>diet<br>µg·kg <sup>-1</sup> bw·d <sup>-1</sup> | Length of<br>exposure | Total<br>dose<br>mg·kg <sup>-1</sup><br>bw | [Tissue]<br>mg·kg <sup>-1</sup><br>ww | Endpoint and Effects  | Reference                 |
|----------------------------------|------------|-----|----|--|-----------------------|--|---------------------------------------|---|---------------------------|
| Mink<br><i>Mustela vison</i>     | Adult      | F   | 5  | 1.8<br>288   | 79 d                  | 22.8                                       | 4.9<br>Muscle<br>18.1 Brain           | Anorexia, ataxia at 50-80 d; death at 59-79 d                                       | Wobeser et al. 1976b      |
| Mink<br><i>Mustela vison</i>     | Adult      | F   | 5  | 4.8<br>768   | 36 d                  | 27.6                                       | 14.1<br>Muscle<br>10.5 Brain          | Anorexia, ataxia at 23-32 d; death at 26-36 d                                       | Wobeser et al. 1976b      |
| Mink<br><i>Mustela vison</i>     | Adult      | F   | 5  | 8.3<br>1328  | 26 d                  | 34.5                                       | 17.4<br>Muscle<br>13.3 Brain          | Anorexia, ataxia at 16-21 d; death at 19-26 d                                       | Wobeser et al. 1976b      |
| Mink<br><i>Mustela vison</i>     | Adult      | F   | 5  | 15.0<br>2250   | 20 d                  | 45.0                                       | 22 Muscle<br>15.6 Brain               | Anorexia, ataxia at 16-18 d; death at 18-20 d                                       | Wobeser et al. 1976b      |
| Mink<br><i>Mustela vison</i>     | 6 mos.     | F   | 20 | 0.1<br>16 <sup>c</sup>   | 101 d                 | 16.2                                       | NR                                    | No effect   | Chamberland et al. 1996   |
| Mink<br><i>Mustela vison</i>     | 6 mos.     | F   | 20 | 0.5<br>80  | 101 d                 | 8.1  | NR                                    | No effect reported.   | Chamberland et al. 1996   |
| Mink<br><i>Mustela vison</i>     | 6 mos.     | F   | 20 | 0.9<br>144   | 101 d                 | 14.5                                       | NR                                    | Lethal to >50% between 80-101 days; signs of neurotoxicity; minor renal damage.     | Chamberland et al. 1996   |
| Otter<br><i>Lutra canadensis</i> | Adult      | M   | 3  | 2.0<br>90  | 159-213 d             | 14.6 to<br>21.4                            | 12 Muscle<br>13.3 Brain               | Lethal to 2/3; Anorexia in 1/3 and ataxia in 2/3 otters (survivors euthanized)      | O'Connor and Nielsen 1981 |
| Otter<br><i>Lutra canadensis</i> | Adult      | M   | 3  | 4.0<br>170   | 113-122 d             | 17.9 to<br>21.3                            | 16.3<br>Muscle<br>18.5 Brain          | Anorexia and ataxia in 3 of 3 otters (euthanized); nephrotic and neurologic lesions | O'Connor and Nielsen 1981 |

Canadian Tissue Residue Guidelines: Methylmercury

| Species  | Life Stage | Sex | N  | Daily dose<br>mg·kg <sup>-1</sup> in<br>diet<br>µg·kg <sup>-1</sup> bw·d <sup>-1</sup> | Length of<br>exposure | Total<br>dose<br>mg·kg <sup>-1</sup><br>bw | [Tissue]<br>mg·kg <sup>-1</sup><br>ww | Endpoint and Effects   | Reference                           |
|--|------------|-----|----|--|-----------------------|--|---------------------------------------|--|-------------------------------------|
| Otter<br><i>Lutra<br/>canadensis</i>                                 | Adult      | M   | 3  | 8.0<br>370   | 45 - 57 d             | 16.1 to<br>22.4                            | 15.2<br>Muscle<br>14 Brain            | Lethal to 1/3<br>(survivors<br>euthanized); nephrotic<br>and neurologic lesions      | O'Connor<br>and<br>Nielsen<br>1981  |
| Otter<br><i>Lutra<br/>canadensis</i>                                 | Adult      | M   | 2  | 0<br>0   | 192-229 d             | 0  | 0.77<br>Muscle<br>0.38 Brain          | Control  | O'Connor<br>and<br>Nielsen.<br>1981 |
| Rat<br><i>Rattus</i> spp.<br>Strain:<br>unknown                      | Adult      | F   | 25 | 2.5<br>100   | 2 years               | 73   | NR                                    | 56% survival at end,<br>6% lower weight gain<br>(60 weeks); 76%<br>survival controls | Verschuure<br>n et al.<br>1976      |
| Rat<br><i>Rattus</i> spp.<br>Strain:<br>unknown                      | Adult      | M   | 25 | 2.5<br>100   | 2 years               | 73   | NR                                    | 48% survival at end;<br>72% survival in<br>controls                                  | Verschuure<br>n et al.<br>1976      |
| Rat<br><i>Rattus<br/>norvegicus</i><br>Strain:<br>Sprague-<br>Dawley | Adult      | F   | 56 | NR<br>340  | 104 wks               | 248  | NR                                    | 30% survival at this<br>dosage   | Mitsumori<br>et al. 1983,<br>1984   |
| Rat<br><i>Rattus<br/>norvegicus</i><br>Strain:<br>Sprague-<br>Dawley | Adult      | M   | 56 | NR<br>280  | 104 wks               | 204  | NR                                    | 10% survival at this<br>dosage   | Mitsumori<br>et al. 1983,<br>1984   |

Canadian Tissue Residue Guidelines: Methylmercury

| Species  | Life Stage | Sex | N  | Daily dose<br>mg·kg <sup>-1</sup> in<br>diet<br>µg·kg <sup>-1</sup> bw·d <sup>-1</sup> | Length of<br>exposure | Total<br>dose<br>mg·kg <sup>-1</sup><br>bw | [Tissue]<br>mg·kg <sup>-1</sup><br>ww | Endpoint and Effects   | Reference                     |
|--|------------|-----|----|--|-----------------------|--|---------------------------------------|--|-------------------------------|
| Rat<br><i>Rattus<br/>norvegicus</i><br>Strain: Wistar<br>rat | Adult      | F   | 50 | NR<br>50   | 26 mos                | 3.9  | NR                                    | NOAEL - females  | Munro et<br>al. 1980          |
| Rat<br><i>Rattus<br/>norvegicus</i><br>Strain: Wistar<br>rat | Adult      | M   | 50 | NR<br>50   | 26 mos                | 3.9  | NR                                    | NOAEL - males  | Munro et<br>al. 1980          |
| Rat<br><i>Rattus<br/>norvegicus</i><br>Strain: Wistar<br>rat | Adult      | F   | 50 | NR<br>250  | 26 mos                | 19.5                                       | NR                                    | Reduced bw gains;<br>nerve damage  | Munro et<br>al. 1980          |
| Rat<br><i>Rattus<br/>norvegicus</i><br>Strain: Wistar<br>rat | Adult      | M   | 50 | NR<br>250  | 26 mos                | 19.5                                       | NR                                    | Overt neurotoxicity;<br>decreased<br>hemoglobin and<br>hematocrit, reduced<br>bw gains, mortality;<br>nerve damage; severe<br>kidney damage. | Munro et<br>al. 1980          |
| Rat<br><i>Rattus<br/>norvegicus</i><br>Strain: Wistar<br>rat | Adult      | F   | 4  | NR<br>5 - 50   | 4 d                   | NR   | NR                                    | Fed days 6-9 of<br>gestation; 4 mo old<br>offspring showed<br>operant behaviour<br>deficits - LOAEL 10<br>µg·kg <sup>-1</sup> bw/d           | Bornhause<br>n et al.<br>1980 |
| Rhesus<br>monkey<br><i>Macaca<br/>mulatta</i>                | Adult      | NR  | NR | NR<br>125  | 3.5 mos               | 13.1                                       | NR                                    | Liver and kidney<br>histological changes   | Rice et al.<br>1989           |

Canadian Tissue Residue Guidelines: Methylmercury

| Species                                | Life Stage | Sex | N  | Daily dose<br>mg·kg <sup>-1</sup> in<br>diet<br>µg·kg <sup>-1</sup> bw·d <sup>-1</sup> | Length of<br>exposure | Total<br>dose<br>mg·kg <sup>-1</sup><br>bw | [Tissue]<br>mg·kg <sup>-1</sup><br>ww | Endpoint and Effects                     | Reference           |
|--|------------|-----|----|--|-----------------------|--|---------------------------------------|--|---------------------|
| Rhesus monkey<br><i>Macaca mulatta</i> | Adult      | NR  | NR | NR<br>80   | 7-12 mos              | 17 to 29                                   | NR                                    | Liver and kidney<br>histological changes | Rice et al.<br>1989 |
| Rhesus monkey<br><i>Macaca mulatta</i> | Adult      | NR  | NR | NR<br>100  | 10 mos                | 30   | NR                                    | Liver normal; kidney<br>effects evident  | Rice et al.<br>1989 |
| Rhesus monkey<br><i>Macaca mulatta</i> | Adult      | NR  | NR | NR<br>80-100   | 15 mos                | 36 to 45                                   | NR                                    | Liver normal; kidney<br>effects evident  | Rice et al.<br>1989 |
| Rhesus monkey<br><i>Macaca mulatta</i> | Adult      | NR  | NR | NR<br>90   | 10 mos                | 27   | NR                                    | Liver normal; kidney<br>effects evident  | Rice et al.<br>1989 |

NR = Not Reported

Notes:

<sup>a</sup> 1 mg·kg<sup>-1</sup> fed daily for 11.5 weeks, then fed every other day for remainder of 6 month study (~15 weeks) (1.0 mg·kg<sup>-1</sup> daily dose caused mortality to 3 F; 6 animals in discomfort were euthanized).

<sup>b</sup> Calculated using food consumption rates of 0.1 and 0.14 kg·kg<sup>-1</sup> bw per day for males (M) and females (F), respectively (Aulerich et al. 1974)

<sup>c</sup> Calculated using a food consumption rate of 0.16 kg·kg<sup>-1</sup> bw per day for female captive mink (Bleavins and Aulerich 1981)

**APPENDIX C-2: SUMMARY OF DATA ON THE REPRODUCTIVE AND DEVELOPMENTAL TOXICITY OF ORALLY-ADMINISTERED METHYLMERCURY IN MAMMALS.**

| Species  | Life Stage | Sex      | N  | Daily dose<br>$\mu\text{g}\cdot\text{kg}^{-1}\text{ bw}$       | Duration of exposure                 | Total dose<br>$\text{mg}\cdot\text{kg}^{-1}\text{ bw}$ | Endpoint and Effects  | Reference                  |
|--|------------|----------|----|--|--------------------------------------|--|---|----------------------------|
| Crab-eating macaque<br><i>Macaca fascicularis</i>            | Adult      | F        | 11 | 50 - 70  | Reproductive cycle                   | NR   | Mothers dosed through pregnancy; Visual recognition deficits in 35d old infants   | Gunderson et al. 1986      |
| Crab-eating macaque<br><i>Macaca fascicularis</i>            | Adult      | F        | 7  | 50   | Reproductive cycle                   | NR   | 2/7 aborted - not statistically significant   | Burbacher et al. 1984      |
| Crab-eating macaque<br><i>Macaca fascicularis</i>            | Adult      | F        | 7  | 90   | Reproductive cycle                   |  | 2/7 aborted, 3/7 no conception, not statistically significant   | Burbacher et al. 1984      |
| Mink<br><i>Mustela vison</i>                                 | Adult      | 4F<br>3M | 7  | 1.0 reduced to 0.5 <sup>a</sup><br>100 to 50 M;<br>180 to 90 F | 11.5 wks @<br>1.0<br>15 wks @<br>0.5 | 8.1 +5.3<br>= 13.4 M<br>14.6 + 9.5<br>=24.1 F          | Fertility of adult male mink, % females whelped, number of kits born per female, kit survival to weaning were not affected    | Wren et al. 1987b          |
| Rat<br><i>Rattus norvegicus</i><br>Strain:<br>Sprague-Dawley | Adult      | F        | NR | 3900   | Reproductive cycle <sup>b</sup>      | NR   | No effect on maternal body weight, litter size, behaviour of dams or offspring; 4% lower bw in offspring; biochemical changes | Sundberg and Okarsson 1992 |

Canadian Tissue Residue Guidelines: Methylmercury

| Species  | Life Stage | Sex | N  | Daily dose<br>$\mu\text{g}\cdot\text{kg}^{-1}$ bw | Duration of exposure  | Total dose<br>$\text{mg}\cdot\text{kg}^{-1}$ bw | Endpoint and Effects  | Reference              |
|--|------------|-----|----|---|-----------------------|---|---|------------------------|
| Rat<br><i>Rattus norvegicus</i><br>Strain:<br>Wistar rat | Adult      | F   | NR | 10-50   | Days 6-9 of gestation | 0.04 - 0.2                                      | Decrease in operant behaviour of 4 mo old offspring; LOAEL $10 \mu\text{g}\cdot\text{kg}^{-1}$ bw | Bornhausen et al. 1980 |

NR = Not Reported

Notes:

<sup>a</sup>  $1 \text{ mg}\cdot\text{kg}^{-1}$  fed daily for 10-12 weeks, then every other day for remainder of 6 month study (daily dose caused mortality to 3; 6 animals in discomfort were euthanized).

<sup>b</sup> Fed to dam, 11 weeks prior to mating, during gestation and lactation; measured levels in various dam and pup tissues.



**APPENDIX C-3: SUMMARY OF DATA ON THE CARCINOGENIC AND MUTAGENIC EFFECTS OF ORALLY-ADMINISTERED METHYLMERCURY IN MAMMALS.**

| Species   | Life Stage | Sex | N  | Daily dose<br>$\mu\text{g}\cdot\text{kg}^{-1}\text{ bw}$ | Duration of exposure | Total dose<br>$\text{mg}\cdot\text{kg}^{-1}\text{ bw}$ | Endpoint and Effects   | Reference               |
|---|------------|-----|----|--|----------------------|--|--|-------------------------|
| Mice<br><i>Mus musculus</i><br>Strain: ICR                | Adult      | M   | 60 | 906<br>(10 ppm)  | 104 weeks            | 942  | Renal epithelial tumours in 13/29 ; none at lower doses at 58 weeks                  | Hirano et al. 1986      |
| Mice<br><i>Mus musculus</i><br>Strain: ICR                | Adult      | F   | 60 | 785<br>(10 ppm)  | 104 weeks            | 816  | Epithelial degeneration of renal proximal tubules; degeneration of the sciatic nerve | Hirano et al. 1986      |
| Rat<br><i>Rattus</i> spp.<br>Strain: unknown              | Adult      | M   | 25 | 0-100<br>2.5 ppm   | 2 years              | to 73  | No difference in tumours among all groups  | Verschuuren et al. 1976 |
| Rat<br><i>Rattus</i> spp.<br>Strain: unknown              | Adult      | F   | 25 | 0-100<br>2.5 ppm   | 2 years              | to 73  | No difference in tumours among all groups  | Verschuuren et al. 1976 |
| Rat<br><i>Rattus norvegicus</i><br>Strain: Sprague-Dawley | Adult      | M   | 56 | 280  | 130 weeks            | 255  | No tumours occurred.   | Mitsumori 1983, 1984    |
| Rat<br><i>Rattus norvegicus</i><br>Strain: Sprague-Dawley | Adult      | F   | 56 | 280  | 130 weeks            | 255  | No tumours occurred.   | Mitsumori 1983, 1984    |
| Rat<br><i>Rattus norvegicus</i><br>Strain: Sprague-Dawley | Adult      | M   | 30 | 8<br>0.2 ppm   | 2 years              | 6  | No difference in tumours between treatment and control. No other effects.            | Newberne et al. 1972    |
| Rat<br><i>Rattus norvegicus</i><br>Strain: Sprague-Dawley | Adult      | F   | 30 | 8<br>0.2 ppm   | 2 years              | 6  | No difference in tumours between treatment and control. No other effects.            | Newberne et al. 1972    |

**APPENDIX C-4: SUMMARY OF DATA ON THE ACUTE, CHRONIC OR REPRODUCTIVE TOXICITY OF METHYLMERCURY IN BIRDS.**

| Species                                       | Life Stage | Sex | N  | Dose<br>mg·kg <sup>-1</sup> in diet<br>µg·kg <sup>-1</sup> bw·d <sup>-1</sup> | Length of<br>exposure | Total dose<br>mg·kg <sup>-1</sup><br>bw | Tissue conc.<br>mg·kg <sup>-1</sup> ww | Endpoint and Effects  | Reference                   |
|---|------------|-----|----|---|-----------------------|---|--|---|-----------------------------|
| Black duck<br><i>Anas rubripes</i>            | Adult      | M/F | 26 | 3<br>(free choice)  | 196 d                 | NR                                      | 3-6 Brain<br>ducklings                 | Dose fed to adult pairs;<br>Fewer eggs laid; reduced<br>hatch success; Reduced<br>survival of offspring | Finley and<br>Stendell 1978 |
| Common loon<br><i>Gavia immer</i>             | Adult      | NR  | NR | >0.3 in perch<br>53 <sup>a</sup>  | Breeding<br>season    | NR                                      | 1.6 Muscle<br>0.5 Brain                | Reduced territory<br>establishment, reduced<br>egg laying   | Barr 1986                   |
| Common loon<br><i>Gavia immer</i>             | Adult      | NR  | NR | >0.36 in perch<br>72  | Breeding<br>season    | NR                                      | 2.3 Muscle<br>0.8 Brain                | Reduced territory<br>establishment, reduced<br>egg laying, no chicks<br>survived                        | Barr 1986                   |
| Double-crested<br>cormorant                   | Young      | NR  | NR | 0.56<br>NR  | Season                | NR                                      | 0.63 Brain                             | No effect observed.   | Wolfe and<br>Norman 1998    |
| Double-crested<br>cormorant                   | Adult      | NR  | NR | 0.56<br>NR  | Season                | NR                                      | NR                                     | No effect observed.   | Wolfe and<br>Norman 1998    |
| Great Blue<br>heron<br><i>Ardea herodias</i>  | Young      | NR  | NR | 0.56<br>73-210  | Season                | NR                                      | 0.35 Brain                             | No effect on growth rates<br>of young.  | Wolfe and<br>Norman 1998    |
| Great. Blue<br>heron<br><i>Ardea herodias</i> | Adult      | NR  | NR | 0.56<br>122   | Season                | NR                                      | NR                                     | Reproductive success<br>comparable to<br>uncontaminated colonies.                                       | Wolfe and<br>Norman 1998    |
| Mallard<br><i>Anas<br/>platyrhynchos</i>      | Eggs       | NR  | 80 | NR  | 18 d                  | 1-90<br>µg/eg<br>g                      | 0.05-0.53 /egg                         | Malformations at 1 µg;<br>decreased growth at 9 µg;<br>68% survival at 27 µg                            | Hoffman and<br>Moore 1979   |

Canadian Tissue Residue Guidelines: Methylmercury

| Species                              | Life Stage  | Sex    | N        | Dose<br>mg·kg <sup>-1</sup> in diet<br>µg·kg <sup>-1</sup> bw·d <sup>-1</sup> | Length of<br>exposure | Total dose<br>mg·kg <sup>-1</sup><br>bw | Tissue conc.<br>mg·kg <sup>-1</sup> ww | Endpoint and Effects   | Reference                    |
|--------------------------------------|---|--------|----------|---|-----------------------|---|--|--|------------------------------|
| Mallard<br><i>Anas platyrhynchos</i> | YOY   | M/F    | 48       | 0.5, 5, 15<br>(free choice)   | 84 d                  | NR                                      | NR                                     | Kidney abnormalities at 5 and 15 mg·kg <sup>-1</sup>   | Snelgrove-Hobson et al. 1988 |
| Mallard<br><i>Anas platyrhynchos</i> | Adult<br>1 <sup>st</sup><br>season                              | F<br>M | 30<br>9  | 0 (control)<br>6.4 <sup>b</sup>   | 21 weeks              | NR                                      | <0.05 Eggs                             | Egg production ceased after 43 weeks.  | Heinz 1974                   |
| Mallard<br><i>Anas platyrhynchos</i> | Adult<br>1 <sup>st</sup><br>season                              | F<br>M | 30<br>9  | 0.5 dw<br>86 <sup>c</sup>   | 21 weeks              | NR                                      | ~1 Eggs                                | Egg production ceased after 31 weeks. Smaller eggs than controls, ducklings hyper-responsive in avoidance tests  | Heinz 1974                   |
| Mallard<br><i>Anas platyrhynchos</i> | Adult<br>1 <sup>st</sup><br>season                              | F<br>M | 30<br>9  | 3.0 dw<br>530 <sup>d</sup>  | 21 weeks              | NR                                      | 6.5-9.2 Eggs                           | Egg production ceased after 21 weeks. Smaller eggs, fewer sound eggs, reduced hatching success, reduced hatching survival to 1 wk (no statistics), ducklings hyper-responsive in avoidance tests | Heinz 1974                   |
| Mallard<br><i>Anas platyrhynchos</i> | Adult<br>2 <sup>nd</sup><br>season<br>, 1 <sup>st</sup><br>gen. | F<br>M | 10<br>10 | 0 (control)<br>6.4 <sup>b</sup>   | 1.2-1.5<br>years      | NR                                      | <0.05 Eggs<br><0.07 Liver              |  | Heinz 1976a                  |

Canadian Tissue Residue Guidelines: Methylmercury

| Species                              | Life Stage  | Sex    | N        | Dose<br>mg·kg <sup>-1</sup> in diet<br>µg·kg <sup>-1</sup> bw·d <sup>-1</sup> | Length of<br>exposure    | Total dose<br>mg·kg <sup>-1</sup><br>bw | Tissue conc.<br>mg·kg <sup>-1</sup> ww   | Endpoint and Effects  | Reference   |
|--------------------------------------|---|--------|----------|---|--------------------------|---|--|---|-------------|
| Mallard<br><i>Anas platyrhynchos</i> | Adult<br>2 <sup>nd</sup><br>season<br>, 1 <sup>st</sup><br>gen. | F<br>M | 9<br>9   | 0.5 dw<br>83 <sup>e</sup>   | 1.2-1.5<br>years         | NR                                      | 0.8-1.1 Eggs<br>1.6 Liver<br>1.8 Kidney<br>0.8 Breast<br>0.5 Brain<br>0.6 Ovary<br>11.2 Feathers   | No consistent differences   | Heinz 1976a |
| Mallard<br><i>Anas platyrhynchos</i> | Adult<br>2 <sup>nd</sup><br>season                              | F<br>M | 10<br>10 | 3.0 dw<br>450 <sup>f</sup>  | 1.2-1.5<br>years         | NR                                      | 5.5-7.4 Eggs<br>11.1 Liver<br>14.7 Kidney<br>5.0 Breast<br>4.6 Brain<br>8.4 Ovary<br>68.7 Feathers | Reduced hatchling survival<br>to 1 wk, Brain lesions.<br>Ducklings hyper-<br>responsive in avoidance<br>tests | Heinz 1976a |
| Mallard<br><i>Anas platyrhynchos</i> | Adult<br>2 <sup>nd</sup> gen                                    | F<br>M | 14<br>14 | 0 (control)<br>6.4  | From 9<br>days of<br>age | NR                                      | <0.05 Eggs<br><0.05 Livers   |   | Heinz 1976b |

## Canadian Tissue Residue Guidelines: Methylmercury

| Species                              | Life Stage                    | Sex    | N        | Dose<br>mg·kg <sup>-1</sup> in diet<br>µg·kg <sup>-1</sup> bw·d <sup>-1</sup> | Length of exposure | Total dose<br>mg·kg <sup>-1</sup> bw | Tissue conc.<br>mg·kg <sup>-1</sup> ww   | Endpoint and Effects   | Reference              |
|--------------------------------------|-------------------------------|--------|----------|---|--------------------|--------------------------------------|--|--|------------------------|
| Mallard<br><i>Anas platyrhynchos</i> | Adult<br>2 <sup>nd</sup> gen  | F<br>M | 14<br>14 | 0.5 dw<br>73 <sup>g</sup>   | From 9 days of age | NR                                   | 0.9 Eggs<br>0.9 Liver<br>1.5 Kidney<br>0.7 Breast<br>0.4 Brain<br>0.5 Ovary<br>9.0 Feathers  | Adults consumed significantly more food than controls. Increase in % eggs laid outside the nest box. Decrease in number of normal 1 wk old ducklings (but not in % normal hatchlings surviving to 1 wk). Ducklings gained less weight. | Heinz 1976b            |
| Mallard<br><i>Anas platyrhynchos</i> | Adult<br>3 <sup>rd</sup> gen. | F<br>M | 14       | 0 (control)   | From 9 days of age | NR                                   | <0.05 Eggs<br>0.1-1.4 Liver  |  | Heinz 1979             |
| Mallard<br><i>Anas platyrhynchos</i> | Adult<br>3 <sup>rd</sup> gen  | F      | 14       | 0.5<br>67 <sup>h</sup>  | From 9 days of age | NR                                   | 0.86 Eggs<br>1.5 Liver<br>1.6 Kidney<br>0.8 Breast<br>0.6 Brain<br>0.6 Ovary<br>9.1 Feathers | Hens laid fewer sound eggs. Egg shells thinner. Offspring displayed impaired behaviour   | Heinz 1979             |
| Mallard<br><i>Anas platyrhynchos</i> | Adult                         | M      | 12       | 10 (4 ww)<br>1000   | 75 d               | 75                                   | NR   | Weak legs after 7 weeks; most barely walking and one dead at end of study.   | Heinz and Hoffman 1998 |
| Mallard<br><i>Anas platyrhynchos</i> | Adult                         | F      | 9        | 10 (4 ww)<br>1000   | 75 d               | 75                                   | NR   | Reduced egg hatch, reduced survival of young to 7 d, fewer young produced.   | Heinz and Hoffman 1998 |

Canadian Tissue Residue Guidelines: Methylmercury

| Species  | Life Stage | Sex | N | Dose<br>mg·kg <sup>-1</sup> in diet<br>µg·kg <sup>-1</sup> bw·d <sup>-1</sup> | Length of exposure | Total dose<br>mg·kg <sup>-1</sup> bw | Tissue conc.<br>mg·kg <sup>-1</sup> ww | Endpoint and Effects  | Reference     |
|--|------------|-----|---|---|--------------------|--------------------------------------|--|---|---------------|
| Ring-necked pheasant<br><i>Phasianus colchicus</i> | Adult      | F   | 7 | <sup>g</sup><br>NR  | 84 d               | 34                                   | 7.8 Liver                              | Weight loss; demyelination of spinal cord. Fewer eggs produced. Shell-less eggs; low hatch success. High chick mortality. | Fimreite 1971 |
| Ring-necked pheasant<br><i>Phasianus colchicus</i> | Adult      | F   | 7 | 4.5<br>NR   | 84 d               | 19                                   | 4.5 Liver                              | Weight loss; demyelination of spinal cord. Fewer eggs produced. Shell-less eggs; low hatch success.                       | Fimreite 1971 |

NR = Not Reported

Notes:

<sup>a</sup> Loons:  $0.3 \mu\text{g}\cdot\text{g}^{-1} \times 730 \text{ g}\cdot\text{d}^{-1} \cdot 4.13 \text{ kg}^{-1} \text{ bw} = 53 \mu\text{g}\cdot\text{kg}^{-1} \text{ bw}\cdot\text{d}^{-1}$  (CCME 1998).

<sup>b</sup> Calculated using a measured dose of  $0.05 \text{ mg}\cdot\text{kg}^{-1} \text{ dw}$  and a food ingestion rate for control mallards of  $0.128 \text{ kg}$  dry weight of food per kg body weight per day (Heinz 1974, 1979)

<sup>c</sup> Calculated using a measured dose of  $0.55 \text{ mg}\cdot\text{kg}^{-1} \text{ dw}$  and a food ingestion rate for treated mallards of  $0.156 \text{ kg}$  dry weight of food per kg body weight per day (Heinz 1974, 1979)

<sup>d</sup> Calculated using a measured dose of  $3.4 \text{ mg}\cdot\text{kg}^{-1} \text{ dw}$  and a food ingestion rate for treated mallards of  $0.156 \text{ kg}$  dry weight of food per kg body weight (Heinz 1974, 1979)

<sup>e</sup> Calculated using a measured dose of  $0.53 \text{ mg}\cdot\text{kg}^{-1} \text{ dw}$  and a food ingestion rate for treated mallards of  $0.156 \text{ kg}$  dry weight of food per kg body weight per day (Heinz 1974, 1979)

<sup>f</sup> Calculated using a measured dose of  $2.88 \text{ mg}\cdot\text{kg}^{-1} \text{ dw}$  and a food ingestion rate for treated mallards of  $0.156 \text{ kg}$  dry weight of food per kg body weight per day (Heinz 1976a, 1979)

<sup>g</sup> Calculated using a measured dose of  $0.47 \text{ mg}\cdot\text{kg}^{-1} \text{ dw}$  and a food ingestion rate for treated mallards of  $0.156 \text{ kg}$  dry weight of food per kg body weight per day (Heinz 1976b, 1979)

<sup>h</sup> Calculated using a measured dose of  $0.43 \text{ mg}\cdot\text{kg}^{-1} \text{ dw}$  and a food ingestion rate for treated mallards of  $0.156 \text{ kg}$  dry weight of food per kg body weight per day (Heinz 1976b, 1979)

<sup>i</sup> Fed contaminated grain because MeHg used as a fungicide on grains at the time (grains are no longer MeHg treated in Canada).

**APPENDIX D: HUMAN CONSUMPTION RESTRICTION CATEGORIES FOR METHYLMERCURY IN FISH FROM CANADIAN WATERS.**

Unrestricted Consumption - Fish may be consumed freely. Children under 15 and women of child-bearing age should eat fish only in this category. Fish in this category have concentrations less than or equal to **0.5  $\mu\text{g}\cdot\text{g}^{-1}$** .

Limited Consumption - Consumption restrictions depend upon the mercury level in the fish, how much fish is consumed and the length of time over which the fish is consumed.

*Group #1:* Fish in this group have mercury concentrations between **0.5 to 1.0  $\mu\text{g}\cdot\text{g}^{-1}$** . Short term consumption (for one week only) - anglers can eat 2300 grams (5.0 lb.) of fillets or about ten meals. Long term Consumption (fish consumed throughout the year) - anglers can eat 200 grams (0.5 lb.) of fillets or about one meal per week.

*Group #2:* Fish in this group have mercury concentrations between **1.0 to 1.5  $\mu\text{g}\cdot\text{g}^{-1}$** . Short term Consumption (one week only) - anglers can eat 1500 grams (3.0 lb.) of fillets or about 7 meals. Long term Consumption (Fish consumed throughout the year) - anglers can eat 140 grams (0.3 lb.) of fillets or about one small meal per week.

No Consumption- No Fish should be consumed from this category. Fish in this category have concentrations greater than **1.5  $\mu\text{g}\cdot\text{g}^{-1}$** .

Canadian Tissue Residue Guidelines: Methylmercury

Health Canada  
Health Protection Branch  
Bureau of Chemical Safety  
1st Floor East, Banting Building  
Postal Locator 2201B1  
Ottawa, Ontario  
K1A 0L2

April 7, 1998

Our *reference* number: ADDCP98032001

Ms. Brenda Miskimmin:  
Limnos Freshwater Consultants

Dear Ms. Miskimmin:

This is in reference to your fax and inquiry of March 19, 1998 regarding the 0.5 ppm guideline for mercury in fish.

This guideline was established in 1969 by the Health Protection Branch, and was based on the estimated consumption of commercial fish by eaters of that food commodity and on the available toxicology relating to mercury, and in particular, to methylmercury. Please note that this guideline relates only to fish which are marketed commercially, and that there are exemptions to this guideline. These exemptions are fresh and frozen fillets of shark, swordfish and tuna, where an advisory of limiting consumption of these species to, at most, one meal per week was recommended. The decision to exclude these species from the 0.5 ppm guideline was based on information relating to typical mercury levels in these species, and that strict application of the guideline would severely restrict the continued availability of these species in Canada to the consumer.

We would point out that there also exists a guideline of 0.2 ppm total mercury, which is recommended when fish constitutes a major subsistence food, as is the case for some aboriginal individuals in some smaller communities. This recommendation was issued by the Medical Services Branch in 1979, and reiterated in 1984.

The 0.5 ppm guideline was based on the provisional tolerable daily intake of methylmercury established by the FAO/WHO Joint Expert Committee on Food Additives (JECFA) in 1978, and reiterated in the 1988 JECFA meeting. It is listed in the 1998 edition of the **General Standard for Contaminants and Toxins in Foods** which is published by the Codex Alimentarius (copy attached).

We trust the above will be of assistance to you.

Yours truly

N.P. McEwen  
Additives & Contaminants Section  
Chemical Health Hazard Assessment Division

Attach.



1998

**CODEX ALIMENTARIUS GENERAL STANDARD FOR CONTAMINANTS AND TOXINS IN FOODS**

**ANNEX IV -B SITUATION REVIEW OF CONTAMINANTS AND TOXINS IN FOODS**

1.13 **Mercury**

(1) Mercury is a trace element which is found widespread in nature, due to volcanic and industrial activities. It has no established function in the body. Complex bound mercury (Hg) and in fish methylmercury are the predominant species. Mercury compounds are relatively toxic, especially organic bound mercury. Organic mercury compounds have been used in the past as fungicides, but are now generally forbidden for that use. Contamination of the environment with mercury compounds has been reduced in most countries by the discontinuation of agricultural uses and by adaptations in industrial processes and in waste management. In the past several tragic cases of poisonings have occurred due to mercury compounds, mostly with fish, due to environmental contamination, but also related to food use of treated cereals intended for sowing.

(2) The 1972 JECFA evaluation establishing a PTWI (tolerable weekly intake) of 5 µg/kg body weight for total mercury (out of which a maximum of 3.3 µg/kg may be methylmercury) was confirmed by the 1988 JECFA.

There are extensive analytical and intake data available, identifying fish and fishery products as the main source of mercury in the diet. Cereals have been contributing in the past too, in relation to the seed treatment with mercury compounds.

Mercury levels in plant products are usually very low. This is also valid for animal products (except fish), but higher levels might occur when fish meal is an important item of the animal feed.

The average daily intake of mercury is reported to be between a few to 20 µg, mainly varying with the amount of fishery products in the diet. For consumers with a high intake of fish, the intake may be higher, but in normal situations still considerably below the PTWI.

(3) Health problems may occur in contaminated areas and/or in relation to contaminated Products. There are indications that naturally occurring higher mercury levels in predatory marine fish are less toxic due to complexing with selenium.

(4) Many countries have national MLs, especially regarding mercury in fish, but sometimes also regarding other foods.

There is a WHO guideline value of 0.001 mg/L for total mercury in drinking water

(5) The exposure of the general population and of groups at risk may be reduced by prohibiting fishing and sale of fish from heavily contaminated waters. Advice to consumers about fish consumption may also be useful in specific situations.

(6) Codex standards exist for mercury in fish. Further action is desirable on that point regarding the definition of the residues and regarding the specification of

## Canadian Tissue Residue Guidelines: Methylmercury

predatory fish. A detailed position paper is necessary as a basis for any other future action regarding possible MLs for mercury in other foods. Presently available information does not seem to justify establishing of MLS in other food commodities.