



Health
Canada

Santé
Canada

Tuberculosis

Drug resistance in Canada

2000

Reported susceptibility results of the
Canadian Tuberculosis Laboratory
Surveillance System

Our mission is to help the people of Canada maintain and improve their health.

Health Canada

► HOW TO REACH US

For more information, copies of this report or other reports, please contact:

**Tuberculosis Prevention and Control
Centre for Infectious Disease Prevention and Control
Population and Public Health Branch
Health Canada
Room 0108 B, Brooke Claxton Building
Tunney's Pasture, Ottawa, Ontario K1A 0L2**

Internal Postal Address: 0900B-1

Telephone: (613) 941-0238

Facsimile: (613) 946-3902

This report can also be accessed on the internet at:

<http://www.hc-sc.gc.ca/hpb/lcdc>

The following text, figures and tables were prepared by:

Howard Njoo, MD, MHSc, FRCPC
Director
Tuberculosis Prevention and Control

Penny Nault
Tuberculosis Database Manager
Tuberculosis Prevention and Control

Melissa Phipers, MSc
Senior Epidemiologist
Tuberculosis Prevention and Control

► ACKNOWLEDGEMENT

Tuberculosis Prevention and Control would like to acknowledge the members of the Canadian Tuberculosis Laboratory Technical Network and their teams for their contribution to and their participation in the Canadian Tuberculosis Laboratory Surveillance System (CTBLSS).

Published by authority of the Minister of Health

© Minister of Public Works and Government Services Canada 2001

Cat. H49-110/2000
ISBN 0-662-65720-9

This publication can be made available in alternative forms.



Tuberculosis

Drug resistance in Canada

2000

**Reported susceptibility results of the
Canadian Tuberculosis Laboratory
Surveillance System**

TABLE OF CONTENTS

- ▶ **INTRODUCTION.** 1

- ▶ **METHODOLOGY** 1

- ▶ **RESULTS** 2

- ▶ **DISCUSSION** 3

- ▶ **FIGURES**
 - Figure 1.** Reported TB drug resistance in Canada by province/territory – 2000 3
 - Figure 2.** Reported MTB isolates in Canada by province/territory – 2000 4
 - Figure 3.** Overall pattern of reported TB drug resistance in Canada – 2000 4
 - Figure 4.** Reported TB drug resistance in Canada by type of drug – 2000 5
 - Figure 5.** Reported TB drug resistance in Canada by gender and year of birth – 2000 5

- ▶ **TABLES**
 - Table A.** Concentrations for routine testing of first-line anti-tuberculosis drugs 2
 - Table 1.** Overall pattern of reported TB drug resistance in Canada – 1998-2000 6
 - Table 2.** Reported MTB isolates by “reporting” and “originating” province/territory, Canada – 2000 7
 - Table 3.** Reported MDR-TB isolates by province/territory, Canada – 2000. 8
 - Table 4.** Reported TB drug resistance by gender and year of birth, Canada – 2000 9
 - Table 5.** Reported results for routine drug susceptibility testing of MTB isolates to first-line anti-tuberculosis drugs, Alberta – 1998-2000 10
 - Table 6.** Reported results for routine drug susceptibility testing of MTB isolates to first-line anti-tuberculosis drugs, British Columbia – 1998-2000 10
 - Table 7.** Reported results for routine drug susceptibility testing of MTB isolates to first-line anti-tuberculosis drugs, Manitoba – 1998-2000 11
 - Table 8.** Reported results for routine drug susceptibility testing of MTB isolates to first-line anti-tuberculosis drugs, New Brunswick – 1998-2000 11

Table 9. Reported results for routine drug susceptibility testing of MTB isolates to first-line anti-tuberculosis drugs, Newfoundland – 1998-2000	11
Table 10. Reported results for routine drug susceptibility testing of MTB isolates to first-line anti-tuberculosis drugs, Northwest Territories – 1998-2000	12
Table 11. Reported results for routine drug susceptibility testing of MTB isolates to first-line anti-tuberculosis drugs, Nova Scotia – 1998-2000	12
Table 12. Reported results for routine drug susceptibility testing of MTB isolates to first-line anti-tuberculosis drugs, Nunavut – 1998-2000	12
Table 13. Reported results for routine drug susceptibility testing of MTB isolates to first-line anti-tuberculosis drugs, Ontario – 1998-2000	13
Table 14. Reported results for routine drug susceptibility testing of MTB isolates to first-line anti-tuberculosis drugs, Prince Edward Island – 1998-2000	13
Table 15. Reported results for routine drug susceptibility testing of MTB isolates to first-line anti-tuberculosis drugs, Québec – 1998-2000	14
Table 16. Reported results for routine drug susceptibility testing of MTB isolates to first-line anti-tuberculosis drugs, Saskatchewan – 1998-2000	14
Table 17. Reported results for routine drug susceptibility testing of MTB isolates to first-line anti-tuberculosis drugs, Yukon Territory – 1998-2000	15

▶ LIMITATIONS	16
--------------------------------	----

▶ CONCLUSIONS	16
--------------------------------	----

▶ REFERENCE	16
------------------------------	----

▶ APPENDICES

Appendix 1 – Participating Laboratories of the Canadian Tuberculosis Laboratory Surveillance System (CTBLSS).	17
Appendix 2 – <i>M. tuberculosis</i> Complex Antimicrobial Susceptibility Reporting Form	22

► INTRODUCTION

Tuberculosis Prevention and Control (TBPC) at the Centre for Infectious Disease Prevention and Control, Health Canada, in collaboration with the Canadian Tuberculosis Laboratory Technical Network and participating laboratories (representing all provinces and territories) in the Canadian Tuberculosis Laboratory Surveillance System (CTBLSS) (Appendix 1), established a laboratory-based national surveillance system in 1998 to monitor tuberculosis (TB) drug resistance patterns in Canada.

Laboratories report their results on anti-tuberculosis drug susceptibility testing to TBPC for every patient for whom a specimen or an isolate has been received for each calendar year. TBPC subsequently produces an annual report. This report presents 2000 and adjusted 1998 and 1999 (to reflect duplicate removal and late reporting) drug susceptibility data for TB isolates across Canada as of March 6, 2001.

► METHODOLOGY

A computerized database containing drug susceptibility test results of *Mycobacterium tuberculosis* (MTB) and MTB complex (MTBC) isolates is maintained at TBPC at the Centre for Infectious Disease Prevention and Control. Data are collected either through manual completion and mailing of a standard reporting form (Appendix 2) or by electronic transmission. Information collected includes gender, year of birth, province/territory from which the report originates, province/territory from which the specimen originates and susceptibility results. Every effort is made to eliminate duplicate specimens; only the most recent susceptibility results for a given patient in the current reporting year are included for analysis.

Manitoba, Ontario and Newfoundland identify the species and test all isolates for drug resistance in their respective provinces. Some provinces identify the species and test their own isolates and those of other provinces/territories (British Columbia: British Columbia and Yukon Territory isolates; Alberta: Alberta, Northwest Territories and some Nunavut isolates; Quebec: Quebec, New Brunswick, Northwest Territories and Nunavut isolates; Nova Scotia: Nova Scotia and Prince Edward Island isolates). Saskatchewan tests for drug resistance on all MTBC isolates; other provinces and territories report results at the species level, excluding *M. bovis* BCG.

Routine susceptibility testing of MTB or MTBC to first-line anti-tuberculosis drugs is generally performed using the radiometric proportion method (Bactec[®]). Table A lists the first-line anti-tuberculosis drugs and the concentrations in mg/L used by the participating laboratories. Results of susceptibility testing for second-line anti-tuberculosis drugs are not included in this report.

As noted in Table A, the number and specific first-line anti-tuberculosis drugs for which routine susceptibility testing is carried out differ among the provinces and territories. Accordingly, the number of isolates included in the descriptive analyses that were carried out vary. Analyses were performed using SAS version 6.12.

Table A: Concentrations for routine testing of first-line anti-tuberculosis drugs

Anti-tuberculosis drugs	mg/L	Comments
Isoniazid (INH)	0.1	
Rifampin (RMP)	2.0	
Ethambutol (EMB)	2.5	British Columbia uses a concentration of 4.0 mg/L.
Streptomycin (SM)	2.0	Routine testing is not performed for isolates from Quebec, Nova Scotia, New Brunswick, Prince Edward Island and for Nunavut isolates tested in Quebec.
Pyrazinamide (PZA)	100.0	Routine testing is not performed for isolates from British Columbia, Saskatchewan and the Yukon Territory.

► RESULTS

In 2000, participating laboratories across Canada reported drug susceptibility results for 1,468 isolates of MTBC. Eight *M. bovis* isolates were reported: four isolates of *M. bovis* BCG and four isolates of *M. bovis* other than BCG. Only the four (0.3%) *M. bovis* other than BCG isolates (two each from Quebec and Ontario) are included in the analyses, leaving a total of 1,464 isolates. The majority of isolates originated from Ontario, Quebec, British Columbia, Alberta and Manitoba.

Of the 1,464 isolates in 2000 included for analysis, 164 (11.2%) were resistant to one or more first-line anti-tuberculosis drug(s). Resistance to INH was the most common type of drug resistance (7.4%). A total of 14 isolates (1.0%) were multi-drug resistant tuberculosis (MDR-TB) strains (defined as resistance to at least INH and RMP); of which five isolates demonstrated resistance to four or five first-line anti-tuberculosis drugs tested. These isolates were reported from Ontario and British Columbia. In addition, British Columbia, Alberta, Saskatchewan, Manitoba and Ontario reported isolates with other patterns of multi-resistance. Six provinces and territories (Northwest Territories, Yukon, Newfoundland, Prince Edward Island, Nova Scotia and New Brunswick) reported that all isolates tested were susceptible to all the first-line anti-tuberculosis drugs.

Demographic information on the individual patients from whom the isolates originated is limited in this laboratory-based surveillance system. Of the 1,402 isolates for which the year of birth was known, 61% reported a year of birth between 1941 and 1980. Among the 156 drug resistant isolates for which year of birth was known, 71% reported a year of birth between 1941 and 1980. Males accounted for 56% of all the isolates and 64% of the drug resistant isolates for which gender was reported.


► DISCUSSION

The number of reported TB isolates in 2000 increased by 3.4% from the previous year (1,414 to 1,464 isolates). The percentage of isolates demonstrating any type of drug resistance decreased from 12.1% in 1999 to 11.2% in 2000 and the proportion of isolates classified as MDR-TB decreased from 1.2% in 1999 to 1.0% in 2000.


Over 90% of the reported laboratory TB isolates in Canada in 2000 originated from five provinces. The three largest provinces (Ontario, Quebec and British Columbia) consistently reported the majority of isolates and MDR-TB in the three years of data collection. Since the initiation of this laboratory-based surveillance system that began January 1, 1998, Saskatchewan, the Atlantic Provinces and the Territories have not reported any MDR-TB cases.

The results observed to date in this surveillance system are consistent with international data. In the latest report of the global TB drug resistance surveillance project jointly conducted by the World Health Organization (WHO) and the International Union Against Tuberculosis and Lung Disease (IUATLD), the median prevalence of overall TB drug resistance among the participating countries was 11.1% (as compared to 11.2% for Canada) and the median prevalence of MDR-TB was 1.8%¹ (as compared to 1.0% for Canada).


► **Figure 1**
Reported TB drug resistance in Canada by province/territory – 2000 (n = 1,464)




► **Figure 2**
 Reported MTB isolates in Canada by province/territory – 2000 (n = 1,464)



► **Figure 3**
 Overall pattern of reported TB drug resistance in Canada – 2000
 (n = 164/1,464 total isolates)



► **Figure 4**
 Reported TB drug resistance in Canada by type of drug – 2000 (n = 164/1,464 total isolates)



* SM and PZA are not part of routine first-line drug testing in some provinces/territories.

► **Figure 5**
 Reported TB drug resistance in Canada by gender and year of birth – 2000
 (n = 164/1,464 total isolates)




Table 1. Overall pattern of reported TB drug resistance in Canada – 1998-2000

	1998 Total (%)	1999 Total (%)	2000 Total (%)
Total number of isolates tested	1,461 (100.0)	1,414 (100.0)	1,464 (100.0)
Isolates susceptible	1,288 (88.2)	1,243 (87.9)	1,300 (88.8)
Any resistance to INH	123 (8.4)	126 (8.9)	108 (7.4)
Any resistance to RMP	19 (1.3)	19 (1.1)	17 (1.2)
Any resistance to EMB	22 (1.5)	20 (1.4)	20 (1.4)
Any resistance to SM**	82 (5.7)	72 (6.5)	63 (5.5)
Any resistance to PZA**	23 (2.0)	27 (2.4)***	24 (2.1)***
Resistance to one or more drugs	173 (11.8)	171 (12.1)	164 (11.2)
 Monoresistance**	116 (7.9)	113 (8.0)	119 (8.1)
 MDR-TB*	18 (1.2)	17 (1.2)	14 (1.0)
 Other patterns	39 (2.7)	41 (2.9)	31 (2.1)

* MDR-TB is defined as resistance to at least INH and RMP.

** Because SM and PZA are not part of the routine first-line drugs in some provinces and territories, denominators for any resistance to these drugs have been modified to reflect this (1998: SM n = 1,450 and PZA n = 1,174) (1999: SM n = 1,110 and PZA n = 1,130) (2000: SM n = 1,142 and PZA n = 1,144).

*** Includes 2 *M. bovis* isolates for 1999 and 4 *M. bovis* isolates for 2000.

Table 2. Reported MTB isolates by “reporting” and “originating” province/territory, Canada – 2000

Reporting Province	Originating Province/Territory													
	CANADA	Alta.	B.C.	Man.	N.B.	Nfld.	N.S.	N.W.T.	Nun.	Ont.	P.E.I.	Que.	Sask.	Y.T.
Number of isolates	1,464	104	252	101	9	10	4	8	29	599	3	278	64	3
Alta.	113	104	-	-	-	-	-	8	1	-	-	-	-	-
B.C.	255	-	252	-	-	-	-	-	-	-	-	-	-	3
Man.	100	-	-	100	-	-	-	-	-	-	-	-	-	-
Nfld.	10	-	-	-	-	10	-	-	-	-	-	-	-	-
N.S.	6	-	-	-	-	-	3	-	-	-	3	-	-	-
Ont.	599	-	-	-	-	-	-	-	-	599	-	-	-	-
Que.	316	-	-	-	9	-	1	-	28	-	-	278	-	-
Sask.	65	-	-	1	-	-	-	-	-	-	-	-	64	-

Table 3. Reported MDR-TB* isolates by province/territory, Canada – 2000

	Originating Province/Territory													
	CANADA	Alta.	B.C.	Man.	N.B.	Nfld.	N.S.	N.W.T.	Nun.	Ont.	P.E.I.	Que.	Sask.	Y.T.
Total number of isolates tested	1,464	104	252	101	9	10	4	8	29	599	3	278	64	3
Total number of MDR-TB* isolates	14	-	4	-	-	-	-	-	-	9	-	1	-	-
INH & RMP	1	-	-	-	-	-	-	-	-	1	-	-	-	-
INH, RMP & EMB	4	-	1	-	-	-	-	-	-	2	-	1	-	-
INH, RMP & SM	4	-	1	-	-	-	-	-	-	3	-	-	-	-
INH, RMP, EMB & SM	4	-	2	-	-	-	-	-	-	2	-	-	-	-
INH, RMP, SM & PZA	1	-	-	-	-	-	-	-	-	1	-	-	-	-

* MDR-TB is defined as resistance to at least INH and RMP.

Table 4. Reported TB drug resistance by gender and year of birth, Canada – 2000						
Year of Birth	Number of isolates		Any type of resistance		MDR-TB*	
	No.	(%)	No.	(%)		
Total Isolates	1464	(100.0)	164	(11.2)	14	(1.0)
≤ 1920	Males	68 (4.6)	7 (0.5)		-	(0.0)
	Females	56 (3.8)	1 (0.1)		-	(0.0)
	Unknown	3 (0.2)	- (0.0)		-	(0.0)
	Total	127 (8.7)	8 (0.5)		-	(0.0)
1921-1940	Males	187 (12.8)	20 (1.4)		2	(0.1)
	Females	148 (10.1)	11 (0.8)		1	(0.1)
	Unknown	9 (0.6)	- (0.0)		-	(0.0)
	Total	344 (23.5)	31 (2.1)		3	(0.2)
1941-1960	Males	217 (14.8)	29 (2.0)		3	(0.2)
	Females	135 (9.2)	13 (0.9)		-	(0.0)
	Unknown	17 (1.2)	5 (0.3)		-	(0.0)
	Total	369 (25.2)	47 (3.2)		3	(0.2)
1961-1980	Males	258 (17.6)	37 (2.5)		3	(0.2)
	Females	207 (14.1)	24 (1.6)		3	(0.2)
	Unknown	16 (1.1)	2 (0.1)		-	(0.0)
	Total	481 (32.9)	63 (4.3)		6	(0.4)
1981-2000	Males	35 (2.4)	3 (0.2)		-	(0.0)
	Females	46 (3.1)	4 (0.3)		-	(0.0)
	Unknown	- (0.0)	- (0.0)		-	(0.0)
	Total	81 (5.5)	7 (0.5)		-	(0.0)
Unknown	Males	28 (1.9)	3 (0.2)		-	(0.0)
	Females	19 (1.3)	3 (0.2)		1	(0.1)
	Unknown	15 (1.0)	2 (0.1)		1	(0.1)
	Total	62 (4.2)	8 (0.5)		2	(0.1)
Total	Males	793 (54.2)	99 (6.8)		8	(0.5)
	Females	611 (41.7)	56 (3.8)		5	(0.3)
	Unknown	60 (4.1)	9 (0.6)		1	(0.1)

* MDR-TB is defined as resistance to at least INH and RMP.

Table 5. Reported results for routine drug susceptibility testing of MTB isolates to first-line anti-tuberculosis drugs, Alberta – 1998-2000

	1998 Total (%)	1999 Total (%)	2000 Total (%)
Total number of isolates tested for INH, RMP, EMB, SM and PZA	119 (100.0)	117 (100.0)	104 (100.0)
Isolates susceptible	107 (89.9)	110 (94.0)	92 (88.5)
Isolates resistant to one or more drugs	12 (10.1)	7 (6.0)	12 (11.5)
Monoresistance	9 (7.6)	6 (5.1)	7 (6.7)
INH	4 (3.4)	2 (1.7)	2 (1.9)
SM	5 (4.2)	4 (3.4)	3 (2.9)
EMB	- (0.0)	- (0.0)	1 (1.0)
PZA	- (0.0)	- (0.0)	1 (1.0)
MDR-TB*	1 (0.8)	0 (0.0)	0 (0.0)
INH, RMP, EMB, SM & PZA	1 (0.8)	0 (0.0)	- (0.0)
Other Patterns	2 (1.7)	1 (0.9)	5 (4.8)
INH & SM	1 (0.8)	1 (0.9)	3 (2.9)
INH, SM & EMB	- (0.0)	- (0.0)	1 (1.0)
INH, SM & PZA	1 (0.8)	0 (0.0)	1 (1.0)

* MDR-TB is defined as resistance to at least INH and RMP.

Table 6. Reported results for routine drug susceptibility testing of MTB isolates to first-line anti-tuberculosis drugs, British Columbia – 1998-2000

	1998 Total (%)	1999 Total (%)	2000 Total (%)
Total number of isolates tested for INH, RMP, EMB and SM**	237 (100.0)	245 (100.0)	252 (100.0)
Isolates susceptible***	212 (89.5)	225 (91.8)	224 (88.9)
Isolates resistant to one or more drugs	25 (10.5)	20 (8.2)	28 (11.1)
Monoresistance	17 (7.2)	15 (6.1)	21 (8.3)
INH	14 (5.9)	11 (4.5)	12 (4.8)
SM	2 (0.8)	2 (0.8)	8 (3.2)
RMP	1 (0.4)	1 (0.4)	1 (0.4)
EMB	- (0.0)	1 (0.4)	- (0.0)
MDR-TB*	2 (0.8)	1 (0.4)	4 (1.6)
INH, RMP & SM	1 (0.4)	- (0.0)	1 (0.4)
INH, RMP & EMB	- (0.0)	- (0.0)	1 (0.4)
INH, RMP, EMB & SM	1 (0.4)	1 (0.4)	2 (0.8)
Other Patterns	6 (2.5)	4 (1.6)	3 (1.2)
INH & SM	5 (2.1)	2 (0.8)	1 (0.4)
INH & EMB	1 (0.4)	1 (0.4)	- (0.0)
INH, SM & EMB	- (0.0)	1 (0.4)	2 (0.8)

* MDR-TB is defined as resistance to at least INH and RMP.
 ** Routine testing for PZA not conducted in British Columbia.
 *** Includes 1 *M. bovis* isolate (1999).

Table 7. Reported results for routine drug susceptibility testing of MTB isolates to first-line anti-tuberculosis drugs, Manitoba – 1998-2000

	1998 Total (%)	1999 Total (%)	2000 Total (%)
Total number of isolates tested for INH, RMP, EMB, SM and PZA	106 (100.0)	100 (100.0)	101 (100.0)
Isolates susceptible	98 (92.5)	89 (89.0)	93 (92.1)
Isolates resistant to one or more drugs	8 (7.5)	11 (11.0)	8 (7.9)
Monoresistance	4 (3.8)	6 (6.0)	6 (5.9)
INH	2 (1.9)	3 (3.0)	6 (5.9)
SM	2 (1.9)	3 (3.0)	- (0.0)
MDR-TB*	2 (1.9)	2 (2.0)	- (0.0)
INH & RMP	- (0.0)	1 (1.0)	- (0.0)
INH, RMP & EMB	1 (0.9)	- (0.0)	- (0.0)
INH, RMP, SM & PZA	- (0.0)	1 (1.0)	- (0.0)
INH, RMP, EMB, SM & PZA	1 (0.9)	- (0.0)	- (0.0)
Other Patterns	2 (1.9)	3 (3.0)	2 (2.0)
INH & SM	2 (1.9)	1 (1.0)	2 (2.0)
INH, SM & EMB	- (0.0)	1 (1.0)	- (0.0)
INH, SM & PZA	- (0.0)	1 (1.0)	- (0.0)

* MDR-TB is defined as resistance to at least INH and RMP.

Table 8. Reported results for routine drug susceptibility testing of MTB isolates to first-line anti-tuberculosis drugs, New Brunswick – 1998-2000

	1998 Total (%)	1999 Total (%)	2000 Total (%)
Total number of isolates tested for INH, RMP, EMB and PZA*	9 (100.0)	12 (100.0)	9 (100.0)
Isolates susceptible	8 (88.9)	12 (100.0)	9 (100.0)
Isolates resistant to one or more drugs	1 (1.1)	- (0.0)	- (0.0)
Monoresistance	1 (1.1)	- (0.0)	- (0.0)
INH	1 (1.1)	- (0.0)	- (0.0)

* Routine testing for SM not conducted in New Brunswick.

Table 9. Reported results for routine drug susceptibility testing of MTB isolates to first-line anti-tuberculosis drugs, Newfoundland – 1998-2000

	1998 Total (%)	1999 Total (%)	2000 Total (%)
Total number of isolates tested for INH, RMP, EMB, SM and PZA	8 (100.0)	9 (100.0)	10 (100.0)
Isolates susceptible	8 (100.0)	9 (100.0)	10 (100.0)

Table 10. Reported results for routine drug susceptibility testing of MTB isolates to first-line anti-tuberculosis drugs, Northwest Territories – 1998-2000

	1998 Total (%)	1999 Total (%)	2000 Total (%)
Total number of isolates tested for INH, RMP, EMB, SM and PZA	27 (100.0)	11 (100.0)	8 (100.0)
Isolates susceptible	27 (100.0)	11 (100.0)	8 (100.0)

Table 11. Reported results for routine drug susceptibility testing of MTB isolates to first-line anti-tuberculosis drugs, Nova Scotia – 1998-2000

	1998 Total (%)	1999 Total (%)	2000 Total (%)
Total number of isolates tested for INH, RMP, EMB and PZA*	9 (100.0)	8 (100.0)	4 (100.0)
Isolates susceptible	8 (88.9)	7 (87.5)	4 (100.0)
Isolates resistant to one or more drugs	1 (11.1)	1 (12.5)	- (0.0)
Mono-resistance	1 (11.1)	1 (12.5)	- (0.0)
INH	1 (11.1)	1 (12.5)	- (0.0)

* Routine testing for SM not conducted in Nova Scotia.

Table 12. Reported results for routine drug susceptibility testing of MTB isolates to first-line anti-tuberculosis drugs, Nunavut* – 1998-2000

	1998 Total (%)	1999 Total (%)	2000 Total (%)
Total number of isolates tested for INH, RMP, EMB, PZA and SM**	N/A	15 (100.0)	29 (100.0)
Isolates susceptible	N/A	15 (100.0)	28 (96.6)
Isolates resistant to one or more drugs	N/A	- (0.0)	1 (3.4)
Mono-resistance	N/A	- (0.0)	1 (3.4)
INH	N/A	- (0.0)	1 (3.4)

* Note: Nunavut began reporting in 1999.
 ** Routine testing for SM not conducted for Nunavut when tested by Quebec. (n=13 for 1999 and n=28 for 2000)

Table 13. Reported results for routine drug susceptibility testing of MTB isolates to first-line anti-tuberculosis drugs, Ontario – 1998-2000

	1998 Total (%)	1999 Total (%)	2000 Total (%)
Total number of isolates tested for INH, RMP, EMB, SM and PZA	629 (100.0)	587 (100.0)	599 (100.0)
Isolates susceptible	538 (85.5)	488 (83.1)	519 (86.6)
Isolates resistant to one or more drugs	91 (14.5)	99 (16.9)	80 (13.3)
Monoresistance	55 (8.7)	57 (9.7)	52 (8.7)
INH	34 (5.4)	34 (5.8)	23 (3.8)
SM	11 (1.7)	19 (3.2)	16 (2.7)
PZA**	6 (1.0)	4 (0.7)	12 (2.0)
EMB	4 (0.6)	- (0.0)	1 (0.2)
MDR-TB*	11 (1.7)	12 (2.0)	9 (1.5)
INH & RMP	2 (0.3)	2 (0.3)	1 (0.2)
INH, RMP & SM	1 (0.2)	3 (0.5)	3 (0.5)
INH, RMP & EMB	- (0.0)	1 (0.2)	2 (0.3)
INH, RMP & PZA	- (0.0)	1 (0.2)	- (0.0)
INH, RMP, EMB & SM	2 (0.3)	- (0.0)	2 (0.3)
INH, RMP, SM & PZA	- (0.0)	- (0.0)	1 (0.2)
INH, RMP, EMB, SM & PZA	6 (1.0)	5 (0.9)	- (0.0)
Other Patterns	25 (4.0)	30 (5.1)	19 (3.2)
INH & EMB	2 (0.3)	4 (0.7)	2 (0.3)
INH & SM	20 (3.2)	20 (3.4)	14 (2.3)
EMB & RMP	- (0.0)	- (0.0)	2 (0.3)
INH, SM & EMB	2 (0.3)	4 (0.7)	1 (0.2)
INH, SM & PZA	1 (0.2)	2 (0.3)	- (0.0)

* MDR-TB is defined as resistance to at least INH and RMP.
** Includes 1 *M. bovis* isolate for 1999 and 2 *M. bovis* isolates for 2000.

Table 14. Reported results for routine drug susceptibility testing of MTB isolates to first-line anti-tuberculosis drugs, Prince Edward Island – 1998-2000

	1998 Total (%)	1999 Total (%)	2000 Total (%)
Total number of isolates tested for INH, RMP, EMB, and PZA*	2 (100.0)	2 (100.0)	3 (100.0)
Isolates susceptible	2 (100.0)	2 (100.0)	3 (100.0)

* Routine testing for SM not conducted in Prince Edward Island.

Table 15. Reported results for routine drug susceptibility testing of MTB isolates to first-line anti-tuberculosis drugs, Québec – 1998-2000

	1998 Total (%)	1999 Total (%)	2000 Total (%)
Total number of isolates tested for INH, RMP, EMB and PZA**	264 (100.0)	268 (100.0)	278 (100.0)
Isolates susceptible	231 (87.5)	236 (88.1)	249 (89.6)
Isolates resistant to one or more drugs	33 (12.5)	32 (11.9)	29 (10.4)
Monoresistance	28 (10.6)	28 (10.4)	28 (10.1)
INH	9 (3.4)	17 (6.3)	19 (6.8)
RMP	- (0.0)	1 (0.4)	- (0.0)
SM	13 (4.9)	NT**	NT**
PZA***	6 (2.3)	10 (3.7)	9 (3.2)
MDR-TB*	2 (0.8)	2 (0.7)	1 (0.4)
INH & RMP	- (0.0)	1 (0.4)	- (0.0)
INH, RMP & SM	1 (0.4)	NT**	NT**
INH, RMP & EMB	1 (0.4)	- (0.0)	1 (0.4)
INH, RMP, EMB & PZA	- (0.0)	1 (0.4)	- (0.0)
Other Patterns	3 (1.1)	2 (0.7)	- (0.0)
INH & SM	2 (0.8)	NT**	NT**
INH & PZA	1 (0.4)	2 (0.7)	- (0.0)

* MDR-TB is defined as resistance to at least INH and RMP.
 ** Routine testing for SM not conducted in Québec effective January 1, 1999. (NT=not tested)
 *** Includes 1 *M. bovis* isolate for 1999 and 2 *M. bovis* isolates for 2000.

Table 16. Reported results for routine drug susceptibility testing of MTB isolates to first-line anti-tuberculosis drugs, Saskatchewan – 1998-2000

	1998 Total (%)	1999 Total (%)	2000 Total (%)
Total number of isolates tested for INH, RMP, EMB and SM*	49 (100.0)	40 (100.0)	64 (100.0)
Isolates susceptible	47 (95.9)	39 (97.5)	58 (90.6)
Isolates resistant to one or more drugs	2 (4.1)	1 (2.5)	6 (9.4)
Monoresistance	1 (2.0)	- (0.0)	4 (6.3)
INH	1 (2.0)	- (0.0)	2 (3.1)
SM	- (0.0)	- (0.0)	1 (1.6)
EMB	- (0.0)	- (0.0)	1 (1.6)
Other patterns	1 (2.0)	1 (2.5)	2 (3.1)
INH & SM	1 (2.0)	1 (2.5)	1 (1.6)
INH & EMB	- (0.0)	- (0.0)	1 (1.6)

* Routine testing for PZA not conducted in Saskatchewan.

Table 17. Reported results for routine drug susceptibility testing of MTB isolates to first-line anti-tuberculosis drugs, Yukon Territory – 1998-2000

	1998 Total (%)	1999 Total (%)	2000 Total (%)
Total number of isolates tested for INH, RMP, EMB and SM*	1 (100.0)	- (0.0)	3 (100.0)
Isolates susceptible	1 (100.0)	- (0.0)	3 (100.0)

* Routine testing for PZA not conducted in Yukon Territory.
 • No isolates were tested for Yukon Territory in 1999.

▶ LIMITATIONS

Susceptibility testing for first-line anti-tuberculosis drugs is not uniform across the country. Therefore, there are limitations in interpreting the data, particularly the percentage of isolates that are resistant to SM and PZA.

More epidemiologic information on the TB cases from which the isolates were submitted would be desirable to critically examine drug resistance patterns in Canada. Demographic information is sparse; only gender and year of birth are routinely reported to this surveillance system. As well, no differentiation can be made between primary and secondary/acquired drug resistance from the data in the system. The participating laboratories attempted to collect data on the country of origin in 1999. However, because of the difficulties in doing so, this variable has subsequently been dropped from the surveillance data set.

▶ CONCLUSIONS

With growing worldwide concern regarding TB drug resistance, this laboratory-based surveillance system is vital in providing the necessary data in a timely fashion to monitor trends in TB drug resistance in Canada. The surveillance data collected to date indicate that the prevalence of TB drug resistance in this country is similar to that in the overall global situation. However, with data collected only for three years, any discussion of “trends” is premature at the present time. Several more years of collected data will be necessary in order to examine the unfolding pattern of TB drug resistance in Canada.

▶ REFERENCE

1. The WHO/IUATLD Global Project on Anti-tuberculosis Drug Resistance Surveillance. Anti-tuberculosis drug resistance in the world; Report No. 2. (WHO/CDS/TB/2000.278). Geneva: World Health Organization, 2000.

► Appendix 1

Participating Laboratories of the Canadian Tuberculosis Laboratory Surveillance System (CTBLSS)

Alberta (Alberta, Northwest Territories and Nunavut)

Lorraine Ingham
Mycobacteriology Technical Supervisor
Provincial Laboratory of Public Health
Calgary
Telephone: (403) 670-1209
Fax: (403) 270-2216
email: Lorraine.Ingham@CRHA-Health.Ab.ca

Dr. Jutta Preiksaitis
Director
Provincial Laboratory of Public Health
for Northern Alberta
Edmonton
Telephone: (780) 407-8903
Fax: (780) 407-8984
email: jkp@bugs.uah.ualberta.ca

British Columbia (British Columbia and Yukon Territory)

Dr. Mabel Rodrigues, PhD
Section Supervisor TB/Mycology
BCCDC Laboratory Services
Vancouver
Telephone: (604) 775-2153
Fax: (604) 660-6073
email: mabel.rodrigues@bccdc.hnet.bc.ca

Dr. W.A. Black
Medical Microbiologist
BCCDC Laboratory Services
Professor, Medical Microbiology, UBC
Vancouver
Telephone: (604) 660-6029
Fax: (604) 660-6073
email: william.black@bccdc.hnet.bc.ca

Dr. Judy Isaac-Renton
Director
BCCDC Laboratory Services
Professor, Medical Microbiology, UBC
Vancouver
Telephone: (604) 660-6032
Fax: (604) 660-6073

Manitoba

Joyce Wolfe
Mycobacteriology Supervisor
Health Sciences Centre
Winnipeg
Telephone: (204) 787-7652
Voice Mail: (204) 787-1760 (7652#)
Fax: (204) 787-4699
email: joyce_wolfe@hc-sc.gc.ca

**New Brunswick
(see also Quebec)**

Joan MacDonald
Supervisor, Microbiology Laboratory
Department of Laboratory Medicine
Saint John Regional Hospital
Saint-John
Telephone: (506) 648-7226
Fax: (506) 648-6537
email: macjoan@reg2.health.nb.ca

Newfoundland

Sandra B. March
Clinical Microbiologist
Newfoundland Public Health Laboratory
St. John's
Telephone: (709) 777-6535
Fax: (709) 777-6611
email: smarch@nf.aibn.com

**Northwest Territories
(see also Alberta and Quebec)**

Norine Fraley, MLT
Supervisor Bacteriology
Stanton Regional Hospital
Yellowknife
Telephone: (867) 669-4162 or 669-4166
Fax: (867) 669-4141
email: norine_fraley@gov.nt.ca

**Nova Scotia
(Nova Scotia and Prince Edward Island)**

Dr. David J.M. Haldane
Director
Bacteriology and Special Pathogens
Queen Elizabeth II Health Sciences Centre
Halifax
Telephone: (902) 473-2392
Fax: (902) 473-4432
email: plmdjh@qe2-hsc.ns.ca

Ontario

Dr. Frances Jamieson
Clinical Microbiologist
Central Public Health Laboratory
Toronto
Telephone: (416) 235-5841
Fax: (416) 235-5951

Pam Chedore
Head
TB and Mycobacteriology Laboratory
Central Public Health Laboratory
Toronto
Telephone: (416) 235-5928
Fax: (416) 235-6013
email: pam.chedore@moh.gov.on.ca

Job Babu
Mycobacteriology
Hamilton Regional Laboratory Medicine Network
McMaster University Medical Centre Site
Department of Laboratory Medicine
Hamilton
Telephone: (905) 521-2100 (76311)
Fax: (905) 577-0198

Prince Edward Island (see also Nova Scotia)

L.P. Abbot
Clinical Head Microbiology
Queen Elizabeth Hospital
Charlottetown
Telephone: (902) 894-2309
Fax: (902) 894-2385
email: labbott@isn.net

Québec (Quebec, New Brunswick, Northwest Territories and Nunavut)

Louise Thibert, MSc
Head
Mycobacteriology
Laboratoire de santé publique du Québec – INSPQ
Sainte-Anne-de-Bellevue
Telephone: (514) 457-2070 (237)
Fax: (514) 457-6346
email: lthibert@lspq.org

Saskatchewan

North Dorothy Cheke, RT
Senior Technologist / TB Laboratory
Clinical Microbiology, Royal University Hospital
Saskatoon
Telephone: (306) 655-1769
Fax: (306) 655-1726

M. Kanchana
Director, TB Laboratory
Clinical Microbiology, Royal University Hospital
Saskatoon
Telephone: (306) 655-1762
Fax: (306) 655-1726
email: manickem@duke.usask.ca

South Evelyn Nagle, RT
Section Head, Bacteriology/Mycobacteriology
Saskatchewan Health, Provincial Laboratory
Regina
Telephone: (306) 787-8634
Fax: (306) 787-9122
email: enagle@health.gov.sk.ca

Dr. P. Pieroni
Microbiologist
Saskatchewan Health, Provincial Laboratory
Regina
Telephone: (306) 787-3195
Fax: (306) 787-1525
email: ppieroni@health.gov.sk.ca

Yukon Territory
(see British Columbia)

Federal

Dr. Howard Njoo
Director
Tuberculosis Prevention and Control
Centre for Infectious Disease Prevention and
Control
Population and Public Health Branch
Ottawa
Telephone: (613) 941-1191
Fax: (613) 946-3902
email: howard_njoo@hc-sc.gc.ca

Penny Nault
TB Database Manager
Tuberculosis Prevention and Control
Centre for Infectious Disease Prevention and
Control
Population and Public Health Branch
Ottawa
Telephone: (613) 941-6121
Fax: (613) 946-3902
email: penny_nault@hc-sc.gc.ca

Melissa Phypers, MSc
Senior Epidemiologist
Tuberculosis Prevention and Control
Centre for Infectious Disease Prevention and
Control
Population and Public Health Branch
Ottawa
Telephone: (613) 946-3920
Fax: (613) 946-3902
email: melissa_phypers@hc-sc.gc.ca

Dr. Amin Kabani
National Reference Centre for Mycobacteriology
Federal Laboratories for Health Canada
Telephone: (204) 787-1928
Fax: (204) 787-4699
email: amin_kabani@hc-sc.gc.ca

Appendix 2



Health Canada Santé Canada

Serial No. - N° de série

The Canadian Tuberculosis Laboratory Surveillance System
M. TUBERCULOSIS COMPLEX ANTIMICROBIAL
SUSCEPTIBILITY REPORTING FORM

Système de surveillance des laboratoires de tuberculose au Canada
RAPPORT SUR LA SENSIBILITÉ DES SOUCHES DU COMPLEXE
M. TUBERCULOSIS AUX ANTIMICROBIENS

FOR INTERNAL USE ONLY - POUR USAGE INTERNE SEULEMENT		Unique Source Laboratory ID No. - Identificateur unique du laboratoire déclarant:			
Date Rec'd at TBPC: Date de réception au LATB: Y / A M D / J		Date specimen / culture received at laboratory: Date de réception échantillon / culture au laboratoire: Y / A M D / J			
TBPC Number: Numéro du LATB:					
Specie: Espèce: <input type="checkbox"/> M. tuberculosis (may include M. africanum or M. microti) (peut inclure M. africanum et M. microti) <input type="checkbox"/> M. bovis <input type="checkbox"/> M. BCG bovis <input type="checkbox"/> MTB Complex (species unknown) (Complexe MTB (espèce inconnu))					
Have susceptibility test results been previously reported for this patient? - Des résultats d'antibiogramme ont-ils déjà été fournis pour ce patient? <input type="checkbox"/> No / Non <input type="checkbox"/> Yes / Oui → What is the previous Unique Source Laboratory ID No.? / Identificateur antérieur? _____ → What is the previous Form No.? (if known) / N° de formulaire antérieur? (Si connu) _____					
Note: Only DRUG TESTING RESULTS OF ONE ISOLATE are to be reported. No subsequent drug testing results for the same patient are to be reported unless the sensitivity pattern changes.		Note: Ne fournir que les RÉSULTATS POUR UNE SEULE SOUCHE par patient à moins d'un changement du profil de sensibilité.			
1	Province / territory from which this report originates: Province / territoire qui soumet ce rapport:	<input type="text"/>	(see code list) (voir liste de codes)	PROV / TERR CODES PROV / TERR 10 = NFLD / TN 46 = MAN 11 = PEI / IPÉ 47 = SASK 12 = NS / NÉ 48 = ALTA / ALB 13 = NB 59 = BC / BC 24 = QUÉ / Qc 60 = YUK 35 = ONT 61 = NWT / TNO 62 = NUN	
2	Province / territory from which specimen originated: Province / territoire d'où provient l'échantillon:	<input type="text"/>	(see code list) (voir liste de codes)		
3	Patient's date of birth: Date de naissance du patient:	Y / A M D / J	(CCYY/MM/DD) (SSAA/MM/JJ) <input type="checkbox"/> Unknown / Inconnu		
4	Patient's gender: Sexe du patient:	<input type="checkbox"/> Male / Masculin <input type="checkbox"/> Female / Féminin <input type="checkbox"/> Unknown / Inconnu			
5	LABORATORY RESULTS RÉSULTATS DE LABORATOIRE	Concentration (if different from on file) Concentration (si autre que spécifiée)	Results (check appropriate box for every drug) Résultats (cocher la case pertinente pour chaque antibiotique)		
	Antituberculous Drugs Agents Antituberculeux		Sensitive / Sensible	Resistant / Résistant	Other (specify) / Autre (préciser)
	SM (Streptomycin) / (Streptomycine)	mg / L	<input type="checkbox"/>	<input type="checkbox"/>	
	INH (Isoniazid) / (isoniazide)	mg / L	<input type="checkbox"/>	<input type="checkbox"/>	
	RMP (Rifampin) / (Rifampicine)	mg / L	<input type="checkbox"/>	<input type="checkbox"/>	
	EMB (Ethambutol)	mg / L	<input type="checkbox"/>	<input type="checkbox"/>	
	PZA (Pyrazinamide)	mg / L	<input type="checkbox"/>	<input type="checkbox"/>	
	2nd line drugs (specify) Antibiotiques de 2° ligne (préciser)	Concentration	Sensitive / Sensible	Resistant / Résistant	Other (specify) / Autre (préciser)
	1.	mg / L	<input type="checkbox"/>	<input type="checkbox"/>	
	2.	mg / L	<input type="checkbox"/>	<input type="checkbox"/>	
	3.	mg / L	<input type="checkbox"/>	<input type="checkbox"/>	
	4.	mg / L	<input type="checkbox"/>	<input type="checkbox"/>	
	5.	mg / L	<input type="checkbox"/>	<input type="checkbox"/>	
	6.	mg / L	<input type="checkbox"/>	<input type="checkbox"/>	
6	Comments - Commentaires				

HC/SC 9061
(07-2000)

Copy 1 (White) - Reporting Laboratory
Copie 1 (Blanche) - Laboratoire déclarant

Copy 2 (Yellow) - Tuberculosis Prevention and Control (TBPC)
Copie 2 (Jaune) - Lutte anti-tuberculeuse (LATB)