

<b>Fish Health Events ( 2004 - First Quarter )<sup>1</sup></b>				
<b>DFO SubZone</b>	<b>Species</b>	<b>Water Source</b>	<b>Veterinary Diagnosis</b>	<b>Count<sup>2</sup></b>
<b>All Zones</b>	<b>Atlantic salmon</b>	<b>Freshwater</b>	<b>Aeromonas salmonicida Infection</b>	<b>8</b>
<b>2-3</b>	<b>Atlantic salmon</b>	<b>Salt/brackish</b>	<b>Aeromonas salmonicida Infection</b>	<b>2</b>
			<b>Lepeophtheirus Infection</b>	<b>2</b>
			<b>Myxobacterial Infection</b>	<b>7</b>
<b>2-4</b>	<b>Atlantic salmon</b>	<b>Salt/brackish</b>	<b>Aeromonas salmonicida Infection</b>	<b>2</b>
			<b>Lepeophtheirus Infection</b>	<b>6</b>
			<b>Myxobacterial Infection</b>	<b>5</b>
<b>3-1 + 3-2</b>	<b>Atlantic salmon</b>	<b>Salt/brackish</b>	<b>Myxobacterial Infection</b>	<b>2</b>
<b>3-3</b>	<b>Atlantic salmon</b>	<b>Salt/brackish</b>	<b>Lepeophtheirus Infection</b>	<b>4</b>
			<b>Renibacterium salmoninarum Infection</b>	<b>2</b>
<b>3-4 + 3-5</b>	<b>Atlantic salmon</b>	<b>Salt/brackish</b>	<b>Aeromonas salmonicida Infection</b>	<b>1</b>
			<b>Myxobacterial Infection</b>	<b>2</b>
			<b>Yersinia ruckeri Infection</b>	<b>1</b>
<b>All Zones</b>	<b>Pacific salmon</b>	<b>Freshwater</b>	<b>Aeromonas salmonicida Infection</b>	<b>1</b>
			<b>Myxobacterial Infection</b>	<b>2</b>
			<b>Renibacterium salmoninarum Infection</b>	<b>1</b>
<b>All Zones</b>	<b>Pacific salmon</b>	<b>Salt/brackish</b>	<b>Nucleospora (Enterocytozoan) salmonis Infection</b>	<b>2</b>
			<b>Renibacterium salmoninarum Infection</b>	<b>10</b>

**Notes**

- 1 Denotes new Fish Health Events for the quarter
- 2 Counts of veterinary diagnosis are based on FISH GROUP (not site); more than one fish group may exist at a site
- 3 See notes 2 and 3 of Average Mortality Rate report