

# Nitrogen Fertilization

Nitrogen fertilizers are most effectively used as part of a balanced fertilization plan that aims to maximize economic return and maintain environmental quality. Within the Canadian prairies, nitrogen (N) is most often the yield-limiting nutrient with respect to crop production. Nitrogen contributes primarily to grain yield and forage biomass production, and at the same time to protein. Once yield requirements for nitrogen have been met, further nitrogen uptake contributes to protein level increases. Optimum economic response to applied nitrogen fertilizer requires that the nitrogen application rate is balanced with moisture and the availability of other plant nutrients. Soil testing to determine availability of nitrogen along with other nutrients and the assessment of stored soil moisture is therefore critical to selecting the appropriate rate of nitrogen fertilizer to match crop yield potential.


In the event of a crop failure, much of the nitrogen left unused in the soil, under Saskatchewan weather conditions, can carry into the following season and can be accounted for by a soil test.

## Yield

For dryland annual crops grown on the prairies, much of the nitrogen uptake that contributes to yield occurs in the first few weeks following germination, and is complete after about two months (Figure 1).

Maximum nutrient uptake occurs earlier than maximum biomass production. After maximum biomass is achieved, nitrogen uptake contributes mainly to protein content.

The response of grain yield to nitrogen fertilization depends greatly on moisture. Improved moisture conditions usually translate to more yield, to a point where other limiting factors come into play. Excess moisture can reduce yield due to leaching losses of nitrate, as well as loss of nitrates by conversion (denitrification) to gases that escape from the soil. High levels of available soil nitrogen early on in the growing season can promote heavy vegetative growth and water use.



**Figure 1.** Changes in nitrogen uptake of cereals with days after emergence in the field experiment at Melfort (Malhi et al., 2004).

## Protein

Additional nitrogen usually results in higher grain protein content. Under conditions of extreme nitrogen deficiency, a small amount of added nitrogen may result in no protein increase, or even a small decrease, due to dilution from the large growth enhancement associated with the added nitrogen. Beyond this point, increased soil nitrogen supplies are normally associated with increased protein content. Timing of moisture is important and in particular the extent and duration of any moisture stress. Moisture stress at flowering can lower yield but increase protein content.

## Nitrogen Fertilizer Sources

Forms of nitrogen fertilizer widely available and used in Saskatchewan include granular urea (46-0-0), anhydrous ammonia (82-0-0) and urea ammonium nitrate (UAN) solutions (28-0-0). See *Table 1 on page 3 of this document*.

Granular ammonium nitrate (34-0-0) may be available but is becoming more difficult to access. Other fertilizer products that serve as sources of nitrogen, as well as other nutrients, include ammonium sulphate (21-0-0-24), mono-ammonium phosphates such as 12-51-0, ammonium phosphate sulphate (16-20-0-14), ammonium polyphosphate (10-34-0), and ammonium thiosulphate (15-0-0-20). Animal manures and composts are also valuable sources of nitrogen, along with other nutrients. A low and often variable content of available nitrogen in animal manures requires special considerations in their use (see *Nutrient Values of Manure and Managing Manure as a Fertilizer* fact sheets).

Fertilizers that contain or produce ammonium result in acidity when the ammonium is converted to nitrate in the soil. In most Saskatchewan soils, this is not an issue; however, it may be of concern in some sandy, acidic soils.

## Application

Nitrogen fertilizers can be permanently lost through ammonia volatilization, denitrification, leaching and run-off. Nitrogen may also be temporarily tied up in the soil organic matter by microorganisms (immobilization), especially when the fertilizer is closely associated with crop residues with low nitrogen content. A rule of thumb is that potential losses of nitrogen fertilizer will be reduced the closer the fertilizer is applied to the time of crop use. Late fall or spring pre-plant banding of anhydrous ammonia or urea can be efficient, depending on soil and climatic conditions. However, applications made as a band at the time of seeding often are most efficient in increasing yield. In general, the application of nitrogen fertilizer in-soil, especially in a band, reduces potential losses compared to surface application. Urea containing fertilizers, including manures, are susceptible to ammonia volatilization losses when surface applied and not incorporated. There has been increasing use of surface application

of solution 28-0-0 nitrogen fertilizer in a dribble-band. Surface placement of nitrogen fertilizer, whether in a band or broadcast, will work best if there is a rain following application so as to move the fertilizer into the root zone. If available, granular ammonium nitrate (34-0-0) is suitable for surface broadcast application to crops like winter wheat or forage grasses, as volatilization losses are lower compared to surface broadcast urea.

When surface application is the only placement option, urea-containing fertilizer (46-0-0 or 28-0-0) may be treated with a urease inhibitor prior to use in order to reduce ammonia volatilization losses. The urease inhibitor (*AGROTAIN*) will prevent the release of ammonia gas from the urea for a period of about two weeks (see product label for application rates and times of inhibition), giving an opportunity for the urea to be moved into the soil by rainfall over this period.

Application of all nitrogen fertilizer at the time of seeding in a single pass (direct seeding) has become a common practice in Saskatchewan. Placing nitrogen fertilizers in the seed-row is a good method, provided the rates are kept low. There is a limit to how much fertilizer nitrogen can be safely placed in the seed-row before injury in the form of reduced germination and emergence occurs. Such injury is due to the salt effect of the fertilizer holding back moisture from the seed and seedling, as well as ammonia gas produced from urea fertilizer. The maximum safe rates of seed-placed nitrogen will vary with the opener spread and row spacing of the seeder, the crop, the soil texture and moisture (the fact sheet *Guidelines for Safe Rates of Fertilizer Applied With the Seed* should be followed to avoid damage.)

To reduce or eliminate risk of crop damage when applying high rates of nitrogen fertilizers at the time of seeding, separation of the seed from the fertilizer is needed. This may be accomplished by placing the fertilizer in a separate furrow from the seed. Various equipment configurations are available to do this. Common separation strategies are placement of the fertilizer band a minimum of one inch below and one inch to the side of the seed-row (side-banding) and placement of the fertilizer band midway between every second seed-row (mid-row banding) (Figure 2).

**Table 1. Common Commercial Nitrogen Fertilizer Forms in Saskatchewan**

Form of nitrogen (N) fertilizer	Common formulations and usage in Saskatchewan	Reaction in soil	Physical properties of the fertilizer	Comments
<b>Dry granular</b>	46-0-0 urea (NH <sub>2</sub> ) <sub>2</sub> CO - Most commonly used source of N.	- Salt index* is 1.62. - Urea converts to ammonia in presence of urease enzyme. - Ammonia then converts to ammonium in presence of water (during this reaction the soil pH around the granule site is raised temporarily). - Ammonium is held on the soil with small amounts in the soil water. - Plants use both ammonium-N and nitrate-N.	- Granular. - Soluble in water. - Bulk density, non-compacted about 47 lb./ft. <sup>3</sup> . - Melting point is 133°C. - Critical relative humidity (RH) at which the fertilizer starts absorbing moisture is 73%.	- See the Farm Facts <i>Guidelines For Safe Rates Of Fertilizer Applied With the Seed</i> for banding with the seed. - Addition of a urease inhibitor can slow the conversion of urea to ammonia for up to two weeks (see product label for AGROTAIN rates.) - Avoid storage or contact with ammonium nitrate because the mixture will draw moisture (hygroscopic) and turn into slush. Critical RH of the mixture is 18%. - Most common method of application is banding into soil (the furrow must be closed to prevent ammonia escape).
	21-0-0-24; 20-0-0-24; 19-2-0-22 Ammonium sulphate (NH <sub>4</sub> ) <sub>2</sub> SO <sub>4</sub> - Most commonly used plant available form of S.	- Salt index is 3.25. - Sulphate remains in the soil water and is mobile.	- Granular or crystal. - Critical RH is 79%.	- Fertilizer sulphur is expressed as S. - Sulphate-S (SO <sub>4</sub> <sup>2-</sup> ) is the plant available form of S.
<b>Gas (liquid when compressed)</b>	82-0-0 - anhydrous ammonia - NH <sub>3</sub> - Second most commonly used source of N.	- Salt index is 0.57. - Ammonia gas reacts with soil water to form ammonium. - Ammonium is held on the soil with some in the soil water.	- Compressed gas. - Sharp pungent odor. - Soluble in water. - Is a liquid at -33°C at one atmosphere of pressure.	- Free ammonia in soil is very toxic and can move through cell walls, injuring plant tissue. - Must be banded into moist soil and the furrow must close to prevent ammonia loss. - For side banding, if the seedbed is fractured because of poor seedbed conditions, ammonia can move into the seed row, injuring seed or seedlings. - It is important to understand and comply with all required safety and labeling guidelines.
<b>Liquid (solution-N fertilizer)</b>	28-0-0 - 50% urea-N, 25% ammonium-N, 25% nitrate-N. - Increasing use	- Salt index is 2.3. - The urea fraction is subject to volatile losses.**	- Has a specific gravity of 1.3 at 15.6°C. - One imperial gallon weighs 13 lb. - Each imperial gallon contains 3.57 lb. of N. - pH is 7.0 to 7.6. - Salt out temperature is -18°C. - Must be above 0°C to be pumped. - Can be blended with other liquid fertilizers (P, K, S) in the spring for immediate application.	- Liquid fertilizers used in the seed row must follow the same guidelines as for granular fertilizer. See Farm Facts <i>Guidelines For Safe Rates Of Fertilizer Applied With Seed</i> . - Blended liquid fertilizers can provide all the nutrient needs for a crop, or liquid fertilizer system can be used in combination with granular or anhydrous ammonia fertilizer systems. - Can also be applied in irrigation water (fertigation). - Can be banded into the soil, spoke injected, dribble banded on the surface, foliar spray (keep N rates low for foliar spray application, water volumes high, apply into evening and/or under cool weather conditions to avoid leaf scorch). - Addition of a urease inhibitor (AGROTAIN) can slow the conversion of the urea portion of the fertilizer to ammonia for up to two weeks (see product label for AGROTAIN rates).

\* Salt index is a measure of the salt effect that the fertilizer induces in the soil solution due to the affinity for water. The higher the salt index number, the greater the salt effect. Based on 20 lb. of actual nutrient N, P<sub>2</sub>O<sub>5</sub>, K<sub>2</sub>O, S.


\*\* Volatile losses refer to volatilization of ammonia.

Recent research in Saskatchewan has shown relatively little difference in agronomic performance of nitrogen fertilizer applied either as anhydrous ammonia or urea at seeding time in mid-row or side band configurations (PAMI research update 761). Unlike phosphorus, nitrogen is considered a “mobile” nutrient in the soil and can move long distances with water moving to the roots. As a result, placement close to the seed is generally not considered as essential for early crop access to nitrogen. However, under conditions of severe deficiency and when nutrient mobility in soil is limited, some nitrogen placed close to or in the seed row can be beneficial to ensure early nitrogen nutrition. Granular fertilizers can be applied in the seed-row or banded in a separate furrow at the time of seeding, but anhydrous ammonia should never be placed where the ammonia can come into contact with the germinating seed or seedling. Liquid nitrogen fertilizers like 28-0-0 may be placed in the soil in furrows separate from the seed in a manner similar to other forms, and similar

restrictions apply as to safe rates of nitrogen in close proximity to the seed. However, with the liquid forms, separation may be more easily achieved by “squirting” (directed flow liquid) the fertilizer away from the seed.

Applications of nitrogen fertilizer may also be made after crop emergence. Such applications may be made as a “rescue treatment” when nitrogen deficiency in the crop is detected by visual observation and/or tissue testing. For most annual crops, early application, before stem extension, will result in the best yield, while delayed application will mainly increase protein in the grain. Top dressing is not generally recommended as the primary method of nitrogen application. However, it can be used when sufficient nitrogen was not applied before or at seeding, when additional nitrogen is warranted due to improved moisture conditions, or when applying nitrogen to increase grain protein content.

*This bulletin was revised by the Saskatchewan Soil Fertility Subcouncil, March 2005.*



**Figure 2.** Common In-Soil Placement Options for Fertilizers Applied at the Time of Seeding.