## **Errata: CAS Highlights**

August 2007

The CAS included a brief six-item screener to measure problematic drug use. The screener, the ASSIST, was developed by the World Health Organization.

An error was recently found in one symptom (represented by two items: CNAS5 and ASSIS5) of the ASSIST scale. Properly stated, the question is "Have you ever tried [AND FAILED] to control, cut down or stop using cannabis, marijuana or hashish {or other drugs}. The phrase "and failed" was not asked of respondents. This error, in turn, affects the following derived variables: ASISTCAN, ASISTCN3, ASISTCN2, ASISTIL and ASISTIL3.

NOTE: A small study to assess the impact of the missing "and failed" phrase suggested that estimates with the missing phrase underestimate the standard ASSIST item, but total scores and their cutoffs do not differ significantly between the two versions. This error should not affect estimates of subgroup differences (i.e., sex, age and regional differences). Caution and warning should be used in making direct comparisons with other studies using the ASSIST.

Page	For	Should read
P.6, column 2, line 22	"The most common concern is failing to control use (4.8%)	"The most common concern is attempts to cut down (4.8%)
	followed by"	followed by"
P.8, Figure 4	"Failed control"	"Attempts to cut down "

This document was published by the Canadian Centre on Substance Abuse (CCSA).

For additional copies, contact CCSA, 75 Albert St., Suite 300 Ottawa ON K1P 5E7 tel.: 613 235-4048 e-mail: info@ccsa.ca

This document can also be downloaded as a PDF at www.ccsa.ca

Ce document est également disponible en français sous le titre Enquête sur les toxicomanies au Canada (ETC) : Une enquête nationale sur la consommation d'alcool et d'autres drogues par les Canadiens : La prévalence de l'usage et les méfaits : Points saillants

ISBN 1-896323-35-9