

2007 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 1

Canada Western Red Spring

Percentage of seeded acres / () = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
Lillian	14.8 (3.6)	0.0 (T)	19.2 (3.9)	17.9 (5.4)	1.4 (0.0)
AC Barrie	13.8 (17.9)	34.2 (40.7)	12.5 (16.6)	3.5 (6.1)	0.0 (0.0)
Superb	12.8 (18.3)	10.3 (13.3)	13.1 (17.6)	14.0 (22.3)	2.9 (0.0)
Harvest	10.3 (5.5)	6.7 (4.0)	2.2 (1.7)	23.2 (11.8)	0.0 (0.0)
McKenzie	7.1 (9.3)	3.7 (4.0)	12.6 (15.4)	1.7 (4.2)	24.5 (11.7)
Prodigy	4.2 (4.7)	0.5 (0.1)	6.0 (6.6)	4.2 (5.0)	0.0 (2.3)
AC Eatonia	4.0 (4.5)	0.0 (T)	2.6 (2.4)	8.2 (10.2)	8.6 (0.0)
AC Domain	3.8 (3.9)	18.1 (18.4)	0.4 (0.4)	0.0 (T)	0.0 (0.0)
AC Intrepid	3.5 (5.4)	2.6 (2.8)	3.9 (5.1)	3.5 (7.3)	24.6 (41.4)
AC Splendor	3.1 (3.7)	0.8 (1.0)	3.0 (3.1)	4.6 (6.0)	13.4 (21.9)
Other	3.1 (3.4)	0.9 (2.2)	3.9 (3.8)	3.3 (3.6)	18.1 (2.2)
CDC Teal	3.0 (3.9)	3.0 (2.2)	2.0 (2.9)	4.3 (6.1)	0.0 (20.5)
CDC Imagine	2.8 (3.2)	1.8 (1.4)	3.8 (4.4)	2.2 (2.8)	0.0 (0.0)
AC Elsa	2.4 (2.3)	0.5 (0.6)	4.2 (3.9)	1.1 (1.1)	0.0 (0.0)
5602 HR	2.3 (0.8)	7.9 (2.0)	0.3 (0.2)	1.5 (0.9)	0.0 (0.0)
AC Cadillac	2.2 (3.0)	1.1 (1.2)	3.5 (5.1)	1.0 (1.2)	0.0 (0.0)
Infinity	1.5 (0.2)	0.7 (0.2)	2.2 (0.2)	1.1 (0.1)	3.6 (0.0)
Columbus	1.2 (1.4)	T (T)	2.1 (2.6)	0.7 (0.6)	0.0 (0.0)
CDC Bounty	0.7 (1.1)	1.1 (1.4)	0.7 (1.2)	0.4 (0.9)	0.0 (0.0)
AC Abbey	0.5 (1.7)	0.0 (0.0)	0.8 (2.1)	0.6 (2.3)	0.0 (0.0)
CDC Go	0.5 (0.1)	0.6 (0.0)	0.1 (0.1)	1.1 (0.2)	0.0 (0.0)
Journey	0.5 (0.4)	T (0.3)	0.1 (0.1)	1.3 (0.9)	0.0 (0.0)
5601 HR	0.5 (0.7)	2.3 (3.0)	0.0 (0.1)	T (T)	0.0 (0.0)
5400 IP	0.4 (0.0)	0.9 (0.0)	0.5 (0.0)	0.1 (0.0)	2.9 (0.0)
Lovitt	0.3 (0.3)	0.8 (0.7)	0.1 (0.2)	0.3 (0.2)	0.0 (0.0)
AC Cora	0.2 (0.0)	0.8 (0.0)	T (0.0)	0.0 (0.0)	0.0 (0.0)
5600 HR	0.1 (0.2)	0.3 (0.1)	0.2 (0.3)	T (0.2)	0.0 (0.0)
Somerset	0.1 (T)	0.2 (0.1)	T (T)	T (0.0)	0.0 (0.0)
5500 HR	T (0.2)	0.1 (T)	T (0.1)	0.0 (0.4)	0.0 (0.0)
CDC Alsask	T (T)	0.0 (0.0)	T (T)	T (0.0)	0.0 (0.0)
CDC Osler	T (0.1)	0.0 (0.0)	T (0.1)	0.0 (T)	0.0 (0.0)
AC Michael	T (0.0)	0.0 (0.0)	0.0 (0.0)	T (0.0)	0.0 (0.0)
Peace	T (T)	0.0 (0.0)	T (0.0)	0.0 (T)	0.0 (0.0)
Alsen	T (0.1)	0.1 (0.1)	0.0 (0.1)	0.0 (0.0)	0.0 (0.0)
AC Minto	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)

* Trace: less than 0.05 percent

2007 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 2

Canada Western Amber Durum

Percentage of seeded acres / () = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
Strongfield	41.5 (18.5)	0.0 (0.0)	42.7 (17.2)	36.4 (24.7)	100.0 (0.0)
AC Avonlea	22.7 (39.9)	0.0 (19.7)	20.2 (38.9)	33.3 (44.4)	0.0 (0.0)
Kyle	19.2 (22.6)	0.0 (50.8)	21.1 (24.3)	11.0 (15.1)	0.0 (0.0)
AC Navigator	11.1 (11.1)	0.0 (0.0)	11.0 (11.9)	11.5 (7.6)	0.0 (0.0)
Other	1.9 (1.1)	0.0 (0.0)	1.7 (1.1)	3.0 (1.3)	0.0 (0.0)
AC Morse	1.5 (2.9)	0.0 (0.0)	0.8 (2.1)	4.7 (6.7)	0.0 (0.0)
Commander	1.2 (1.7)	0.0 (0.0)	1.5 (2.0)	0.0 (0.2)	0.0 (0.0)
Napoleon	0.9 (2.2)	0.0 (29.5)	1.0 (2.6)	0.2 (0.0)	0.0 (0.0)

* Trace: less than 0.05 percent

2007 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 3

Canada Western Hard White Spring

Percentage of seeded acres / () = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
Snowbird	99.5 (96.6)	98.6 (98.5)	100.0 (96.8)	100.0 (92.8)	0.0 (0.0)
Kanata	0.5 (3.4)	1.4 (1.5)	0.0 (3.2)	0.0 (7.2)	0.0 (0.0)

* Trace: less than 0.05 percent

2007 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 4

Canada Western Extra Strong

Percentage of seeded acres / () = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
CDC Rama	37.5 (16.9)	0.0 (0.0)	48.3 (22.9)	0.0 (0.0)	0.0 (0.0)
AC Corinne	25.5 (6.5)	0.0 (0.0)	32.8 (8.9)	0.0 (0.0)	0.0 (0.0)
Burnside	17.8 (0.0)	41.6 (0.0)	17.7 (0.0)	0.0 (0.0)	0.0 (0.0)
Glenavon	12.0 (0.0)	0.0 (0.0)	0.0 (0.0)	94.1 (0.0)	0.0 (0.0)
Laser	5.6 (14.5)	58.4 (0.0)	0.0 (0.0)	0.0 (55.9)	0.0 (0.0)
Glenlea	1.7 (43.3)	0.0 (100.0)	1.2 (58.2)	5.9 (0.0)	0.0 (0.0)
Bluesky	0.0 (12.0)	0.0 (0.0)	0.0 (0.8)	0.0 (44.1)	0.0 (0.0)
Amazon	0.0 (6.8)	0.0 (0.0)	0.0 (9.3)	0.0 (0.0)	0.0 (0.0)
Walrus	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)
Wildcat	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)

* Trace: less than 0.05 percent

2007 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 5

Canada Prairie Spring White

Percentage of seeded acres / () = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
AC Vista	94.0 (89.1)	100.0 (100.0)	93.5 (88.1)	75.9 (0.0)	0.0 (100.0)
AC Karma	6.0 (10.9)	0.0 (0.0)	6.5 (11.9)	24.1 (0.0)	0.0 (0.0)
Other	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)

* Trace: less than 0.05 percent

2007 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 6

Canada Prairie Spring Red

Percentage of seeded acres / () = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
5700PR	39.0 (38.8)	39.9 (63.7)	32.8 (36.1)	42.8 (39.1)	0.0 (100.0)
AC Crystal	30.3 (30.7)	2.5 (0.0)	41.1 (42.2)	25.1 (25.5)	0.0 (0.0)
5701PR	14.9 (12.6)	49.9 (7.9)	21.5 (19.2)	8.6 (9.1)	0.0 (0.0)
AC Foremost	14.0 (16.0)	0.0 (0.0)	1.9 (1.2)	22.5 (25.2)	0.0 (0.0)
AC Taber	1.8 (1.9)	7.7 (28.4)	2.7 (1.3)	0.9 (1.2)	0.0 (0.0)
Other	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)

* Trace: less than 0.05 percent

2007 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 7

Canada Western Red Winter

Percentage of seeded acres / () = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
CDC Falcon	28.5 (32.7)	66.8 (72.4)	5.5 (6.7)	0.2 (0.9)	0.0 (0.0)
CDC Buteo	23.6 (13.2)	15.4 (9.1)	44.6 (25.9)	1.5 (0.0)	0.0 (0.0)
AC Bellatrix	10.8 (16.6)	0.0 (0.0)	9.4 (11.6)	33.1 (55.8)	0.0 (0.0)
Radiant	9.4 (6.4)	0.0 (0.0)	1.9 (3.6)	39.9 (23.2)	0.0 (0.0)
CDC Raptor	7.2 (7.7)	1.6 (3.3)	16.9 (17.5)	0.2 (0.0)	0.0 (0.0)
CDC Clair	5.6 (9.4)	1.3 (1.0)	11.8 (18.9)	2.4 (9.8)	0.0 (0.0)
CDC Osprey	5.0 (3.4)	0.0 (0.0)	5.0 (7.6)	14.3 (2.9)	0.0 (0.0)
McClintock	4.5 (3.7)	10.7 (7.0)	0.8 (2.1)	0.0 (0.0)	0.0 (0.0)
CDC Harrier	2.0 (2.8)	3.5 (4.9)	1.7 (2.2)	0.0 (0.0)	0.0 (0.0)
CDC Kestrel	1.1 (0.9)	0.8 (0.9)	2.1 (1.3)	0.0 (0.2)	0.0 (0.0)
AC Readymade	1.0 (0.7)	0.0 (0.0)	0.0 (0.0)	4.7 (3.1)	0.0 (0.0)
AC Tempest	0.7 (1.3)	0.0 (0.0)	0.1 (0.9)	2.9 (4.2)	0.0 (0.0)
Norstar	0.2 (1.2)	0.0 (1.4)	0.2 (1.6)	0.7 (0.0)	0.0 (0.0)
Other	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)

* Trace: less than 0.05 percent

2007 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 8

Canada Western Soft White Spring

Percentage of seeded acres / () = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
AC Andrew	95.5 (84.7)	100.0 (0.0)	98.7 (100.0)	79.3 (50.2)	0.0 (0.0)
Bhishaj	2.4 (1.8)	0.0 (0.0)	0.8 (0.0)	10.3 (5.9)	0.0 (0.0)
AC Meena	1.4 (10.2)	0.0 (0.0)	0.3 (0.0)	7.3 (33.2)	0.0 (0.0)
AC Nanda	0.7 (2.4)	0.0 (0.0)	0.3 (0.0)	3.0 (7.7)	0.0 (0.0)
AC Reed	0.0 (0.9)	0.0 (0.0)	0.0 (0.0)	0.0 (2.9)	0.0 (0.0)
AC Phil	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)

* Trace: less than 0.05 percent

2007 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 9

Malting -- Two-Row

Percentage of seeded acres / () = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
AC Metcalfe	59.1 (60.4)	62.0 (63.4)	58.0 (58.6)	60.0 (61.6)	100.0 (90.2)
CDC Copeland	17.6 (14.5)	10.8 (10.7)	18.1 (16.1)	18.5 (13.1)	0.0 (0.0)
CDC Kendall	12.1 (13.5)	0.1 (0.5)	15.5 (18.6)	9.0 (9.4)	0.0 (0.0)
Newdale	4.6 (2.4)	24.2 (18.5)	2.9 (0.9)	2.4 (0.9)	0.0 (0.0)
Harrington	3.3 (4.6)	0.3 (0.1)	2.6 (3.1)	5.5 (7.6)	0.0 (9.8)
Other	1.5 (2.1)	2.0 (6.9)	1.5 (1.2)	1.4 (2.4)	0.0 (0.0)
Merit	0.7 (1.7)	0.0 (0.0)	0.6 (1.1)	1.1 (3.0)	0.0 (0.0)
CDC Select	0.4 (0.1)	0.0 (0.0)	0.0 (T)	1.3 (0.2)	0.0 (0.0)
Calder	0.3 (T)	0.5 (0.0)	0.4 (0.0)	0.1 (0.1)	0.0 (0.0)
Stein	0.3 (0.7)	0.0 (0.0)	0.2 (0.2)	0.6 (1.6)	0.0 (0.0)

* Trace: less than 0.05 percent

2007 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 10

Malting -- Six-Row

Percentage of seeded acres / () = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
Legacy	49.0 (43.8)	40.7 (33.1)	55.6 (54.0)	26.0 (22.3)	0.0 (0.0)
Tradition	22.5 (12.7)	21.0 (7.2)	25.2 (18.0)	0.9 (1.2)	0.0 (0.0)
Lacey	11.0 (9.1)	16.1 (12.0)	6.5 (8.5)	30.2 (2.7)	0.0 (0.0)
Excel	7.8 (11.7)	8.5 (13.8)	7.2 (10.7)	10.1 (10.3)	0.0 (0.0)
Robust	5.8 (17.8)	10.8 (30.0)	2.7 (6.3)	11.0 (41.9)	0.0 (0.0)
CDC Yorkton	1.5 (0.0)	2.4 (0.0)	0.2 (0.0)	10.4 (0.0)	0.0 (0.0)
Other	1.1 (2.7)	0.0 (3.2)	1.3 (0.8)	5.4 (11.4)	0.0 (100.0)
CDC Battleford	0.9 (1.3)	0.5 (0.7)	0.9 (0.3)	3.9 (10.3)	0.0 (0.0)
CDC Sisler	0.2 (0.8)	0.0 (0.0)	0.2 (1.5)	2.2 (0.0)	0.0 (0.0)
CDC Laurence	0.1 (0.0)	0.0 (0.0)	0.2 (0.0)	0.0 (0.0)	0.0 (0.0)
CDC Clyde	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)
CDC Springside	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)

* Trace: less than 0.05 percent

2007 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 11

Two-Row and Six-Row Malting Barley

Percentage of seeded acres / () = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
Malting -- Two-Row	82.3 (84.2)	55.6 (55.8)	82.2 (83.6)	96.2 (95.7)	100.0 (96.2)
Malting -- Six-Row	17.7 (15.8)	44.4 (44.2)	17.8 (16.4)	3.8 (4.3)	0.0 (3.8)

* Trace: less than 0.05 percent

2007 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 12

Hulless -- Two-Row

Percentage of seeded acres / () = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
CDC McGwire	86.5 (86.9)	100.0 (96.2)	86.0 (85.5)	100.0 (100.0)	0.0 (0.0)
Other	7.3 (12.2)	0.0 (3.8)	7.6 (13.4)	0.0 (0.0)	0.0 (0.0)
Millhouse	3.7 (0.0)	0.0 (0.0)	3.8 (0.0)	0.0 (0.0)	0.0 (0.0)
CDC Fibar	2.5 (0.0)	0.0 (0.0)	2.6 (0.0)	0.0 (0.0)	0.0 (0.0)
CDC Gainer	0.0 (1.0)	0.0 (0.0)	0.0 (1.1)	0.0 (0.0)	0.0 (0.0)
CDC Rattan	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)

* Trace: less than 0.05 percent

2007 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 13

Hulless -- Six-Row

Percentage of seeded acres / () = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
AC Bacon	100.0 (9.5)	0.0 (0.0)	100.0 (0.0)	100.0 (33.3)	0.0 (0.0)
Tyto	0.0 (86.4)	0.0 (0.0)	0.0 (94.4)	0.0 (66.7)	0.0 (0.0)
Other	0.0 (4.0)	0.0 (0.0)	0.0 (5.6)	0.0 (0.0)	0.0 (0.0)
Falcon	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)

* Trace: less than 0.05 percent

2007 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 14

Feed Barley -- Two-Row

Percentage of seeded acres / () = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
Xena	54.9 (61.6)	4.3 (3.2)	64.6 (73.9)	59.5 (67.8)	0.0 (0.0)
Conlon	16.5 (12.1)	82.3 (85.9)	7.4 (4.9)	9.9 (2.6)	0.0 (0.0)
CDC Dolly	8.7 (9.1)	0.4 (0.8)	14.2 (12.9)	8.5 (9.6)	0.0 (0.0)
Ponoka	6.3 (2.4)	0.0 (0.0)	0.2 (0.0)	8.3 (3.2)	74.4 (100.0)
Seebe	3.2 (5.0)	0.0 (0.0)	0.2 (0.8)	4.2 (6.4)	25.6 (0.0)
Bridge	2.9 (0.0)	0.0 (0.0)	3.1 (0.0)	3.3 (0.0)	0.0 (0.0)
CDC Bold	2.5 (2.7)	0.0 (0.0)	0.7 (0.1)	3.3 (3.6)	0.0 (0.0)
Other	2.4 (4.2)	0.7 (0.1)	3.8 (2.5)	2.3 (5.1)	0.0 (0.0)
TREY	2.1 (1.5)	11.3 (7.9)	3.5 (1.6)	0.5 (0.5)	0.0 (0.0)
CDC Helgalson	0.6 (1.5)	0.9 (2.1)	2.2 (3.1)	0.1 (1.1)	0.0 (0.0)
RIVERS	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)
CDC Thompson	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)

* Trace: less than 0.05 percent

2007 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 15

Feed Barley -- Six-Row

Percentage of seeded acres / () = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
AC RANGER	23.7 (22.1)	75.5 (61.8)	19.7 (13.0)	8.3 (15.0)	0.0 (0.0)
Vivar	15.9 (7.5)	1.4 (0.0)	0.0 (0.0)	31.0 (14.6)	0.0 (0.0)
AC Rosser	13.5 (8.7)	7.0 (9.3)	37.0 (20.1)	2.3 (2.0)	0.0 (0.0)
Stander	9.3 (12.9)	1.9 (8.2)	12.1 (10.0)	10.6 (16.1)	0.0 (14.8)
AC Lacombe	7.2 (7.8)	3.7 (1.9)	4.7 (9.5)	10.1 (8.5)	0.0 (16.2)
AC Harper	6.3 (2.8)	0.0 (0.0)	2.5 (0.4)	11.0 (5.3)	0.0 (0.0)
Other	5.9 (11.4)	0.0 (6.4)	9.1 (11.2)	5.4 (13.1)	27.3 (13.0)
Brier	5.2 (10.6)	0.0 (0.0)	10.2 (28.1)	4.2 (4.0)	0.0 (0.0)
Trochu	5.1 (5.3)	0.0 (0.0)	1.6 (0.2)	9.1 (10.3)	0.0 (0.0)
AC Albright	3.5 (3.3)	0.0 (0.0)	0.1 (0.0)	5.1 (4.8)	58.1 (56.0)
Bedford	1.7 (1.8)	9.6 (10.3)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)
CDC Earl	1.2 (1.5)	0.0 (0.0)	0.0 (0.0)	2.4 (2.9)	0.0 (0.0)
Virden	1.0 (2.6)	1.0 (2.1)	2.8 (7.4)	0.0 (0.0)	0.0 (0.0)
Kasota	0.5 (1.5)	0.0 (0.0)	0.0 (0.0)	0.5 (2.8)	14.6 (0.0)
MANNY	0.0 (0.3)	0.0 (0.0)	0.0 (0.0)	0.0 (0.5)	0.0 (0.0)

* Trace: less than 0.05 percent

2007 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 16

All Barley (Malting, Feed and Hulless)

Percentage of seeded acres / () = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
Malting -- Two-Row	50.5 (51.5)	37.0 (34.9)	67.5 (68.9)	36.1 (40.9)	47.7 (75.9)
Feed -- (2 & 6 Row)	37.6 (37.9)	33.4 (37.1)	15.7 (15.7)	62.4 (57.0)	52.3 (21.1)
Malting -- Six-Row	10.8 (9.7)	29.6 (27.7)	14.6 (13.5)	1.4 (1.9)	0.0 (3.0)
Hulless -- (2 & 6 Row)	1.0 (0.9)	0.1 (0.4)	2.2 (1.9)	0.1 (0.2)	0.0 (0.0)

* Trace: less than 0.05 percent

2007 CANADIAN WHEAT BOARD VARIETY SURVEY

Table 17

All Wheat

Percentage of seeded acres / () = last year

Variety	Prairie Average	Manitoba	Saskatchewan	Alberta	British Columbia
Canada Western Red Spring	65.8 (74.0)	83.4 (86.6)	55.6 (65.8)	74.7 (80.2)	98.6 (93.7)
Canada Western Amber Durum	23.7 (17.3)	0.0 (0.2)	35.5 (27.3)	15.1 (10.3)	1.4 (0.0)
Canada Western Red Winter	5.0 (3.0)	12.5 (7.2)	3.6 (2.1)	3.6 (2.1)	0.0 (0.0)
Canada Prairie Spring Red	2.5 (2.5)	0.6 (0.3)	1.7 (1.7)	4.9 (4.9)	0.0 (3.9)
Canada Western Soft White Spring	1.4 (0.4)	0.1 (0.0)	2.2 (0.5)	0.8 (0.4)	0.0 (0.0)
Canada Western Hard White Spring	1.2 (2.7)	2.9 (5.7)	0.9 (2.1)	0.8 (1.9)	0.0 (0.0)
Canada Prairie Spring White	0.3 (0.2)	0.4 (0.1)	0.4 (0.3)	0.1 (0.0)	0.0 (2.4)
Canada Western Extra Strong	0.1 (0.1)	0.1 (T)	0.2 (0.2)	T (0.1)	0.0 (0.0)

* Trace: less than 0.05 percent