

Gonorrhoea

What is gonorrhoea?

Gonorrhoea is a disease caused by germs or bacteria. The infection is usually found at the opening of the uterus, or in the tube that carries urine from the bladder. It can also infect the rectum, throat, and pelvic organs. To find out if you have gonorrhoea, you must be examined by a doctor or nurse and have tests taken.

How is it spread?

Gonorrhoea can be spread easily from one infected person to another by having unprotected sex - not using a condom - with someone who has this disease. Gonorrhoea can be spread at any time while a person is infected, whether or not he or she has symptoms. A person can still become infected again in the future even if the person and infection are treated.

What are the symptoms?

In some cases there are no symptoms, and a person can have gonorrhoea and not know it. When symptoms develop, they usually occur two to five days after the person becomes infected, but may not appear for up to thirty days.

Women may notice a change in the amount and colour of fluid from the vagina, pain while urinating, and/or unusual menstrual bleeding. Men may notice a creamy white or yellow fluid from the penis and a burning feeling while urinating. In both men and women, a gonorrhoea infection in the rectum may cause itching, pain, bleeding, or a stringy white fluid when having a bowel movement. Gonorrhoea infections in the throat may cause a sore throat.

What are the complications?

If treated early before any complications begin, gonorrhoea causes no lasting problems. Untreated gonorrhoea can lead to many complications. If this infection is left untreated, it can spread and damage organs in the body.

Women may have difficulty getting pregnant. Men may develop an infection in the testicles, which can make it difficult to make a woman pregnant. Pregnant women may pass the infection to their baby's eyes during childbirth. This may lead to blindness if the baby is untreated. In both men and women, untreated gonorrhoea can cause joint, skin and eye problems.

What is the treatment?

Both the person diagnosed with gonorrhoea, and their sexual partner(s) within the last sixty days, must take antibiotic pills. Any sexual partner(s) of the person diagnosed with gonorrhoea must be treated, regardless of their test results. Follow up tests are recommended six months after treatment.

Important: Do not have sex until you and your sexual partner(s) have finished all the medication. Take all of the medication exactly as instructed.

Should I be tested for other sexually transmitted diseases (STDs)?

If you have different sexual partners, or if your partner has different partners, you should also be tested for other STDs including chlamydia, syphilis and HIV (the virus linked to AIDS). You should also consider getting vaccine shots to prevent hepatitis B infection.

Please remember: The more sexual partners you have, the higher your risk of getting a STD.

Birth control pills

Birth control pills may not work very well when you are taking some antibiotic medicines. Keep taking your birth control pills while taking any medication, and also use a second form of birth control, such as a condom, until your next period after completing the antibiotics.

Ways to reduce your risk of getting a sexually transmitted disease

- Have sex with only one partner who has been tested for sexually transmitted diseases (STDs), who has been treated if necessary and who is having sex only with you. The more partners you have, the higher your risk of getting an STD.
- Use a female or male condom every time you have sex. Condoms offer protection against STDs, but they must be used properly.
- Have regular check-ups for STDs.
- Decide not to have sex.

Important facts about condoms

- A condom acts like a barrier that helps prevent the exchange of body fluids, the transmission of sexually transmitted infections, and pregnancy.
- A new condom should be used each time you have sexual intercourse (anal, vaginal, or oral sex).
- Check the expiry date on the condom package.
- Use only water-based lubricants with the male latex condom. Oil-based lubricants, such as petroleum jelly, lotion or baby oil, can weaken and destroy latex.
- Female condoms are made of polyurethane. This material can be used with any type of lubricant, water-based or oil-based.
- Some lubricants contain chemicals called spermicides to help protect against unwanted pregnancy. If they irritate your genitals, don't use them.
- Do not use a male condom together with a female condom as the friction created may cause tearing of either product.
- If a condom breaks during sex remove it immediately and apply a new condom.
- Remember! Condoms do not offer 100 per cent protection from STDs and unwanted pregnancy. It will not consistently prevent transmission of STDs passed through skin-to-skin contact – for example, syphilis, human papilloma virus (warts) and herpes. However, if used properly, they are very effective and can reduce the risk of transmission of these STDs.

Putting on a male condom

- Take the condom carefully out of the packet.
- Place the condom on the tip of the penis when it is hard and erect, but before it touches the partner's body. Make sure that the rolled-up condom rim faces outward.
- With the other hand, pinch the tip of the condom to remove any trapped air, and unroll the condom to the base of the erect penis.

- After intercourse and before the penis becomes soft, withdraw the penis carefully, holding the rim of the condom against the penis, so that semen does not spill out.
- Slide the condom gently off the penis, and knot the open end.
- After using the condom, throw it in the garbage.

Putting on a female condom

- A condom can be inserted up to eight hours before sexual intercourse.
- Open the package carefully. Hold the small ring at the closed end of the condom between the thumb and middle finger.
- Find a comfortable position, either lying down, sitting with your knees apart or standing with one foot raised on a stool, squeeze the small ring and insert it into the vagina as far as you can.
- Put a finger inside the condom and push the small ring inside as far as possible. It is also possible to insert the condom by putting it onto the erect penis before intercourse.
- Make sure that the part of the condom with the outer ring is outside the body. The outer ring will lie flat against the body when the penis is inside the condom.
- When the penis enters the vagina, make sure that the penis is inside the condom.
- Immediately after sexual intercourse, remove the condom by gently twisting the outer ring and pulling the condom out, making sure that no semen is spilt and throw it in the garbage.



BC Centre for Disease Control
AN AGENCY OF THE PROVINCIAL HEALTH SERVICES AUTHORITY

For more BC HealthFile topics visit
www.bchealthguide.org/healthfiles/index.stm,
or visit your local public health unit.

Call the BC NurseLine to speak to a
registered nurse, available 24-hours every
day:

- In Greater Vancouver, call 604-215-4700
- In BC, call toll-free 1-866-215-4700
- Deaf and hearing-impaired, call 1-866-889-4700
- Pharmacist available 5pm to 9am every day
- Translation services in over 130 languages upon request.

Visit BC HealthGuide OnLine – a world of
health information you can trust at
www.bchealthguide.org