FISH FACTS 2003

Results from fishery surveys conducted during the summer of 2003 are shown on the following pages. Lakes are mainly surveyed using two types of equipment; gill nets or downhaul trap nets. Both types of equipment are effective in capturing fish. Fish caught in downhaul traps can generally be released after being measured. Fish caught in gill nets are not releasable but can be used to gather additional information. For this reason, the size of fish caught in downhaul trap nets is recorded by length, while size of fish captured in gill nets is recorded by weight.

If you have any questions on survey methods or equipment, or if you have concerns about the health of a specific fishery, please contact your nearest Saskatchewan Environment fisheries biologist.

Waterbody and Location	Species Caught (Weight)	Date Tested/Survey Gear/
	and Population Status	Remarks
Anglin Lake	Walleye: (40–1800g) Fair	June 17/Gill nets. Angling
53°44' 105°56'	Northern Pike: (100–	success for walleye and
55 - 27W2	1725g) Moderate	pike was reported fair at
N of Prince Albert	Perch: (30–140g) Fair	time of testing.
Atchison Lake	Rainbow Trout: (14-38	July 23-24/Gill nets set at 3
54°50′ 108°10′	cm) Fair	locations. Survey indicates
N. of Meadow Lake	Brook Trout: (24-47 cm)	a good population of brook
	Good	trout, fair population of
		rainbow trout.
Atton Lake	Northern Pike: (1880g)	July 16/Gill nets. Although
52°51' 108°52'	Sparse	net catches poor, angling
4,5-45-20W3	Perch: (175-280g) Poor	fairly successful and scuba
NE of Cut Knife		divers were seeing lots of
		fish.
Baldy Lake	Walleye: (100–1800g) Poor	June 11/Gill nets. Fair
54°06' 104°38'	Northern Pike: (475–	success for pike reported.
16, 21–59–18W2	1175g) Moderate	
Narrow Hills Provincial	_	
Park		
Bittern Lake	Walleye: (450–1100g)	May 23/Gill nets. Some
53°56' 105°45'	Moderate	winterkill took place during
57,58–26W2	Northern Pike: (130–	the winter of 2002/03 but
S.of Montreal Lake	1375g) Good	not enough to affect the fish
	Perch: (35–300g) Fair	population.
	Whitefish: 2	

Burtlein Lake 54°15' 104°38'	Tiger Trout: (80–700g) Moderate	June 13/Gill nets. Stocked every second year with
3, 4–61–18W2 S. of Little Bear Lake	Splake: (200–675g) Moderate	splake and with tiger trout when available. Fishing is reported to be fair.
Burton Lake 52°16' 105°07' 17,18-38-22W2 N of Humboldt	Walleye: (50-1900g) Good Northern Pike: (1200g) Sparse	June20/Gill nets. Walleye are stocked every two years: last stocked in 2002. In the past perch were also present in net catches.
Bushy Lake 52°51' 108°49' 3-45-20W3 NE of CutKnife	Walleye: (155-600g) Good Perch: (up to 400g) Fair	July 17/Gill net. Walleye fry stocked every second year. Perch are a very good size. Angling pressure has been light.
Canoe Lake 55°10' 108°15' W of Beauval	Walleye: (18-57 cm) Good Northern Pike: (36-88 cm) Good	August 18-25/Trap nets. Set at 3 sites. Good numbers of pike and walleye at most test net locations. Good angling success reported for pike and walleye. Walleye limit of 3.
Chris Johnson Lake 51°52' 102°08' 30, 31–33–1W2 S.W. of Norquay	Northern Pike: (375–1400g) Good	September 11/Gill nets. No perch were caught in the test nets.
Clarence Lake 54°14' 105°00' 60, 61–20-21W2 S of East Trout Lake	Walleye: (1550–3700g) Moderate Northern Pike: (500– 1725g) Fair Perch: (30–50g) Sparse	August 6/Gill nets. No reports of angling success.
Crooked Lake 50°36' 102°44' 19-5,6W2 S of Melville	Walleye: (120-2000g) Good Northern Pike: (500- 6000g) Fair	July 15/Gill nets. Water levels are low due to a compensation claim regarding flooding First
5 OI MENTILE	Perch: (100-350g) Good	Nations land. Access at boat launches is difficult. Moderate angling success is reported.

Crystal Lake	Walleye: (140–1200g)	September 11/Gill nets. Fair
51°51' 102°26'	Moderate	angling success reported for
33, 3, 4–W2	Northern Pike: (700 g)	walleye and pike.
N of Canora	Poor	
	Perch: (40–50 g) Poor	
Cutarm Reservoir	Walleye: (200-1000g)	August 7/ Gill nets. Stocked
50°57' 102°10'	Good	annually with 400,000
9,16-20-32W1	Perch: (100-330g) Good	walleye. Perch were
E of Esterhazy		introduced in 2000. Good
		angling success for perch
		and walleye is reported.
D'Amour Lake	Walleye: (300–1200g)	June 24/Gill nets. Stocked
53°04' 107°00'	Moderate	with 200,000 walleye fry
17,18–47–7W3	Northern Pike: (500–	annually.
W of Leask	1350g) Fair	
22	Perch: (40 g) Sparse	
Dog Lake	Rainbow Trout: (350–	June 19/Gill nets. Good
54°28' 106°05'	1300g) Moderate	angling success reported.
28–63–1W3	1300g) Woderate	Stocked annually with 7700
W of Weyakwin		rainbow trout.
Dore Lake	Walleye: (15-78 cm)	June 9-25/Trap nets. Set at
54°46' 107°17'	Recovering	6 sites. Pike - Fair to poor
NE of Meadow Lake	Northern Pike: (44-94 cm)	angling success reported.
TVL of Weddow Lake	Fair	Trap net catch per unit
	T un	effort (CPUE) decreased
		citori (Ci OL) decreased
i		, ,
		from 1998 to 2003. Walleye
		from 1998 to 2003. Walleye – recovering. Trap net
		from 1998 to 2003. Walleye – recovering. Trap net CPUE numbers increasing
		from 1998 to 2003. Walleye – recovering. Trap net CPUE numbers increasing from 1999-2003, walleye
		from 1998 to 2003. Walleye – recovering. Trap net CPUE numbers increasing from 1999-2003, walleye continue to show recovery,
		from 1998 to 2003. Walleye – recovering. Trap net CPUE numbers increasing from 1999-2003, walleye continue to show recovery, strong 1, 2, 3, 5+ year old
		from 1998 to 2003. Walleye – recovering. Trap net CPUE numbers increasing from 1999-2003, walleye continue to show recovery, strong 1, 2, 3, 5+ year old age classes, but a weak 4
		from 1998 to 2003. Walleye – recovering. Trap net CPUE numbers increasing from 1999-2003, walleye continue to show recovery, strong 1, 2, 3, 5+ year old age classes, but a weak 4 year old age class evident in
		from 1998 to 2003. Walleye – recovering. Trap net CPUE numbers increasing from 1999-2003, walleye continue to show recovery, strong 1, 2, 3, 5+ year old age classes, but a weak 4 year old age class evident in sample. Walleye limit of 1.
		from 1998 to 2003. Walleye – recovering. Trap net CPUE numbers increasing from 1999-2003, walleye continue to show recovery, strong 1, 2, 3, 5+ year old age classes, but a weak 4 year old age class evident in sample. Walleye limit of 1. Fair angling success
		from 1998 to 2003. Walleye – recovering. Trap net CPUE numbers increasing from 1999-2003, walleye continue to show recovery, strong 1, 2, 3, 5+ year old age classes, but a weak 4 year old age class evident in sample. Walleye limit of 1. Fair angling success reported in spring.
Elaine Lake	Walleye: (850–1950g)	from 1998 to 2003. Walleye recovering. Trap net CPUE numbers increasing from 1999-2003, walleye continue to show recovery, strong 1, 2, 3, 5+ year old age classes, but a weak 4 year old age class evident in sample. Walleye limit of 1. Fair angling success reported in spring. June 18/Gill nets. Stocked
54°27' 106°22'	Sparse	from 1998 to 2003. Walleye recovering. Trap net CPUE numbers increasing from 1999-2003, walleye continue to show recovery, strong 1, 2, 3, 5+ year old age classes, but a weak 4 year old age class evident in sample. Walleye limit of 1. Fair angling success reported in spring. June 18/Gill nets. Stocked every second year with 1.3
		from 1998 to 2003. Walleye recovering. Trap net CPUE numbers increasing from 1999-2003, walleye continue to show recovery, strong 1, 2, 3, 5+ year old age classes, but a weak 4 year old age class evident in sample. Walleye limit of 1. Fair angling success reported in spring. June 18/Gill nets. Stocked

Fairy Glen Lake	Walleye: (65–1600g) Fair	August 26/Gill nets. Fair
54°05' 104°46'	Northern Pike: (225–450g)	angling success was
10, 59-10W2	Poor	reported for walleye.
W of Upper Fishing Lake		
From Lake	Walleye: (175-350g)	October 7/Gill nets. Stocked
34-19-31W1	Excellent	annually with 20,000
E of Esterhazy		walleye. Complete
		winterkill occurred in
		2000/2001, so walleye are
		small yet.
Good Spirit Lake	Walleye: (55-950g)	August 1/Gill nets. Access
51°34' 102°40'	Excellent	at boat launches is difficult
29,30-5W2	Northern Pike: (750-	due to low water levels.
Good Spirit Provincial Park	2250g) Poor	Good angling success in the
	Perch: (325-420g) Poor	spring and summer was
		reported.
Grenfell Reservoir	Walleye: (100-850g)	August 27/Gill nets.
30-16-7W2	Excellent	Stocked annually with
S of Grenfell		50,000 walleye. Most
		walleye are currently too
		small to interest anglers, but
		an excellent fishery is
		developing.
Hackett Lake	Walleye: (19-52 cm) Good	August 7-12/Trap nets.
59°29' 105°18'	Northern Pike: (38-98 cm)	Sets at 4 locations. Good
	Fair	angling reported for pike
		and walleye; trophy pike
		present but scarce (at levels
		similar to past surveys).
		<u> </u>
		Proposal introduced in 2004
		Proposal introduced in 2004 to reduce walleye angling
		Proposal introduced in 2004 to reduce walleye angling limits from 5 to 3, over
		Proposal introduced in 2004 to reduce walleye angling limits from 5 to 3, over concern of increased
	W. H. (20.72)	Proposal introduced in 2004 to reduce walleye angling limits from 5 to 3, over concern of increased walleye harvest.
Helene Lake	Walleye: (28-50 cm) Good	Proposal introduced in 2004 to reduce walleye angling limits from 5 to 3, over concern of increased walleye harvest. August 19-21/Trap nets.
53°33' 108°12'	Walleye: (28-50 cm) Good	Proposal introduced in 2004 to reduce walleye angling limits from 5 to 3, over concern of increased walleye harvest. August 19-21/Trap nets. One location. Lake has a
	Walleye: (28-50 cm) Good	Proposal introduced in 2004 to reduce walleye angling limits from 5 to 3, over concern of increased walleye harvest. August 19-21/Trap nets. One location. Lake has a history of partial winterkills
53°33' 108°12'	Walleye: (28-50 cm) Good	Proposal introduced in 2004 to reduce walleye angling limits from 5 to 3, over concern of increased walleye harvest. August 19-21/Trap nets. One location. Lake has a history of partial winterkills since 1999, however,
53°33' 108°12'	Walleye: (28-50 cm) Good	Proposal introduced in 2004 to reduce walleye angling limits from 5 to 3, over concern of increased walleye harvest. August 19-21/Trap nets. One location. Lake has a history of partial winterkills since 1999, however, survey shows good numbers
53°33' 108°12'	Walleye: (28-50 cm) Good	Proposal introduced in 2004 to reduce walleye angling limits from 5 to 3, over concern of increased walleye harvest. August 19-21/Trap nets. One location. Lake has a history of partial winterkills since 1999, however, survey shows good numbers of walleye 3 to 4+ years
53°33' 108°12'	Walleye: (28-50 cm) Good	Proposal introduced in 2004 to reduce walleye angling limits from 5 to 3, over concern of increased walleye harvest. August 19-21/Trap nets. One location. Lake has a history of partial winterkills since 1999, however, survey shows good numbers of walleye 3 to 4+ years old, but 2 and 3 year old
53°33' 108°12'	Walleye: (28-50 cm) Good	Proposal introduced in 2004 to reduce walleye angling limits from 5 to 3, over concern of increased walleye harvest. August 19-21/Trap nets. One location. Lake has a history of partial winterkills since 1999, however, survey shows good numbers of walleye 3 to 4+ years old, but 2 and 3 year old fish absent in sample. Poor
53°33' 108°12'	Walleye: (28-50 cm) Good	Proposal introduced in 2004 to reduce walleye angling limits from 5 to 3, over concern of increased walleye harvest. August 19-21/Trap nets. One location. Lake has a history of partial winterkills since 1999, however, survey shows good numbers of walleye 3 to 4+ years old, but 2 and 3 year old fish absent in sample. Poor to fair angling success
53°33' 108°12'	Walleye: (28-50 cm) Good	Proposal introduced in 2004 to reduce walleye angling limits from 5 to 3, over concern of increased walleye harvest. August 19-21/Trap nets. One location. Lake has a history of partial winterkills since 1999, however, survey shows good numbers of walleye 3 to 4+ years old, but 2 and 3 year old fish absent in sample. Poor to fair angling success reported for walleye. Poor
53°33' 108°12'	Walleye: (28-50 cm) Good	Proposal introduced in 2004 to reduce walleye angling limits from 5 to 3, over concern of increased walleye harvest. August 19-21/Trap nets. One location. Lake has a history of partial winterkills since 1999, however, survey shows good numbers of walleye 3 to 4+ years old, but 2 and 3 year old fish absent in sample. Poor to fair angling success

Incomois I als:	Wellers (40, 2000 -) C 1	Angust 12/C:11 E-:-
Iroquois Lake	Walleye: (40–2800g) Good	August 13/Gill nets. Fair
53°10′ 107°01′	Northern Pike: (240-	angling success was
48 – 7,8W3	1250g) Fair	reported for walleye,
S of Shell Lake Town	Perch: (25–40g) Poor	northern pike and whitefish.
	Whitefish: (200–1900g)	
	Good	
Jan Lake	Walleye: (25–2400g) Fair	June 18-August 1, 2002/Gill
54°56' 102°55'	Northern Pike: (182-	nets. Good angling success
SE of Pelican Narrows	5500g) Fair	for walleye and pike.
	Perch: Poor	
Jean Lake	Perch: Nil	July 22/Gill nets. A
52°48' 103°13'		winterkill took place during
22, 23–44–9W2		the winter of 2002/03
S. E. of Mistatim		possibly due to low water
		conditions.
Jeannette Lake	Walleye: (18-40 cm) Fair	July 16-25/Trap nets and
54°33' 108°32'	Northern Pike: (52-81	Gill nets. Large walleye
Meadow Lake Provincial	cm) Poor	absent in population, but
	CIII) POOI	average size slightly larger
Park		than 2000 and 2002
		surveys. Only 5 pike
		caught in 2003. Good
		angling success reported for
T7 1 7 1	XX II (26.64)	walleye.
Keeley Lake	Walleye: (26-64 cm)	May 21-June 6/Trap nets.
54°54' 108°08'	Excellent	Set at 6 locations caught
NE of Meadow Lake	Northern Pike: (20-96 cm)	good to excellent numbers
	Good	of walleye and pike at most
		sites. Good angling
		success reported for pike
		and walleye. Walleye limit
		of 3.
Lac la Plonge	Lake Trout: (44-88 cm)	Sept.22-Oct.7/Trap nets. Set
55°08' 107°20'	Excellent	at 2 locations. Lake trout
E. of Beauval		Spawning surveys of 2002-
		03 indicate good CPUE
		numbers of lake trout in all
		year classes. Anglers report
		fair success for lake trout
		and pike.
		and pixe.

Little Jackfish Lake	Perch: (100-290g)	June 12/Gill nets. Has been
25,26-30-30W1	Excellent	stocked annually with 6,000
Duck Mountain Provincial	Rainbow Trout: Sparse	rainbow trout and 3,000
Park	Tiger Trout: Sparse	tiger trout. Perch were
		illegally introduced in the
		late 1990's, and have out-
		competed the trout. Further
		trout stocking will be ceased.
Little Loon Lake	Walleye: (80-640g) Good	Aug 8/Gill nets. Walleye
53°21' 108°15'		fry stocked every two years.
		Angling pressure fairly
34,35-50-16W3		heavy but poor success for
E of Glaslyn		walleye. Pike have not
		been caught in gillnets since
		1993.
Margo Lake	Walleye: (200-1580g)	July 23/Gill nets. Walleye
51°49' 103°21'	Good	populations remain good,
9-33-10W2	Northern Pike: (480-880g)	although decreased slightly
SW of Margo	Moderate	since the 1999 survey.
_	Perch: (50g) Sparse	Perch was captured for the
		first time in gillnets.
Martyns Lake	Walleye: (325–1300g) Poor	July 22/Gill nets. No reports
52°37′ 103°22′	Northern Pike: (475–850g)	of fishing success.
23–42–10W2	Fair	
N. W. of Porcupine Plain	Perch: (25–85 g) Moderate	1.1.22/631
Moosomin Reservoir	Walleye: (120-3150g)	July22/Gill nets. Stocked
50°04' 101°42'	Good	every second year with 1
13-31W1 SW of Moosomin	Northern Pike: (1200-	million walleye. Poor
S W OI MOOSOIIIII	6800g) Poor Perch: (100-400g)	angling success for perch
	Moderate	and walleye, and good success for pike is reported.
Narrow Lake	Walleye: (800–1100g) Poor	July 29/Gill nets. No reports
54°25' 104°22'	Northern Pike: (200–550g)	of angling success.
62,63–16W2	Fair	or ungining success.
N of Little Bear Lake		
Nelson Lake	Walleye: (800–2450g)	July 23/Gill nets. Stocked
52°01' 102°45'	Moderate	annually with 100,000
14–35–6W2	Northern Pike: (2000 g)	walleye fry.
N. W. of Preeceville	Poor	
	Perch: (30–130g) Fair	

Nemeiben Lake	Walleye: (200–2600g) Poor	May 29–August 20/Gill
55°20' 105°20'	Northern Pike: (54-5500g)	nets.
NW of La Ronge.	Fair	nets.
NW of La Ronge.	Lake Trout: (50–5000g)	
	Poor (30–3000g)	
	Perch: Poor	
Oscar Creek	Brook Trout: Nil	Oct 1-2/Gill nets and
25-44-8W3	Brook frout; Mil	
		backpack electroshocker. Brook trout not stocked
N of Krydor		since 2000. No evidence of
		survival – low waters in last
		few years may have had
	77 77 77 77	impact.
Oyama Reservoir	Northern Pike: (550-	August 21/Gill nets. A
36-15-17W2	4350g) Good	partial winterkill occurred
E of Kronau	Perch: (80-95g) Fair	in 2002/2003, mostly
		affecting perch. Good
		angling success for pike
		was reported this spring.
Pagan Lake	Northern Pike: (26-62 cm)	May 13-15, Sept. 11-
54°19' 108°16'	Fair	13/Trap nets. Sets at 3
		locations caught fair
		numbers of pike; no large
		pike caught in recent
		surveys, low numbers of
		perch caught.
Pinkney Lake	Walleye: (80–1425g)	July 30/Gill nets. Fair
54°03' 105°05'	Moderate	success reported for walleye
28,32,33–58–21W2	Northern Pike: (500–	and pike.
N of Whiteswan Lake	1075g) Moderate	
Rafferty Reservoir	Walleye: (40-2850g) Good	September 12/Gill nets.
49°09' 103°15'	Northern Pike: (850-	Fair to good angling success
2-9W2	4900g) Poor	for walleye and northern
NW of Estevan	Perch: (325g) Fair	pike. Northern pike size
	_	and number are probably
		larger than survey indicates.
		Yellow perch becoming
		good size.
Ridge Lake	Brook Trout: (170-1375g)	June 10/Gill nets. Stocked
54°16′ 105°02′	Good	every second year with
11–61–21W2		4,000 brook trout
N. W. of Piprell Lake		fingerlings.
or ripron Dune	l .	

Road Lake 54°20' 104°33'	Rainbow Trout: (240-1000g) Moderate	June 11/Gill nets. Good success reported by anglers.
1–63–18W2 E of Little Bear Lake		
Round Lake	Walleye: (60-3700g) Good	July 16-17/Gill nets. Water
50°32' 102°22'	Northern Pike: (750-	levels are low due to a
18-3W2	2750g) Moderate	compensation claim
SE of Melville	Perch: (100-400g)	regarding flooding First
	Moderate	Nations land. Access at boat
	Catfish: (5650-5850g) 2	launches is difficult.
	_	Moderate angling success is
		reported.
Round Lake	Walleye: Nil	May 13/Gill nets. A
52°22' 103°24'	Northern Pike: Nil	complete winterkill took
16,17,20,21–39–20W2		place 2002/03. Stocked with
S.W of Porcupine Plain		500,000 walleye fry spring
		2003.
Sealey Lake	Rainbow Trout: (350–	June 13/Gill nets. Stocked
54°16' 104°36'	1225g) Moderate	annually with 12,000
10,14,15–61–18W2	Brook Trout: (70–450g)	rainbow and 6,000 brook
S. E of Little Bear Lake	Good	trout fingerlings. Fair
		success reported for both
	D 15 (60 220)	trout species.
Sedge Lake	Brook Trout: (60–220g)	June 19/Gill nets. Stocked
54° 27' 106° 05'	Good	annually with 1,600 brook
22–63–1W3		trout fingerlings.
W of Weyakwin Lake	D 1 (42.79) C 1	A 15.6/E 1. C.1
Sled Lake	Pike (42-78 cm) Good	August 5-6/Trap nets. Sets
54°27' 107°25' S. of Dore Lake		at 3 locations. Abundant
S. of Dore Lake		pike population, although no trophy pike caught in
		any location.
Smuts Lake	Walleye: Nil	June 24/Gill nets. No
52°29' 106°05'	Perch: Nil	gamefish caught in gill nets,
33-40,41-1W3	2 02 044 1 (11	and seining in July caught
SW of Alvena		no perch. Walleye and
		perch have been restocked.
		Lake severely winterkilled
		during 2002/03.
Steepbank Lake	Walleye: (2100 g) Sparse	August 7/Gill nets. No
54°16' 105°06'	Northern Pike: (190–	reports of angler success.
9,17,18–61–21W2	2200g) Poor	
S of East Trout Lake		

Steep Creek 53°14' 105°21' 18–49–23W2 East of Prince Albert	Brook Trout: (40–190g) Moderate	July 24/Gill nets. Stocked annually with 500 brook trout fingerlings.
Struthers Lake 52°51' 105°15' 2, 3–45–23W2 S. E. of Birch Hills	Walleye: Nil Northern Pike: Nil Perch: Nil	July 7/Gill nets. Stocked with 300,000 walleye fry. A complete winterkill took place during the winter of 2002/03
Sturgeon Lake 53°25' 106°05' 51–1,2W3 North of Holbien	Walleye: (160–1550g) Moderate Northern Pike: (425– 1425g) Moderate Perch: (25–130g) Moderate	June 25/Gill nets. Stocked every second year with 1.4 million walleye fry.
Tobin Lake 53°35' 103°30' E. of Nipawin	Walleye: (60–5400g) Good Northern Pike: (925– 8600g) Moderate Sauger: (130g) Sparse Perch: (30–450g) Moderate Goldeye: (50–450g) Moderate	June 3-6 July 15-17 August 19-21/Gill nets. Good success for walleye and pike reported.
Turtle Lake 53°36' 108°38' SW of Meadow Lake	Walleye (16-71 cm) Good Pike (32-102 cm) Good	July 2-15/Trap nets. Sets at 5 locations indicate walleye numbers increasing compared with previous surveys. Pike: Good percentage of large pike caught in sample, with strong year classes of young pike. Good angling for pike reported, fair for walleye.
Usinneskaw Lake 51°47' 103°22' 33-32-10W2 S of Margo	Walleye: (80-1520g) Excellent Northern Pike: (620- 1100g) Fair	July 25/Gill nets. Walleye populations continue to be excellent. Locals report a muddy taste in the larger walleye in late summer.
Whitesand Lake 51°46' 103°20' 21,28-32-10W2 S of Margo	Walleye: (275-1500g) Excellent Northern Pike: (640- 1275g) Fair	July 24/Gill nets. Walleye populations have increased in abundance and are rated excellent. Lake is stocked with fry every three years.

Wishtikan Lake 52°50' 108°53' 31,32-44-21W3 NE of Cut Knife	Perch: (80-280g) Excellent	July 15/Gill nets. An excellent population of angling-sized perch is present in this waterbody. Good fishing has been reported.
Wolseley Reservoir 36-16-10W2 S of Wolseley	Walleye: (150-600g) Excellent Northern Pike: (800- 2950g) Good Perch: (140-150g) Poor	August 22/Gill nets. Stocked annually with 30,000 - 50,000 walleye fry. This is the first time in many years an excellent walleye population has developed due to heavy pike predation and periodic winterkill.