

# Urine collection practices in a small rural hospital: Evaluation of alignment with antimicrobial stewardship guidelines

Jennifer Irwin, RN;<sup>1</sup> Leslie Forrester, BA (Hons), MA, MSc<sup>2</sup>

<sup>1</sup> Infection Control Practitioner, Vancouver Coastal Health Authority, Powell River General Hospital, Powell River, BC, Canada

<sup>2</sup> Regional Director Data Analytics, Vancouver Coastal Health Authority, Powell River General Hospital, BC, Canada

## Corresponding author:

Jennifer Irwin, RN, Infection Control Practitioner  
Vancouver Coastal Health Authority  
Powell River General Hospital  
5000 Joyce Ave.  
Powell River, BC V8A 5R3 Canada  
Tel.: 604-485-3214  
Fax: 604-485-3245  
jennifer.irwin@vch.ca

## ABSTRACT

**Background:** The diagnosis of a symptomatic urinary tract infection (UTI) can be challenging among elderly patients, resulting in an increased risk for specimen collection and treatment of asymptomatic patients. The Vancouver Coastal Health Authority's Antimicrobial Stewardship Guidelines provide a systematic approach to support clinicians in the assessment and treatment of UTIs. The purpose of this quality review was to evaluate if urine collection practices were in alignment with the guidelines.

**Methods:** The review involved a retrospective examination of all urine specimens collected from medical and surgical patients of a small rural hospital from September 1, 2015 to August 31, 2016. A survey of staff was also conducted to evaluate the interdisciplinary team's knowledge of urine collection and interpretation and to assess concordance with the guidelines.

**Results:** 318 urine cultures were reviewed, of which only 78 (24.5%) met microbiologic criteria and were considered to be positive cultures. A large proportion of samples resulted in mixed organisms. There was a highly statistically significant relationship between urinalysis and a positive culture (Fisher's exact test  $p < 0.0001$ ). A positive urine culture ( $> 100$  million CFU/L) was statistically significantly more likely to be observed when there was a positive urinalysis.

**Conclusions:** Older adults represent a large and growing population of hospitalized patients. Diagnosis of a symptomatic UTI in the elderly can pose challenges. Urine cultures are frequently obtained and a high proportion sent for culturing with missing or negative urinalysis. Urinalysis results are pertinent in the diagnosis of a UTI as there is correlation between a positive urinalysis and a positive urine culture. Adoption of a systematic approach in the management of UTIs would result in consistent and appropriate assessment and treatment of UTIs for elderly patients.

## KEYWORDS

Elderly; urinary tract infections; urinalysis; antimicrobial stewardship; systematic approach

## INTRODUCTION

Urinary tract infections (UTIs) have been identified as the single most common healthcare-acquired infection, accounting for 40% of all hospital-acquired infections and resulting in a substantial burden for patients and the healthcare system [1-3]. UTIs impact patient morbidity and mortality, which can lead to delirium, falls, and an increased length of stay in hospital [4, 5].

Older adults are at a greater risk for UTIs due to increased post-void residual volume, prostatic hypertrophy, age-associated changes in immune function, and a growing number of comorbidities [4, 6-8]. Hospitalized older adults generally have more functional impairments, higher cognitive deficits, and a greater number of medical comorbidities. Medical

comorbidities, such as a stroke and dementia, may predispose individuals to bowel and bladder incontinence, which has been associated with symptomatic UTIs and persistent asymptomatic bacteriuria [6]. However, the most significant risk factor associated with UTIs in hospitalized patients is the presence of a urinary catheter. Approximately 80% of healthcare-acquired UTIs are catheter-associated [1, 4, 8].

The diagnosis of a symptomatic UTI requires both the presence of genitourinary symptoms in addition to a positive urine culture [2, 6]. Elderly patients who suffer from significant cognitive deficits that impair their ability to communicate and chronic genitourinary symptoms (e.g., incontinence, urgency, and frequency) make the diagnosis of symptomatic UTIs

**Acknowledgements:** None.

**Conflicts of interest:** None.

**Funding:** None.

**FIGURE 1: The ASPIRES Urinary Tract Infections Management Algorithm.**



**Antimicrobial Stewardship Programme:**  
**Innovation, Research, Education, and Safety**  
 Quality and Patient Safety, Vancouver Coastal Health  
 Vancouver General Hospital  
 855 West 12<sup>th</sup> Avenue, Vancouver, BC, V5Z1M9, Canada

**VCH Management of Urinary Tract Infections (UTI) in Non-pregnant Adults**

**KEY POINTS:**

1. Malodorous/cloudy urine alone is **NOT** a sign/symptom of UTI and is **NOT** an indication to obtain urine cultures <sup>(1)</sup>
2. Changes in cognitive function and activities of daily living **REQUIRE** clinical assessment; never assume these are due to UTI
3. Urine should **ALWAYS** be collected midstream, by in/out catheterization, or through a new catheter (unless contraindicated) <sup>(2)</sup>
4. Positive urine cultures in asymptomatic patients should **NOT** be treated except in pregnancy or prior to urologic/gynecologic surgery


**Signs and Symptoms of Suspected UTI <sup>(3)</sup>**

-One of the following in febrile patients (oral temperature >37.8 °C [or 1°C above baseline in Spinal Cord Injury]) or **two** of the following in afebrile patients:

- |  |  |
|--|--|
| i. Acute dysuria                               | vi. Suprapubic pain  |
| ii. New or marked increase in incontinence     | vii. Gross hematuria   |
| iii. New or marked increase in urgency         | viii. Swelling, or tenderness of the testes, epididymis, or prostate |
| iv. New or marked increase in frequency        | ix. New-onset of acute costovertebral angle pain or tenderness       |
| v. New or marked increase in urinary retention | x. Episode of autonomic dysreflexia (with no other apparent cause)   |

**NOTE:** Only after clinical assessment and ruling-out of other possible causes should changes in mental status and functional decline, and sudden fever, rigors or new-onset hypotension suggest UTI in patients; use clinical judgment. <sup>(3)</sup>

**For Geriatric and Spinal Cord Injury (including conus/cauda equina):** UTI may present atypically; use clinical assessment to guide decision for urine culture & urinalysis.



VCH UTI Algorithm

ASPIRES

Revised: August 8, 2017

challenging [6]. Based on these challenges, there is an increased risk for specimen collection and treatment of asymptomatic patients.

The Antimicrobial Stewardship Programme: Innovation, Research, Education and Safety (ASPIRES) has been part of the Vancouver Coastal Health Authority's (VCH) Quality and Patient Safety department since 2012. One of the many contributions made by the ASPIRES team was the development of the Urinary Tract Infections Management Algorithm, which provides a systematic approach in the assessment and treatment of UTIs in non-pregnant adults, as seen in Figure 1.

As urine testing often drives prescribing, a key component of antibiotic stewardship around UTIs must be in the ordering and interpretation of urine tests. In the absence of a standardized approach, there is an increased risk of over-collection of samples and treatment of asymptomatic patients. Interpretation of the significance of the bacterial culture results requires evaluation of a number of factors, including clinical signs and symptoms, urinalysis results, specimen collection and storage, and antibiotic treatment [7, 9].

Urinalysis specimens provide critical information for interpretation of results and should be collected in coordination with urine for culturing [4, 8, 10]. According to recent studies, the minimum laboratory evaluation for a suspected UTI should include urinalysis for determination of leukocyte esterase (LE) and nitrate (NIT) levels by use of dipstick and microscopic evaluation for white blood cells. If the urinalysis is negative for LEs and NIT, a positive culture is very unlikely [6, 10, 11].

To prevent contamination, urine specimens should be collected using a clean-catch midstream technique or by intermittent catheter. The clean-catch midstream approach is influenced by the patient's physical ability to perform the task and adhere to the instructions provided by the healthcare worker. Specimens should not be collected while using a bedpan, as they are not sterile and there is risk for urine to contact the surrounding perineum, resulting in contamination. Indwelling catheters must be replaced prior to obtaining specimens, as the inner and outer surfaces of the lumen develop a biofilm within days of insertion. Once a biofilm has been established, it protects the uropathogens from antimicrobial treatment and provides an environment that supports bacterial growth [8].

With respect to storage, urine specimens should be kept refrigerated and transported to the microbiology lab without delay. A delay in processing can lead to inaccurate results due to bacterial growth [10]. This is particularly pertinent to Powell River General Hospital (PRGH), as the Microbiology department was recently closed. Since June 2014, all urine specimens for culturing have been sent to the microbiology laboratory in Vancouver General Hospital (VGH) for processing. Urinalysis specimens continue to be processed at PRGH.

Although the ASPIRES algorithm outlines a systematic approach for the assessment and treatment of UTIs from the evaluation of clinical signs and symptoms, specimen collection, and treatment, the primary objective of this quality review was to assess the use of urinalysis in the diagnostic assessment of patients for UTIs.

## METHODS

The quality review was undertaken in PRGH, a small community hospital located on the northern Sunshine Coast of British Columbia. PRGH is a 33-bed facility that supports a regional district with an estimated population size of 20,000. The annual number of admissions for 2015-2016 was 1,599, with an average length of stay of 6.89 patient days. Older adults ( $\geq 65$  years) accounted for 819 (51%) of the admissions, with an average length of stay of 8.7 patient days.

### Design

The review involved two separate components: a staff educational survey and a retrospective analysis of electronic microbiology data.

Ethical review was not required for this quality assurance project as it involved evaluation of expected knowledge and practice.

### Staff survey

The objective of the staff survey was to evaluate the interdisciplinary team's knowledge of urine specimen collection and interpretation of results, which would identify opportunities for improvement. The survey was distributed during the week of March 13 to March 17, 2017 to 19 staff members working day shifts on the medical/surgical unit. 17 of the 19 staff were nurses (LPN/RN), one staff member was an occupational therapist, and one was a physiotherapist. Completed surveys were returned and results were entered into an Excel spreadsheet.

The survey consisted of 16 items based on the VCH Professional Practice guidelines *Indwelling urinary catheter: Guideline to prevent catheter associated urinary tract infections (CAUTI) – Adult* and the ASPIRES algorithm *VCH Management of Urinary Tract Infections in Non-pregnant Adults*. Survey items 1 to 7 evaluated sample collection; items 8 to 16 evaluated the interpretation of urine specimen results.

### Electronic data review

Microbiology data, including the date and time of urine culture specimen results, was extracted for all patients admitted to the medical/surgical unit over a one-year period (September 1, 2015 to August 31, 2016). Corresponding urinalysis data was retrieved manually from the local electronic patient care information system.

Data was imported into Statistical Package for Social Sciences (SPSS, version 18) and reviewed for data quality prior to conducting the descriptive and statistical analyses.

### Definitions

For the purpose of analysis, the following case definitions were used.

#### Urinalysis

"Urinalysis test" was defined as a physical, chemical, or microscopic analysis or examination of urine. It determines the levels of LEs and NITs by use of a dipstick and a microscopic examination for white blood cells.

“Positive urinalysis” was defined as a urine specimen resulting in LEs and/or NITs of any value, whereas a “negative urinalysis” was defined as a urine specimen resulting in the absence of LEs (-) and NITs (-).

#### Urine cultures

“Positive urine culture with significance” was defined as a single urine specimen with the isolation of one predominate bacterial strain with a count of > 100 million CFU/L [4, 8]. “Negative urine cultures” included those with no growth, an insignificant quantity of growth, or mixed growth due to probable contaminants.

## RESULTS

### Education survey


All 19 surveys distributed were returned for a response rate of 100%. No survey items were left blank. There was a general consensus overall: staff scored higher on Section B (interpretation of results) at 80.7% than on Section A (specimen collection) at 69.2%, as summarized in Table 1. The difference was statistically significant (Fisher’s exact test,  $p = 0.02$ ). Four questions were answered with less than 54% accuracy and were distributed between both sections. The question that received the lowest number of correct answers was A7 (16%), which measured knowledge on urine collection methods for the catheterized patient.

### Microbiology data

#### Sample characteristics

318 urine specimens were collected from September 1, 2015 to August 31, 2016 and submitted to the VGH microbiology laboratory for culturing, as noted in Figure 2. 148 (46.5%) specimens were collected from males and 170 (53.5%) specimens were collected from females. The mean age was 73.6 years, with a median of 77 years and a range of 18 to 101 years. Statistical analysis using a non-parametric median test showed that the difference in the median age for males (78.5) vs females (77.0) was not statistically significant ( $p = 0.742$ ).

As noted in Table 2, 62 (19.5%) of the 318 specimens did not have a corresponding urinalysis. Of the 256 urinalysis specimens processed, 136 (53.1%) were negative for both LEs and NITs,



and 120 (46.9%) were positive by definition. 78 (24.5%) of the 318 urine cultures met microbiologic criteria for significant growth of organisms and were considered to be positive urine cultures; of these, 50 (64.1%) corresponded to a positive urinalysis, in comparison to nine (11.5%) corresponding to negative urinalysis. The remaining 19 (24.5%) positive cultures were associated with an absent or missing urinalysis.

184 (57.9%) of the 318 specimens resulted in no growth, of which 45 (24.4%) were associated with a positive urinalysis and 105 (57.1%) were associated with a negative urinalysis. The remaining 34 (18.5%) specimens, which had no growth, were associated with a missing urinalysis. 56 (17.6%) of the 318 cultures submitted did not grow organisms with numbers greater than 100 million CFU/L and were not considered to be positive based on our case definition.

There was a highly statistically significant relationship between urinalysis and clinical significance (Fisher’s exact test  $p < 0.0001$ ). Specifically, a positive urine culture (> 100 million CFU/L) was statistically significantly more likely to be observed when there was a positive urinalysis, and a negative culture (< 100 million CFU/L and No growth) was statistically significantly more likely with a negative urinalysis.


**TABLE 1: Educational survey.**

Question	Correct	Percentage
A1 Elderly population and risk factors for developing a UTI.	17	89%
A2 Change in cognitive function – indication of a UTI.	18	95%
A3 Malodorous/cloudy urine – indication of a UTI.	9	47%
A4 Signs and symptoms of a UTI.	17	89%
A5 Collection of a urinalysis specimen.	14	74%
A6 Ideal urine specimen for culture.	14	74%
A7 Urine specimen from a catheterized patient.	3	16%
<b>Section A subtotal</b>		<b>69.1%</b>
B8 Urinalysis results and interpretation.	18	95%
B9 UTI vs asymptomatic bacteriuria.	15	79%
B10 Definition of a positive urine culture.	10	53%
B11 Prevalence of bacteriuria in elderly patients.	17	89%
B12 Treatment of a positive culture in asymptomatic patients.	15	79%
B13 Asymptomatic bacteriuria vs a UTI.	16	84%
B14 Treatment of asymptomatic bacteriuria.	18	95%
B15 Location of urinalysis results.	19	100%
B16 Location of urine cultures results.	10	53%
<b>Section B subtotal</b>		<b>80.8%</b>

**TABLE 2: Culture significance vs urinalysis.**

Urinalysis		Culture Significance			Total
		Greater than 100 million CFU/L	Less than 100 million CFU/L	No growth	
Not collected	Count	19	9	34	62
	Column %	24.4%	16.1%	18.5%	
(-) Leukocyte and (-) Nitrate	Count	9	22	105	136
	Column %	11.5%	39.3%	57.1%	
(+) Leukocyte and/or (+) Nitrate	Count	50	25	45	120
	Column %	64.1%	44.6%	24.4%	
Total	Count	78	56	184	318

The organisms isolated in urine cultures with significance greater than 100 million CFU/L are summarized in Figure 3. The predominant individual organisms identified were *Escherichia coli* (n = 23; 29.5%), and *Enterococcus faecalis* (n = 13; 16.7%). Mixed organisms contributed the second-highest percentage (21.8%).

**FIGURE 3: Distribution by organism.**

## DISCUSSION

The ASPIRES algorithm provides a systematic approach in the assessment, diagnosis, and treatment of UTIs. Findings from this quality assurance review reinforce the importance of adherence to evidence-driven best practices to ensure consistently high-quality care for our patients. Results revealed areas where concordance with the guidelines needs to be improved.

Firstly, the practical and effective use of a urinalysis specimen in combination with a urine culture in the assessment and diagnosis of a UTI needs to be recognized. Evidence shows that specimens resulting in a negative urinalysis are unlikely to be associated with a positive culture (less than 20%) [7, 10, 11].

In this review, 62 (19.5%) of the 318 cultures did not have a corresponding urinalysis. Of the 256 urinalysis specimens collected, 136 (53.1%) resulted in a negative urinalysis and of

those, 105 (77.2%) were associated with no growth. These results correspond to what is reported in the literature and support the intent of the ASPIRES algorithm to reduce unnecessary testing and treatment of urine cultures, which can lead to significant side effects and expense and can drive antibiotic resistance [2, 6, 10].

As with many rural hospitals affected by the centralization of services, the local microbiology laboratory has been closed, impacting the processing of urine specimens. In response, two separate workflows have evolved to manage urine specimens. All urine for culture and sensitivity is sent to the regional laboratory irrespective of an absent or negative urinalysis. This workflow limits the ability of front-line staff to be compliant with the algorithm and results in an increased number of urine specimens sent for culturing.

Secondly, opportunities for improvement in staff knowledge and practice specific to specimen collection became apparent with the high proportion of cultures that resulted in mixed organisms, which is suggestive of contamination. In this review, mixed organisms accounted for 22% of all organisms with a culture of significance. This was second to *E. coli*, which was the most predominant organism. "Mixed growth" is defined as the presence of more than one bacterium in the urine sample [8, 9]. Cultures that contain more than one organism are usually considered contaminated. Contamination occurs when samples are not collected properly or when there is a substantial delay in processing. Mixed organisms do not meet the clinical definition of a positive culture and therefore impede diagnosis, resulting in a delay of treatment and/or inappropriate antimicrobial use [8, 9, 12].

The challenges of midstream urine collections without contamination are greatly underestimated. Fecal and urinary incontinence have an impact on specimen collection and may play a significant role in contamination. Further investigation is required to assess urine specimen collection techniques, as it is unclear if staff were able to provide adequate instructions to patients in the correct methods of proper cleansing prior to obtaining a specimen.

The results of the staff survey provided insight into the nurses' knowledge of sample collection in the presence of an indwelling catheter. Only 16% of the staff surveyed recognized the need to replace the catheter prior to obtaining urine for culturing. This may have contributed to the large proportion of specimens that resulted in mixed organisms.

Another contributing factor that may influence contamination is the extended period of time between collection of the specimen on the unit and the processing of the urine culture in the regional laboratory.

Lastly, the diagnosis of symptomatic UTIs in older adults is challenging, as most accepted definitions require the presence of localized genitourinary symptoms. However, underlying medical comorbidities such as dementia and stroke may impair peoples' ability to communicate symptoms. As a result, clinicians rely on nonspecific symptoms such as a change in behaviour and laboratory data for diagnosis [2, 6, 13, 14].

A disproportionate number of urine cultures ( $n = 251$ ; 78.9%) were collected from adults ( $\geq 65$  years), suggesting that older adults were more likely to be investigated for possible UTIs. This may be attributed to both the increased risk of UTI for older adults as well as the challenges associated with diagnosis [2, 14-16]. As a chart review was not completed as part of the review, it is unclear if urine specimen collection was influenced by a change in patient behaviour or based on signs and symptoms. The survey suggests opportunity for education in the clinical assessment of a UTI, as only 47% of the participants answered the question pertaining to malodorous and cloudy urine correctly.

This quality review had limitations. Firstly, chart reviews were not conducted to verify the signs and symptoms associated with the urine specimen collection, nor did the review investigate the appropriateness of antimicrobial therapy. Urine specimens obtained from catheterized patients were not distinguished from non-catheterized patients, as this is not reliably documented in the electronic patient care information system.

In summary, older adults represent a large and growing population of hospitalized patients who are at high risk of complications during their stay. UTIs have been identified as the single most common healthcare-acquired infection [1, 2, 16, 17]. Studies have shown that the diagnosis of a symptomatic UTI is challenging due to chronic genitourinary symptoms and cognitive deficits among elderly patients. Urinalysis specimens can provide critical information for interpretation of results in conjunction with urine for culturing.

In the absence of a standardized approach, there is an increased risk of over-collection of samples and treatment of asymptomatic patients, which may lead to a variety of negative consequences, including the development of multidrug-resistant organisms [2].

## REFERENCES

- Nicolle, L. E. (2008). The prevention of hospital-acquired urinary tract infection. *Clinical Infectious Disease*, 46(2) 251-253. doi: 10.1086/524663
- Silver, S. A., Baillie, L., & Simor, A. E. (2009). Positive urine cultures: A major cause of inappropriate antimicrobial use in hospitals? *Canadian Journal of Infectious Diseases Medical Microbiology*, 20(4), 107-111. Retrieved from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2807252/>
- Foxman, B. (2002). Epidemiology of urinary tract infections: Incidence, morbidity, and economic cost. *Disease-a-Month*, 49(2), 53-70. doi: <https://doi.org/10.1067/mda.2003.7>
- Vancouver Coastal Health. (2013). *ASPIRES urinary tract infection algorithm*. Retrieved from [http://vhpharmsci.com/Resouces/Pocket/Anti-microbial-Stewardship-Programme-ASPIRES-UTI-Education-Slides%20\(Feb%202013\).pdf](http://vhpharmsci.com/Resouces/Pocket/Anti-microbial-Stewardship-Programme-ASPIRES-UTI-Education-Slides%20(Feb%202013).pdf)
- Soliman, Y., Meyer, R., & Baum, N. (2016). Falls in the elderly secondary to urinary symptoms. *Reviews in Urology*, 18(1), 28-32. Retrieved from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4859925/>
- Rowe, T. A., & Juthani-Mehta, M. (2013). Urinary tract infection in older adults. *Aging Health*, 9(5). doi: 10.2217/ahe.13.38
- Vancouver Coastal Health (VCH). (2017). *VCH Management of urinary tract infections (UTI) in Non-pregnant Adults*. Retrieved from [http://www.vhpharmsci.com/Resouces/Pocket/VCH%20UTI%20Algorithm%20\\_24OCT15\\_%20\\_FINAL\\_.pdf](http://www.vhpharmsci.com/Resouces/Pocket/VCH%20UTI%20Algorithm%20_24OCT15_%20_FINAL_.pdf)
- Vancouver Coastal Health (VCH). (2017). *Indwelling urinary catheter: Guideline to prevent catheter associated urinary tract infections (CAUTI) – Adult*. Retrieved from <http://shop.healthcarebc.ca/vch/VCHDSTs/D-00-07-30110.pdf>
- Goering, R., Dockrell, H., Zuckerman, M., Wakelin, D., Roitt, I., Mims, C., & Chiodini, P. (2008). *Mims' medical microbiology* (4th ed.). Philadelphia, PA: Elsevier.
- Meyrier, A. (2015, July). *Sampling and evaluation of voided urine in the diagnosis of urinary tract infection in adults*. Retrieved from [http://www.uptodate.com/contents/sampling-and-evaluation-of-voided-urine-in-the-diagnosis-of-urinary-tract-infection-in-adult?source=search\\_result&search=UTI+and+urine+sampling&selectTitle=1%7E150](http://www.uptodate.com/contents/sampling-and-evaluation-of-voided-urine-in-the-diagnosis-of-urinary-tract-infection-in-adult?source=search_result&search=UTI+and+urine+sampling&selectTitle=1%7E150)
- Ouslander, J. G., Schapira, M., Fingold, S., & Schnelle, J. (1995). Accuracy of rapid urine screen tests among incontinent nursing home residents with asymptomatic bacteria. *Journal of the American Geriatrics Society*, 43(7), 772-775.
- Bekeris, L. G., Jones, B. A., Walsh, M. K., & Wagar, E. A. (2008). Urine culture contamination: A College of American Pathologists Q-Probes study of 127 laboratories. *Archives of Pathology & Laboratory Medicine*, 132(6), 913-917. doi: 10.1043/1543-2165(2008)132[913:UCCACO]2.0.CO;2
- Detweiler, K., Mayers, D., & Fletcher, S. (2015). Bacteriuria and urinary tract infections in the elderly. *Urologic Clinics of North America*, 42(4), 561-568. doi: 10.1016/j.ucl.2015.07.002
- Daley, P., Penney, C., Wakeham, S., Compton, G., McKim, A., O'Keefe, J., Barrett, B., & Nicolle, L. (2015). Urinary tract infection diagnosis and response to therapy in long-term care: A prospective observational study. *Canadian Journal of Infectious Diseases and Medical Microbiology*, 26(3), 133-136. Retrieved from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4507838/>
- Fekete, T., & Hooton, T. M. (2016, August). *Approach to the adult with asymptomatic bacteriuria*. Retrieved from <https://www.uptodate.com/contents/approach-to-the-adult-with-asymptomatic-bacteriuria>
- Rowe, T. A., & Juthani-Mehta, M. (2014). Diagnosis and management of urinary tract infection in older adults. *Infectious Disease Clinics of North America*, 28(1), 75-89. doi: 10.1016/j.idc.2013.10.004
- Nicolle, L. E. (2016). Urinary tract infections in the older adult. *Clinics in Geriatric Medicine*, 32(3), 523-538. doi: 10.1016/j.cger.2016.03.002