

**Pan-Canadian Healthcare Infection Surveillance Definition Standardization Project
Long Term Care Infection Surveillance Definitions**

In partnership with:

Accreditation Canada
 Association of Medical Microbiology and Infectious Diseases Canada
 Canadian Patient Safety Institute
 Centre for Communicable Disease and Infection Control, Public Health Agency of Canada
 Infection Prevention and Control Canada

WORKING COMMITTEE

Alisa Cuff	Faith Stoll (Co-chair)	Marilyn Weinmaster
Betty Anne Elford	Gwen Cerkowniak	Monica Sephton
Blair Ranns	Jean Clark	Nathalie Pigeon
Blanda Chow	Jennifer Happe (Co-chair)	Paula Stagg
Caroline Meguerditchian	Karen Cargill	Smit Patel
Darlene Fawcett	Laurel Biluk	Sukhpreet Jagpal
Daphne Murray	Lesley McLeod	Yvette Gable

The Canadian Patient Safety Institute and the Public Health Agency of Canada (PHAC) hosted a national infection prevention and control (IPAC) summit in November 2014. Over 40 participants came together with the goal of advancing IPAC practices and reducing healthcare-associated infections in Canada. At this meeting, measurement and surveillance, specifically improving consistency in surveillance practices across the country, surfaced as a key theme and an action plan was created. Under the leadership of Infection Prevention and Control Canada (IPAC Canada) and the Association of Medical Microbiology and Infectious Diseases Canada (AMMI Canada) a committee was created to help establish and implement standard healthcare infection surveillance definitions for healthcare associated infections (HAI) in acute care and long term care (LTC).

Members of IPAC's Surveillance and Epidemiology (SAEIG) and Long Term Care (LTC) Interest Groups and the L'Association des infirmières en prévention des infections (APII) formed a working group to revise the existing Stone et al. (2012) LTC infection surveillance definitions based on the Canadian healthcare system and an increase in evidence-based literature about some types of infection in the elderly in LTC settings. The Centers for Disease Control (CDC) Healthcare Infection Control Practices Advisory Committee (HICPAC) guideline development methodology (2013) was used to revise the definitions.

TABLE 1: Definitions for Constitutional Criteria
<p>A. Fever</p> <ol style="list-style-type: none"> 1. Single oral temperature >37.8°C OR 2. Repeated oral temperatures >37.2°C or rectal temperatures >37.5°C OR 3. Single temperature >1.1°C over baseline from any site (oral, tympanic, auxiliary) <p>B. Leukocytosis > 10 x 10⁹ leukocytes/L</p> <p>C. Acute change in mental status from baseline (all criteria must be present; see Table 2)</p> <ol style="list-style-type: none"> 1. Acute onset 2. Fluctuating course 3. Inattention 4. Either disorganized thinking or altered level of consciousness <p>D. Acute functional decline</p> <p>A new 3-point increase in total activities of daily living (ADL) score (range, 0–28) from baseline, based on the following 7 ADL items, each scored from 0 (independent) to 4 (total dependence)¹⁴</p> <ol style="list-style-type: none"> 1. Bed mobility 2. Transfer 3. Locomotion within LTC facility 4. Dressing 5. Toilet use 6. Personal hygiene

7. Eating

TABLE 2. Confusion Assessment Method Criteria	
NOTE. Criteria must be assessed during a formal interview with the client. Criteria are adapted from studies by Inouye <i>et al.</i> (1990) and Lim and MacFarlane (2001)	
Criteria	Comments
Acute onset	Evidence of acute change in resident’s mental status from baseline
Fluctuating	Behavior fluctuating (e.g., coming and going or changing in severity during the assessment)
Inattention	Resident has difficulty focusing attention (e.g., unable to keep track of discussion or easily distracted)
Disorganized thinking	Resident’s thinking is incoherent (e.g., rambling conversation, unclear flow of ideas, unpredictable switches in subject)
Altered level of consciousness	Resident’s level of consciousness is described as different from baseline (e.g., hyperalert, sleepy, drowsy, difficult to arouse, nonresponsive)

TABLE 3. Surveillance Definitions for Respiratory Tract Infections (RTIs)	
NOTE. Epidemiological confirmation, instead of a laboratory positive specimen, can be used to meet case definition criteria. A case is considered epidemiologically confirmed by direct contact to a laboratory-confirmed case through person-to-person transmission or through a common exposure (e.g. food)	
Criteria	Comments
<p>A. Common cold syndrome or pharyngitis (at least 2 criteria must be present)</p> <ol style="list-style-type: none"> 1. Runny nose or sneezing 2. Stuffy nose (i.e., congestion) 3. Sore throat or hoarseness or difficulty in swallowing 4. Dry cough 5. Swollen or tender glands in the neck (cervical lymphadenopathy) 6. N/P swab positive for a respiratory pathogen 	Fever may or may not be present. Symptoms must be new and not attributable to allergies.
<p>B. Influenza-like illness (criteria 1 and/or 2 must be present, AND 3 or 4)</p> <ol style="list-style-type: none"> 1. Fever 2. New and or increased cough 3. At least 2 of the following influenza-like illness subcriteria <ol style="list-style-type: none"> a. Chills b. New headache or eye pain c. Myalgias or body aches d. Malaise or loss of appetite e. Sore throat f. Arthralgia (joint pain) 4. N/P swab positive for Influenza virus 	Fever may not be present in the elderly. If criteria for influenza-like illness and another upper or lower RTI are met at the same time, only the diagnosis of influenza-like illness should be recorded. Because of increasing uncertainty surrounding the timing of the start of influenza season, the peak of influenza activity, and the length of the season, “seasonality” is no longer a criterion to define influenza-like illness.
<p>C. Pneumonia (criteria 1 and 2 must be present, OR criteria 1 and 3)</p> <ol style="list-style-type: none"> 1. Interpretation of a chest radiograph as demonstrating pneumonia or the presence of a new infiltrate 2. At least 1 of the following respiratory subcriteria <ol style="list-style-type: none"> a. New or increased cough 	For both pneumonia and lower RTI, the presence of underlying conditions that could mimic the presentation of a RTI (e.g., congestive heart failure or interstitial lung diseases) should be excluded by a review of clinical records and an assessment of presenting symptoms and signs.

<ul style="list-style-type: none"> b. New or increased sputum production c. O₂ saturation <94% on room air or a reduction in O₂ saturation of >3% from baseline d. New or changed lung examination abnormalities e. Pleuritic chest pain f. Respiratory rate of ≥25 breaths/min <p>3. At least 1 constitutional criteria (see Table 1)</p>	
<p>D. Lower respiratory tract infection (bronchitis or tracheobronchitis; all 3 criteria must be present)</p> <ul style="list-style-type: none"> 1. Chest radiograph not performed or negative results for pneumonia or new infiltrate 2. At least 2 of the respiratory subcriteria (a–f) listed in section C above 3. At least 1 of the constitutional criteria (see Table 1) 	(See comment for section C above.)

TABLE 4. Surveillance Definitions for Urinary Tract Infections (UTIs)	
NOTE. A urinalysis negative for leukocytes effectively rules out a UTI while a urinalysis positive for leukocytes does not differentiate symptomatic UTI from asymptomatic bacteriuria. cfu, colony-forming units.	
Criteria	Comments
<p>A. For residents without an indwelling catheter (criteria 1 and 2 must be present with no other identified source of infection, OR criteria 2 and 3)</p> <ul style="list-style-type: none"> 1. At least 1 of the following sign or symptom subcriteria <ul style="list-style-type: none"> 1. Acute pain, swelling, or tenderness of the testes, epididymis, or prostate, in males 2. Fever or leukocytosis (see Table 1) and at least 1 of the following localizing urinary tract subcriteria <ul style="list-style-type: none"> a. Acute dysuria b. Acute costovertebral angle pain or tenderness c. Suprapubic pain d. Gross hematuria e. New or marked increase in incontinence f. New or marked increase in urgency g. New or marked increase in frequency 3. In the absence of fever or leukocytosis, then 2 or more of the following localizing urinary tract subcriteria <ul style="list-style-type: none"> a. Acute dysuria b. Suprapubic pain c. Gross hematuria d. New or marked increase in incontinence e. New or marked increase in urgency f. New or marked increase in 	<p>UTI should be diagnosed when there are localizing genitourinary signs and symptoms and a positive urine culture result. A diagnosis of UTI can be made without localizing symptoms if a blood culture isolate is the same as the organism isolated from the urine and there is no alternate site of infection. In the absence of a clear alternate source of infection, fever or rigors with a positive urine culture result in the noncatheterized resident or acute confusion in the catheterized resident will often be treated as UTI. However, evidence suggests that most of these episodes are likely not due to infection of a urinary source.</p>

frequency	
2. 10^8 cfu/L of no more than 2 species of microorganisms from a specimen collected by an in and out catheter, or alternately a midstream urine	Urine specimens for culture should be processed as soon as possible, preferably within 1–2 h. If urine specimens cannot be processed within 30 min of collection, they should be refrigerated. Refrigerated specimens should be cultured within 24 h. In and out catheter collection is the gold standard for urine collection.
3. A blood culture isolate is the same as the organism isolated from the urine and there is no alternate site of infection	
B. For residents with an indwelling in a single catheter urine specimen or in a midstream voided urine specimen from a resident whose catheter has been removed within the previous 48 h (criteria 1 and 2 must be present with no other identified source of infection, OR criteria 2 and 3) 1. At least 1 of the following sign or symptom subcriteria a. Fever, rigors, or new-onset hypotension, with no alternate site of infection b. Either acute change in mental status or acute functional decline, with no alternate diagnosis and leukocytosis c. New-onset suprapubic pain or costovertebral angle pain or tenderness d. Purulent discharge from around the catheter or acute pain, swelling, or tenderness of the testes, epididymis, or prostate in males	Recent catheter trauma, catheter obstruction, or new onset hematuria are useful localizing signs that are consistent with UTI but are not necessary for diagnosis.
2. Urinary catheter specimen culture with $\geq 10^8$ cfu/L of any organism(s)	Urinary catheter specimens for culture should be collected following replacement of the catheter (if current catheter has been in place for >14 d).
3. A blood culture isolate is the same species as the organism isolated from the urine, with the same resistance pattern, and there is no alternate site of infection	

TABLE 5. Surveillance Definitions for Skin, Soft Tissue, and Mucosal Infections	
NOTE. For wound infections related to surgical procedures, LTC facilities should use the Centers for Disease Control and Prevention’s National Healthcare Safety Network Surgical Site Infection criteria and report these infections back to the institution where the original surgery was performed.	
Criteria	Comments
A. Cellulitis, soft tissue, or wound infection (at least 1 of the following criteria must be present) 1. Pus present at a wound, skin, or soft tissue site 2. New or increasing presence of at least 4 of the following sign or symptom subcriteria a. Heat at the affected site b. Redness at the affected site c. Swelling at the affected site d. Tenderness or pain at the affected site e. Serous drainage at the affected site f. One constitutional criterion (see Table 1) 3. Non-commensal organism isolated with 1 or more signs or symptoms from criteria 2	Presence of organisms cultured from the surface (e.g., superficial swab sample) of a wound is not sufficient evidence that the wound is infected. More than 1 resident with streptococcal skin infection from the same serogroup (e.g., A, B, C, G) in a long-term care facility (LTCF) may indicate an outbreak. Common commensal organisms include diphtheroids (<i>Corynebacterium spp.</i> not <i>C. diphtheria</i>), <i>Bacillus spp.</i> [not <i>B. anthracis</i>], <i>Propionibacterium spp.</i> , coagulase-negative staphylococci (including <i>S. epidermidis</i>), viridans group streptococci, <i>Aerococcus spp.</i> , and <i>Micrococcus spp.</i>

<p>B. Scabies (both criteria 1 and 2 must be present)</p> <ol style="list-style-type: none"> 1. A maculopapular and/or itching rash characteristic of scabies 2. At least 1 of the following scabies subcriteria <ol style="list-style-type: none"> a. Physician diagnosis b. Laboratory confirmation (scraping or biopsy) c. Epidemiologic linkage to a case of scabies with laboratory confirmation 	<p>Consider the appearance and distribution of the rash. The most common symptom of scabies is itching (pruritis) especially at night and pimple (papular) like rash. The itching and rash may affect much of the body or be limited to common sites such as wrists, elbow, armpit, webbing between the fingers, nipple, penis, waist, beltline and buttocks. Tiny burrows that are raised and crooked, grayish white or skin coloured lines on the skin surface. They are found most often in the webbing between the fingers, in the skin folds of the wrist, elbow or knee and on the penis, breast or shoulder blades. If rash presentation is atypical, lab confirmation is recommended.</p> <p>An epidemiologic linkage to a case can be considered if there is evidence of geographic proximity in the facility, temporal relationship to the onset of symptoms, or evidence of common source of exposure (i.e., shared caregiver). Care must be taken to rule out rashes due to skin irritation, allergic reactions, eczema, and other noninfectious skin conditions.</p>
<p>C. Fungal oral or perioral and skin infections</p> <ol style="list-style-type: none"> 1. Oral candidiasis (both criteria a and b must be present) <ol style="list-style-type: none"> a. Presence of raised white patches on inflamed mucosa or plaques on oral mucosa b. Diagnosis by a medical or dental provider 2. Fungal skin infection (both criteria a and b must be present) <ol style="list-style-type: none"> a. Characteristic rash or lesions b. Either a diagnosis by a medical provider or a laboratory confirmed fungal pathogen from a scraping or a medical biopsy 	<p>Mucocutaneous <i>Candida</i> infections are usually due to underlying clinical conditions such as poorly controlled diabetes or severe immunosuppression. Although they are not transmissible infections in the healthcare setting, they can be a marker for increased antibiotic exposure.</p> <p>Dermatophytes have been known to cause occasional infections and rare outbreaks in the LTCF setting.</p>
<p>D. Herpesvirus skin infections</p> <ol style="list-style-type: none"> 1. Herpes simplex infection (both criteria a and b must be present) <ol style="list-style-type: none"> a. A vesicular rash b. Either physician diagnosis or laboratory confirmation 2. Herpes zoster infection (both criteria a and b must be present) <ol style="list-style-type: none"> a. A vesicular rash b. Either physician diagnosis or laboratory confirmation 	<p>Reactivation of herpes simplex (“cold sores”) or herpes zoster (“shingles”) is not considered a healthcare-associated infection. Primary herpesvirus skin infections are very uncommon in a LTCF except in pediatric populations, where it should be considered healthcare associated.</p>
<p>E. Conjunctivitis (at least 1 of the following criteria must be present)</p> <ol style="list-style-type: none"> 1. Pus appearing from 1 or both eyes, present for at least 24 h 2. New or increased conjunctival erythema, with or without itching 3. New or increased conjunctival pain, present for at least 24 h 	<p>Conjunctivitis symptoms (“pink eye”) should not be due to allergic reaction or trauma.</p>

TABLE 6. Surveillance Definitions for Gastrointestinal (GI) Tract Infections	
<p>NOTE. Epidemiological confirmation, instead of a laboratory positive specimen, can be used to meet case definition criteria. A case is considered epidemiologically confirmed by direct contact to a laboratory-confirmed case through person-to-person transmission or through a common exposure (e.g. food)</p>	
Criteria	Comments
<p>A. Gastroenteritis (at least 1 of the following criteria must be present)</p> <ol style="list-style-type: none"> 1. Diarrhea: 3 or more loose or watery stools above what is normal for the resident within a 24-h period 2. Vomiting: 2 or more episodes in a 24-h period 3. Both of the following sign or symptom subcriteria <ol style="list-style-type: none"> a. A stool specimen testing positive for a pathogen (e.g., <i>Salmonella</i>, <i>Shigella</i>, <i>Escherichia coli</i> O157:H7, <i>Campylobacter</i> species, rotavirus) b. At least 1 of the following GI subcriteria <ol style="list-style-type: none"> i. Nausea ii. Vomiting iii. Abdominal pain or tenderness iv. Diarrhea v. mucous in stool 	<p>Care must be taken to exclude noninfectious causes of symptoms. For instance, new medications may cause diarrhea, nausea, or vomiting; initiation of new enteral feeding may be associated with diarrhea; and nausea or vomiting may be associated with gallbladder disease. Presence of new GI symptoms in a single resident may prompt enhanced surveillance for additional cases. In the presence of an outbreak, stool specimens should be sent to confirm the presence of norovirus or other pathogens (e.g., rotavirus or <i>E. coli</i> O157:H7).</p>
<p>B. Norovirus gastroenteritis (both criteria 1 and 2 must be present)</p> <ol style="list-style-type: none"> 1. At least 1 of the following GI subcriteria <ol style="list-style-type: none"> a. Diarrhea: 3 or more loose or watery stools (i.e. Conforming to the shape of the specimen collection container) above what is normal for the resident within a 24-h period b. Vomiting: 2 or more episodes of in a 24-h period 2. A stool specimen for which norovirus is positively detected by electron microscopy, enzyme immunoassay, or molecular diagnostic testing such as polymerase chain reaction (PCR) 	
<p>C. <i>Clostridium difficile</i> infection (both criteria 1 and 2 must be present)</p> <ol style="list-style-type: none"> 1. One of the following GI subcriteria <ol style="list-style-type: none"> a. Diarrhea: 3 or more loose or watery stools (i.e. Conforming to the shape of the specimen collection container) above what is normal for the resident within a 24-h period b. Presence of toxic megacolon (abnormal dilatation of the large bowel, documented radiologically) 2. One of the following diagnostic subcriteria <ol style="list-style-type: none"> a. A stool sample yields a positive laboratory test result for <i>C. difficile</i> toxin A or B, or a toxin-producing <i>C. difficile</i> organism is identified from a stool sample culture or by a molecular diagnostic test such as PCR b. Pseudomembranous colitis is identified during endoscopic examination or surgery 	<p>A “primary episode” of <i>C. difficile</i> infection is defined as one that has occurred without any previous history of <i>C. difficile</i> infection or that has occurred 8 weeks after the onset of a previous episode of <i>C. difficile</i> infection. A “recurrent episode” of <i>C. difficile</i> infection is defined as an episode of <i>C. difficile</i> infection that occurs 8 weeks or sooner after the onset of a previous episode, provided that the symptoms from the earlier (previous) episode have resolved. Individuals previously infected with <i>C. difficile</i> may continue to remain colonized even after symptoms resolve. In the setting of an outbreak of GI infection, individuals could have positive test results for presence of <i>C. difficile</i> toxin because of ongoing colonization and also be co-infected with another pathogen. It is important that other surveillance criteria be used to differentiate infections in this situation.</p>

or in histopathologic examination of a biopsy specimen	
--	--

Table 7. Blood Stream Infections
Adhere to CDC's National Healthcare Safety Network (NHSN) definitions

References

- Agata, E. D., Loeb, M. B., and Mitchell, S. L. (2013). Challenges in assessing nursing home residents with advanced dementia for suspected urinary tract infections. *Journal of the American Geriatrics Society*, 61(1), 62-66.
- Association for Professionals in Infection Control and Epidemiology. (2014) APIC Text of Infection Control and Epidemiology 4th Edition. Washington, DC: Association for Professionals in Infection Control and Epidemiology Inc.
- Bain, B. J., & England, J. M. (1975). Normal haematological values: sex difference in neutrophil count. *British Medical Journal*, 1(5953), 306-309.
- Balogun, S. A., and Philbrick, J. T. (2013). Delirium, a symptom of UTI in the elderly: fact or fable? A systematic review. *Canadian Geriatrics Journal*, 17(1), 22-26.
- Bennett, N. J., Johnson, S. A., Richards, M. J., Smith, M. A., and Worth, L. J. (2016). Infections in Australian Aged-Care Facilities: Evaluating the Impact of Revised McGeer Criteria for Surveillance of Urinary Tract Infections. *Infection Control and Hospital Epidemiology*, 37(5), 610.
- Bernard, H., Höhne, M., Niendorf, S., Altmann, D., and Stark, K. (2014). Epidemiology of norovirus gastroenteritis in Germany 2001–2009: eight seasons of routine surveillance. *Epidemiology and Infection*, 142(01), 63-74.
- Bernatchez, S. F. (2014). Care of peripheral venous catheter sites: advantages of transparent film dressings over tape and gauze. *Journal of the Association for Vascular Access*, 19(4), 256-261.
- British Columbia Provincial Infection Control Network. (2011, February) Respiratory Infection Outbreak Guidelines for Healthcare Facilities. Reference Document for use by Health Care: Reference Document for use by Health Care Organizations for Internal Policy/Protocol Development. Retrieved from http://www.bccdc.ca/resource-gallery/Documents/Guidelines%20and%20Forms/Guidelines%20and%20Manuals/Epid/CD%20Manual/Chapter%203%20-%20IC/RI_Outbreak_Guidelines_March17_Final.pdf
- Canadian Patient Safety Institute Safer Healthcare Now (2015, July) Prevent Ventilator Associated Pneumonia: Getting Started Kit. Retrieved from [http://www.patientsafetyinstitute.ca/en/Topic/Pages/Ventilator-Associated-Pneumonia-\(VAP\).aspx](http://www.patientsafetyinstitute.ca/en/Topic/Pages/Ventilator-Associated-Pneumonia-(VAP).aspx)
- Capobussi, M., Sabatino, G., Donadini, A., Tersalvi, C. A., and Castaldi, S. (2014). Control of scabies outbreaks in an Italian hospital: an information-centered management strategy. *American Journal of Infection Control*, 42(3), 316-320.
- Cardona, A. F., and Wilson, S. E. (2015). Skin and soft-tissue infections: a critical review and the role of telavancin in their treatment. *Clinical Infectious Diseases*, 61(suppl 2), S69-S78.
- Carman, W. F., Elder, A. G., Wallace, L. A., McAulay, K., Walker, A., Murray, G. D., & Stott, D. J. (2000). Effects of influenza vaccination of health-care workers on mortality of elderly people in long-term care: a randomized controlled trial. *The Lancet*, 355(9198), 93-97.
- Centers for Disease Control and Prevention's National Healthcare Safety Network. (2017, January). Pneumonia (Ventilator-associated [VAP] and non-ventilator-associated Pneumonia [PNEU]) Event. Retrieved from <https://www.cdc.gov/nhsn/pdfs/pscmanual/6pscvapcurrent.pdf>
- Chitnis, A. S., Edwards, J. R., Ricks, P. M., Sievert, D. M., Fridkin, S. K., and Gould, C. V. (2012). Device-Associated Infection Rates, Device Utilization, and Antimicrobial Resistance in Long-Term Acute Care Hospitals Reporting to the National Healthcare Safety Network, 2010. *Infection Control and Hospital Epidemiology*, 33(10), 993-1000.
- Chopra, V., Montoya, A., Joshi, D., Becker, C., Brant, A., McGuirk, H., et al. (2015). Peripherally inserted central catheter use in skilled nursing facilities: a pilot study. *Journal of the American Geriatrics Society*, 63(9), 1894-1899.
- Chuang, S., Howley, P. P., and Lin, S. H. (2015). Implementing systems thinking for infection prevention: the cessation of repeated scabies outbreaks in a respiratory care ward. *American journal of infection control*, 43(5), 499-505.
- Cornbleet, P. J. (2002). Clinical utility of the band count. *Clinics in laboratory medicine*, 22(1), 101-136.
- Cornbleet, P. J., & Novak, R. W. (1995). Lack of reproducibility of band neutrophil identification despite the use of uniform identification criteria. *Lab Hematol*, 1(2), 89-96.
- Cotter, M., Donlon, S., Roche, F., Byrne, H., & Fitzpatrick, F. (2012). Healthcare-associated infection in Irish long-term care facilities: results from the First National Prevalence Study. *Journal of Hospital Infection*, 80(3), 212-216.
- Daley, P., Penney, C., Wakeham, S., Compton, G., McKim, A., O'Keefe, J., et al. (2015). Urinary tract infection diagnosis and response to therapy in long-term care: a prospective observational study. *Canadian Journal of Infectious Diseases and Medical Microbiology*, 26(3), 133-136.

- De Wit, M. A. S., Koopmans, M. P. G., Kortbeek, L. M., Wannet, W. J. B., Vinje, J., Van Leusden, F., et al. (2001). Sensor, a population-based cohort study on gastroenteritis in the Netherlands: incidence and etiology. *American Journal of Epidemiology*, 154(7), 666-674.
- Drees, M., Kanapathippillai, N., & Zubrow, M. T. (2012). Bandemia with normal white blood cell counts associated with infection. *The American Journal of Medicine*, 125(11), 1124-e9.
- Earhart, A., and Kaminski, D. (2006). Reducing Catheter-Related Bloodstream Infections: An Extended-Care Facility's Process Improvement Experience. *Journal of the Association for Vascular Access*, 11(2), 90-95.
- Edwards, M., Purpura, J., and Kochvar, G. (2014). Quality improvement intervention reduces episodes of long-term acute care hospital central line-associated infections. *American Journal of Infection Control*, 42(7), 735-738.
- Eikelenboom-Boskamp, A., Cox-Claessens, J. H. M., Boom-Poels, P. G. M., Drabbe, M. I. J., Koopmans, R. T. C. M., and Voss, A. (2011). Three-year prevalence of healthcare-associated infections in Dutch nursing homes. *Journal of Hospital Infection*, 78(1), 59-62.
- Friesema, I. H. M., Vennema, H., Heijne, J. C. M., De Jager, C. M., Morroy, G., Van Den Kerkhof, J. H. T. C., et al. (2009). Norovirus outbreaks in nursing homes: the evaluation of infection control measures. *Epidemiology and Infection*, 137(12), 1722-1733.
- Gaspard, P., Ambert-Balay, K., Mosnier, A., Aho-Glélé, S., Roth, C., Larocca, S., et al. (2015). Burden of gastroenteritis outbreaks: specific epidemiology in a cohort of institutions caring for dependent people. *Journal of Hospital Infection*, 91(1), 19-27.
- Gravel, D., Gardam, M., Taylor, G., Miller, M., Simor, A., McGeer, A., et al. (2009). Infection control practices related to Clostridium difficile infection in acute care hospitals in Canada. *American journal of infection control*, 37(1), 9-14.
- Grima, A., Gatt, A., Zahra, G., and Gambin, A. (2009). Outbreak of norovirus infection in a nursing home for the elderly in Malta, November-December 2008. *Euro surveillance: European Communicable Disease Bulletin*, 14(4), 854-858.
- Han, A., and Jump, R. L. (2014). Discrepancies between surveillance definition and the clinical incidence of Clostridium difficile infection in a Veterans Affairs long-term care facility. *Infection Control and Hospital Epidemiology*, 35(11), 1435-1436.
- Harada, N., Higuchi, K., and Mori, T. (2008). Assessment of nosocomial transmission of tuberculosis in a psychiatric hospital using a whole blood interferon-gamma assay. *Japan Journal of Infectious Disease*, 61(5), 415-418.
- Hayden, M. K., Lin, M. Y., Lolans, K., Weiner, S., Blom, D., Moore, N. M., et al. (2015) Prevention of colonization and infection by Klebsiella pneumoniae carbapenemase-producing enterobacteriaceae in long-term acute-care hospitals. *Clinical Infectious Diseases*, 60(8), 1153-1161.
- Heudorf, U., Boehlcke, K., & Schade, M. (2012). Healthcare-associated infections in long-term care facilities (HALT) in Frankfurt am Main, Germany, January to March 2011. *Euro Surveill*, 17(35), 20256.
- Hewitt, K. A., Nalabanda, A., and Cassell, J. A. (2015). Scabies outbreaks in residential care homes: factors associated with late recognition, burden and impact. A mixed methods study in England. *Epidemiology and Infection*, 143(07), 1542-1551.
- High, K. P., Bradley, S. F., Gravenstein, S., Mehr, D. R., Quagliarello, V. J., Richards, C., & Yoshikawa, T. T. (2009). Clinical practice guideline for the evaluation of fever and infection in older adult residents of long-term care facilities: 2008 update by the Infectious Diseases Society of America. *Journal of the American Geriatrics Society*, 57(3), 375-394.
- Huang, M. Y., Chang, W. H., Hsu, C. Y., Tsai, W., Chen, Y. J., Lee, C. H., et al. (2012). Bloodstream infections in the elderly: effects of nursing homes on antimicrobial-resistant bacteria. *International Journal of Gerontology*, 6(2), 93-100.
- Hunter, J. C., Mu, Y., Dumyati, G. K., Farley, M. M., Winston, L. G., Johnston, H. L., et al. (2016, January). Burden of Nursing Home-Onset Clostridium difficile Infection in the United States: Estimates of Incidence and Patient Outcomes. In *Open forum infectious diseases* (Vol. 3, No. 1, p. ofv196). Oxford University Press.
- Infection Prevention and Control Canada Eastern Ontario Chapter. (2012) Guideline for the Diagnosis and Management of Pneumonia in LTC. Retrieved from www.ipac-canada.org/IPAC-EO/2012 ASP LTC PROTOCOLS.pdf
- Inouye, S. K., Van Dyck, C. H., Alessi, C. A., Balkin, S., Siegal, A. P., & Horwitz, R. I. (1990). Clarifying confusion: the confusion assessment method: a new method for detection of delirium. *Annals of internal medicine*, 113(12), 941-948.
- Institut National de Santé Publique du Québec (2014, April) Définitions pour la surveillance des infections nosocomiales dans les milieux d'hébergement et de soins de longue durée. Retrieved from [https://www.inspq.qc.ca/pdf/publications/1871 Définitions Surveillance Nosocomiales.pdf](https://www.inspq.qc.ca/pdf/publications/1871_Définitions_Surveillance_Nosocomiales.pdf)
- Jungbauer, F. H., Veenstra-Kyuchukova, Y. K., Koeze, J., KruijtSpanjer, M. R., and Kardaun, S. H. (2015). Management of nosocomial scabies, an outbreak of occupational disease. *American Journal of Industrial Medicine*, 58(5), 577-582.
- Khan, A., O'Grady, S., and Muller, M. P. (2012). Rapid control of a scabies outbreak at a tertiary care hospital without ward closure. *American journal of infection control*, 40(5), 451-455.
- Kirk, M. D., Lalor, K., Raupach, J., Combs, B., Stafford, R., Hall, G. V., and Becker, N. (2011). Food-and waterborne disease outbreaks in Australian long-term care facilities, 2001-2008. *Foodborne Pathogens and Disease*, 8(1), 133-139.
- Leduc, A. (2014). Reducing the treatment of asymptomatic bacteriuria in seniors in a long-term care facility. *Canadian Nurse*, 110(7).

- Lim, C. J., McLellan, S. C., Cheng, A. C., Culton, J. M., Parikh, S. N., Peleg, A. Y., and Kong, D. C. (2012). Surveillance of infection burden in residential aged care facilities. *Medical journal of Australia*, 196(5), 327-331.
- Lim, W. S., & Macfarlane, J. (2001). A prospective comparison of nursing home acquired pneumonia with community acquired pneumonia. *European Respiratory Journal*, 18(2), 362-368.
- Loeb, M., Bentley, D. W., Bradley, S., Crossley, K., Garibaldi, R., Gantz, N., et al. (2001). Development of minimum criteria for the initiation of antibiotics in residents of long-term-care facilities: results of a consensus conference. *Infection Control and Hospital Epidemiology*, 22(02), 120-124.
- Lubart, E., Segal, R., Haimov, E., Dan, M., Baumohl, Y., and Leibovitz, A. (2011). Bacteremia in a multilevel geriatric hospital. *Journal of the American Medical Directors Association*, 12(3), 204-207.
- Luxmore, B., Powell, K. R., Díaz, S. R., & Novak, R. W. (2002). Absolute band counts in febrile infants: know your laboratory. *Pediatrics*, 110(1), e12-e12.
- Lynch, D. (2012). Achieving Zero Central Line-Associated Bloodstream Infections: Connector Design Combined with Practice in the Long-Term Acute Care Setting. *Journal of the Association for Vascular Access*, 17(2), 75-77.
- Manitoba Public Health. (2016, August). Communicable Disease Management Protocol: Seasonal Influenza. Retrieved from <http://www.gov.mb.ca/health/publichealth/cdc/protocol/influenza1.pdf>
- McAlearney, A. S., Hefner, J. L., Robbins, J., Harrison, M. I., and Garman, A. (2015). Preventing Central Line-Associated Bloodstream Infections: A Qualitative Study of Management Practices. *Infection Control and Hospital Epidemiology*, 36(05), 557-563.
- McGeer, A., Campbell, B., Emori, T. G., Hierholzer, W. J., Jackson, M. M., Nicolle, L. E., et al. (1991). Definitions of infection for surveillance in long-term care facilities. *American Journal of Infection Control*, 19(1), 1-7.
- Medical Council of Canada. (2017). Objectives for the Qualifying Examination 3rd Ed. 3.3.16. Clinical Laboratory Tests Normal Values. Retrieved from http://apps.mcc.ca/Objectives_Online/objectives.pl?lang=english&loc=values#Urine
- Menezes, F. G. D., Correa, V. M. D. S. D., Franco, F. G. D. M., Ribeiro, M. I., Cardoso, M. F. D. S., Morillo, S. G., et al. (2010). An outbreak of norovirus infection in a long-term care facility in Brazil. *Einstein (São Paulo)*, 8(4), 410-413.
- Micali, G., Lacarrubba, F., Verzi, A. E., and Nasca, M. R. (2014). Low-cost equipment for diagnosis and management of endemic scabies outbreaks in underserved populations. *Clinical Infectious Diseases*, ciu826.
- Munoz-Price, L. S., Hota, B., Stemer, A., and Weinstein, R. A. (2009). Prevention of bloodstream infections by use of daily chlorhexidine baths for patients at a long-term acute care hospital. *Infection Control and Hospital Epidemiology*, 30(11), 1031-1035.
- Mylotte, J. M., Tayara, A., and Goodnough, S. (2002). Epidemiology of bloodstream infection in nursing home residents: evaluation in a large cohort from multiple homes. *Clinical infectious diseases*, 35(12), 1484-1490.
- Nace, D. A., Drinka, P. J., & Crnich, C. J. (2014). Clinical uncertainties in the approach to long term care residents with possible urinary tract infection. *Journal of the American Medical Directors Association*, 15(2), 133-139.
- Nicolle, L. E. (2014). Catheter associated urinary tract infections. *Antimicrobial resistance and infection control*, 3(1), 1.
- O'Horo, J. C., Silva, G. L., Munoz-Price, L. S., and Safdar, N. (2012). The efficacy of daily bathing with chlorhexidine for reducing healthcare-associated bloodstream infections: a meta-analysis. *Infection Control and Hospital Epidemiology*, 33(03), 257-267.
- Ontario Public Health. (2014, December) Infectious Diseases Protocol Appendix B: Provincial Case Definitions for Reportable Diseases, Disease: Influenza. Retrieved from http://www.health.gov.on.ca/en/pro/programs/publichealth/oph_standards/docs/influenza_cd.pdf
- Orfanakis, N. G., Ostlund, R. E., Bishop, C. R., & Athens, J. W. (1970). Normal blood leukocyte concentration values. *American Journal of Clinical Pathology*, 53(5), 647-651.
- Pawar, D., Tsay, R., Nelson, D. S., Elumalai, M. K., Lessa, F. C., McDonald, L. C., and Dumyati, G. (2012). Burden of Clostridium difficile infection in long-term care facilities in Monroe County, New York. *Infection Control and Hospital Epidemiology*, 33(11), 1107-1112.
- Perry, M. (2012). How the signs and symptoms of common infections vary with age. *Practice Nursing*, 23(4), 176-182.
- Pollack, C. V., Amin, A., Ford, W. T., Finley, R., Kaye, K. S., Nguyen, H. H., et al. (2015). Acute bacterial skin and skin structure infections (ABSSI): practice guidelines for management and care transitions in the emergency department and hospital. *The Journal of Emergency Medicine*, 48(4), 508-519.
- Provincial Infectious Disease Advisory Committee. (2013, January). Annex C: Testing, surveillance and management of Clostridium difficile in all healthcare settings. Retrieved from https://www.publichealthontario.ca/en/eRepository/PIDAC-IPC_Annex_C_Testing_SurveillanceManage_C_difficile_2013.pdf
- Public Health Agency of Canada (2011, May) The Canadian Pandemic Plan for the Healthcare Sector, Annex F Prevention and Control of Influenza during Pandemics in all Healthcare Settings. Retrieved from <http://www.phac-aspc.gc.ca/cpip-pclcpi/ann-f-eng.php>

- Public Health Agency of Canada. (2016, November 21). Canadian Pandemic Influenza Preparedness: Planning Guidance for the Health Sector. Retrieved from <http://www.phac-aspc.gc.ca/cpip-pclcpi/index-eng.php>
- Rhee, S. M., and Stone, N. D. (2014). Antimicrobial stewardship in long-term care facilities. *Infectious Disease Clinics of North America*, 28(2), 237-246.
- Rotjanapan, P., Dosa, D., & Thomas, K. S. (2011). Potentially inappropriate treatment of urinary tract infections in two Rhode Island nursing homes. *Archives of Internal Medicine*, 171(5), 438-443.
- Rowe, T. A., and Juthani-Mehta, M. (2013). Urinary tract infection in older adults. *Aging health*, 9(5), 519-528.
- Rowe, T. A., and Juthani-Mehta, M. (2014). Diagnosis and management of urinary tract infection in older adults. *Infectious Disease Clinics of North America*, 28(1), 75-89.
- Sakon, N., Yamazaki, K., Nakata, K., Kanbayashi, D., Yoda, T., Mantani, M., et al.. (2014). Impact of genotype-specific herd immunity on the circulatory dynamism of norovirus: a 10-year longitudinal study of viral acute gastroenteritis. *Journal of Infectious Diseases*, 211(6): 879-888.
- Salgado, C. D., Chinnnes, L., Paczesny, T. H., and Cantey, J. R. (2007). Increased rate of catheter-related bloodstream infection associated with use of a needleless mechanical valve device at a long-term acute care hospital. *Infection Control and Hospital Epidemiology*, 28(06), 684-688.
- Sampathkumar, P., Drage, L. A., and Martin, D. P. (2009, March). Herpes zoster (shingles) and postherpetic neuralgia. In *Mayo Clinic Proceedings* (Vol. 84, No. 3, pp. 274-280). Elsevier.
- Schwartz, S., Vergoulidou, M., Schreier, E., Loddenkemper, C., Reinwald, M., Schmidt-Hieber, M., et al.. (2011). Norovirus gastroenteritis causes severe and lethal complications after chemotherapy and hematopoietic stem cell transplantation. *Blood*, 117(22), 5850-5856.
- Schweon, S. J., Edmonds, S. L., Kirk, J., Rowland, D. Y., & Acosta, C. (2013). Effectiveness of a comprehensive hand hygiene program for reduction of infection rates in a long-term care facility. *American Journal of Infection Control*, 41(1), 39-44.
- Seebach, J. D., Morant, R., Rüegg, R., Seifert, B., & Fehr, J. (1997). The diagnostic value of the neutrophil left shift in predicting inflammatory and infectious disease. *American Journal of Clinical Pathology*, 107(5), 582-591.
- Shapiro, M. F., Hatch, R. L., & Greenfield, S. (1984). Cost containment and labor-intensive tests: the case of the leukocyte differential count. *JAMA*, 252(2), 231-234.
- Shi, Q., Warren, L., Saposnik, G., & MacDermid, J. C. (2013). Confusion assessment method: a systematic review and meta-analysis of diagnostic accuracy. *Neuropsychiatr Dis Treat*, 9, 1359-70.
- Simonetti, A. F., Viasus, D., Garcia-Vidal, C., & Carratalà, J. (2014). Management of community-acquired pneumonia in older adults. *Therapeutic advances in infectious disease*, 2(1), 3-16.
- Stevens, D. L., Bisno, A. L., Chambers, H. F., Dellinger, E. P., Goldstein, E. J., Gorbach, S. L., et al. (2014). Practice guidelines for the diagnosis and management of skin and soft tissue infections: 2014 update by the Infectious Diseases Society of America. *Clinical Infectious Diseases*, 59(2), e10-e52.
- Stone, N. D., Ashraf, M. S., Calder, J., Crnich, C. J., Crossley, K., Drinka, P. J., et al. (2012). Surveillance definitions of infections in long-term care facilities: revisiting the McGeer criteria. *Infection Control and Hospital Epidemiology*, 33(10), 965-977.
- Szabó, R., & Böröcz, K. (2015). Infections and antimicrobial use among institutionalized residents in Hungary: increasing need of microbiological surveillance. *Acta Microbiologica et Immunologica Hungarica*, 62(1), 29-44.
- Tavianini, H. D., Deacon, V., Negrete, J., and Salapka, S. (2014). Up for the challenge: eliminating peripherally inserted central catheter infections in a complex patient population. *Journal of the Association for Vascular Access*, 19(3), 159-164.
- Tebbutt, G. M., Wilson, D., & Holtby, I. (2009). A study of 279 general outbreaks of gastrointestinal infection in the north-East region of England. *International Journal of Environmental Research and Public Health*, 6(2), 547-557.
- Thompson, N. D., and Perz, J. F. (2009). Eliminating the blood: ongoing outbreaks of hepatitis B virus infection and the need for innovative glucose monitoring technologies. *Journal of diabetes science and technology*, 3(2), 283-288.
- Thompson, N. D., Barry, V., Alelis, K., Cui, D., and Perz, J. F. (2010). Evaluation of the potential for bloodborne pathogen transmission associated with diabetes care practices in nursing homes and assisted living facilities, Pinellas County. *Journal of the American Geriatrics Society*, 58(5), 914-918.
- Van Esch, G., Van Broeck, J., Delmée, M., & Catry, B. (2015). Surveillance of *Clostridium difficile* infections in a long-term care psychogeriatric facility: outbreak analysis and policy improvement. *Archives of Public Health*, 73(1), 18.
- Van Gessel, H., Riley, T., & McGregor, A. (2009). *Clostridium difficile* infection: an update for infection control practitioners. *Healthcare Infection*, 14(3), 115-118.
- Vega, E., Barclay, L., Gregoricus, N., Shirley, S. H., Lee, D., & Vinjé, J. (2014). Genotypic and epidemiologic trends of norovirus outbreaks in the United States, 2009 to 2013. *Journal of Clinical Microbiology*, 52(1), 147-155.
- Venkatachalam, I., Yang, H. L., Fisher, D., Lye, D. C., Lin, L. M., Tambyah, P., and Perl, T. M. (2014). Multidrug-resistant gram-negative bloodstream infections among residents of long-term care facilities. *Infection Control and Hospital Epidemiology*, 35(05), 519-526.

- Viale, P., Tumietto, F., Giannella, M., Bartoletti, M., Tedeschi, S., Ambretti, S., et al. (2015). Impact of a hospital-wide multifaceted programme for reducing carbapenem-resistant Enterobacteriaceae infections in a large teaching hospital in northern Italy. *Clinical Microbiology and Infection*, 21(3), 242-247.
- Vigil, H. R., and Hickling, D. R. (2016). Urinary tract infection in the neurogenic bladder. *Translational Andrology and Urology*, 5(1), 72.
- Viseur, N., Lambert, M. L., Delmee, M., Van Broeck, J., & Catry, B. (2011). Nosocomial and non-nosocomial *Clostridium difficile* infections hospitalised patients in Belgium-compulsory surveillance data from 2008 to 2010. *Euro Surveillance*, 16(43):pii=20000.
- Wei, L. A., Fearing, M. A., Sternberg, E. J., & Inouye, S. K. (2008). The Confusion Assessment Method: a systematic review of current usage. *Journal of the American Geriatrics Society*, 56(5), 823-830.
- Wiksw, M. E., Kambhampati, A., Shioda, K., Walsh, K. A., Bowen, A., & Hall, A. J. (2015). Outbreaks of acute gastroenteritis transmitted by person-to-person contact, environmental contamination, and unknown modes of transmission—United States, 2009–2013. *MMWR Surveillance Summer*, 64(12), 1-16.
- World Health Organization (2014, January) Global Epidemiological Surveillance Standards for Influenza. Retrieved from http://www.who.int/influenza/resources/documents/influenza_surveillance_manual/en/
- Wu, F. T., Chen, H. C., Yen, C., Wu, C. Y., Katayama, K., Park, Y., et al (2015). Epidemiology and molecular characteristics of norovirus GII. 4 Sydney outbreaks in Taiwan, January 2012–December 2013. *Journal of Medical Virology*, 87(9), 1462-1470.
- Yen, C., Wiksw, M. E., Lopman, B. A., Vinje, J., Parashar, U. D., & Hall, A. J. (2011). Impact of an emergent norovirus variant in 2009 on norovirus outbreak activity in the United States. *Clinical Infectious Diseases*, 53(6), 568-571.
- Zacharski, L. R., Elveback, L. R., & Linman, J. W. (1971). Leukocyte counts in healthy adults. *American Journal of Clinical Pathology*, 56(2), 148-150.